

**Apoptotic induction induces *Leishmania aethiopica* and *L. mexicana* spreading in terminally differentiated THP-1 cells**

Rajeev Rai<sup>1</sup>, Paul Dyer<sup>1</sup>, Simon Richardson<sup>1</sup>, Laurence Harbige<sup>1</sup>, Giulia Getti<sup>1</sup>

<sup>1</sup>Department of Life and Sport Science, University of Greenwich, Chatham Maritime, Kent, ME4 4TB, United Kingdom. Corresponding email: G.T.M.Getti@greenwich.ac.uk

## SUMMARY

Leishmaniasis develops after parasites establish themselves as amastigotes inside mammalian cells and start replicating. As relatively few parasites survive the innate immune defence, intracellular amastigotes spreading towards uninfected cells is instrumental to disease progression. Nevertheless the mechanism of *Leishmania* dissemination remains unclear, mostly due to the lack of a reliable model of infection spreading. Here, an *in vitro* model representing the dissemination of *Leishmania* amastigotes between human macrophages has been developed. Differentiated THP-1 macrophages were infected with GFP expressing *L. aethiopica* and *L. mexicana*. The percentage of infected cells was enriched via camptothecin treatment to achieve  $64.1\pm 3\%$  (*L. aethiopica*) and  $92\pm 1.2\%$  (*L. mexicana*) at 72 hours, compared to  $35\pm 4.2\%$  (*L. aethiopica*) and  $36.2\pm 2.4\%$  (*L. mexicana*) in untreated population. Infected cells were co-cultured with a newly differentiated population of THP-1 macrophages. Spreading was detected after 12 hours of co-culture. Live cell imaging showed inter-cellular extrusion of *L. aethiopica* and *L. mexicana* to recipient cells took place independently of host cell lysis. Establishment of secondary infection from *Leishmania* infected cells provided an insight into the cellular phenomena of parasite movement between human macrophages. Moreover, it supports further investigation into the molecular mechanisms of parasites spreading, which forms the basis of disease development.

## KEYWORDS

*In vitro* model; spreading; *Leishmania aethiopica*; *Leishmania mexicana*; human monocytes; infection; THP-1, Retinoic acid

## KEY FINDINGS

- Co-culture of enriched populations of infected THP-1 cells and freshly differentiated THP-1 cells were established to mimic late stages of *Leishmania* infection.
- Flow cytometry analysis detected a 50% increase in the percentage of infected cells 24 hours after co-culture.
- Live imaging showed dissemination of *L. aethiopica* and *L. mexicana* between human macrophages via non-lytic mechanisms.

## INTRODUCTION

Unicellular protozoan parasites belonging to the genus *Leishmania* are the causative agent of a spectrum of human diseases known as leishmaniasis. Over 350 million people worldwide are at risk of contracting the disease which is endemic in 98 countries (Alvar et al., 2012). Moreover, current long term conflicts and consequent migration of infected individuals from endemic areas has increased not only the number of reported cases but also the risk of an increase in the number of countries becoming endemic of the disease (Kedzierski, 2010 and Stamm, 2016). Depending on the species of parasite and host genetic background, disease manifestation ranges from asymptomatic to cutaneous skin ulceration to life threatening infection of liver and spleen, leading to death, if untreated. No vaccine is currently available and effectiveness of existent drugs is limited due to toxicity and emergence of resistant parasites, making research on disease development and host pathogen interaction a major priority in public health. The transmission of leishmaniasis is initiated when the sandfly injects the parasite infective stage (metacyclic promastigotes) during a blood meal. The total number of promastigotes inoculated is low (100-1000) (Kimblin et al., 2008 and Rogers et al., 2004) with approximately 90% killed instantaneously by the host complement system despite the protective action of the sandfly saliva (Dominguez et al., 2003 and Gomez and Oliveira, 2012). Disease can only develop in those

cases where the surviving parasites successfully establish themselves within the definitive host cells, the macrophages.

Survival inside host macrophages depends on the ability of well characterised virulence factors (e.g. GIPLs, GP63, LPG, PPG and KMP-11) to inhibit macrophages anti-microbial activity (Bifeld and Clos, 2015). *Leishmania* inhibits multiple host pathways including lysosomal fusion (Lodge and Descoteaux, 2005), inflammatory cytokine and chemokine production (Cameron *et al.*, 2004 and Gregory *et al.*, 2008), and MHC class II expression (De Souza *et al.*, 1995). These virulent functions provide sufficient time for promastigotes to successfully transform into amastigotes, which establish themselves within the parasitophorous vacuole and start to actively replicate. Eventually intracellular parasites are released and spread to uninfected cells. In mice models, a silent phase, lasting 4–5 wk was identified whereby the amplification of parasites in the dermis takes place without the formation of a lesion. The development of a lesion coincides with the killing of the parasites (Belkaid *et al.*, 2000). Although the cellular and molecular mechanism behind *Leishmania* spread remains largely unknown, the above findings indicate a silent spreading takes place. The prime reason for this slow advancement is the lack of an effective *in vitro* model to represent this stage of host-parasite interaction. It has been widely assumed that *Leishmania* could egress via host cell lysis as previously observed for numerous human pathogens such as *Shigella flexneri* and *Listeria monocytogenes* (Ashida *et al.*, 2011). Nevertheless, multidimensional live cell imaging has shown evidence that spreading of *L. (L.) amazonensis* amastigotes between mice macrophages can also take place via a regulated mechanism resembling apoptosis (Real *et al.*, 2014). Interestingly the existence of this mechanism is species-specific with no evidence of apoptotic induction detected during *L. (V.) guyanensis* (DaMata *et al.*, 2015). However, no data is available for other *Leishmania* species or, more importantly, on whether a similar mechanism exists in human macrophages. The choice of host cell to support *Leishmania* infection makes significant

differences to the parasites susceptibility to drugs (Seifert *et al.*, 2010) and it cannot be excluded that it would also impact on the parasites behaviour during spreading. Studies carried out directly on human cells are needed but are complicated by the fact that available models allow study of infection over a relatively short period of times (72 hours maximum) which is generally insufficient for spreading of amastigotes to take place (Getti *et al.*, 2008).

This study aimed to develop an *in vitro* model, which allowed the study of long-term *Leishmania* infection and dissemination in human macrophages. This was successfully achieved by employing a cell-to-cell infection strategy whereby enriched populations of *L. aethiopica* and *L. mexicana* infected cells were used to infect uninfected recipient macrophages. The percentage of infected cells in the mixed population doubled within 12 hours from co-culture without significantly affecting viable cell number, suggesting infection spreading was taking place. Live microscopy confirmed parasite movement between infected and recipient cells and gave an initial insight into the process of *Leishmania* extrusion.

## MATERIALS AND METHODS

### *Parasite culture*

Previously described promastigotes of *Leishmania aethiopica* (MHOM/ET/72/L100) and *Leishmania mexicana* (MNYC/BZ/62/M379) constitutively expressing GFP were used in this study (Patel *et al.*, 2014). The reporter gene, GFP, was integrated in the genome, downstream of the promoter of the 18S rRNA. The presence of several copies of the 18S rRNA gene tandemly arrayed in the *Leishmania* parasite genome is advantageous in that introduction of transgenes into the chromosome has no detrimental effect on parasite survival (Mißlitz *et al.* 2000).

For simplicity, GFP expressing clones were named as: L8G for *L. aethiopica* and M5G for *L. mexicana*. The promastigotes were grown in Schneider's Drosophila Medium (Thermo Fischer Scientific, UK) supplemented with 23% Foetal Calf Serum (FCS) (Thermo Fischer Scientific) and 1× Penicillin-Streptomycin-Glutamine (Thermo Fischer Scientific) at neutral pH. The parasites were sub-cultured once a week at a concentration of  $1 \times 10^6$  per ml, supplemented with 700 µg/ml of G418 (Sigma-Aldrich, UK) and were incubated at 24°C cooling incubator.

#### *Human cell culture*

Human THP-1 monocytes (ATCC, TIB-202) were sub-cultured every 3 days at a concentration of  $2.5 \times 10^5$  per ml in complete RPMI 1640 medium (Thermo Fischer Scientific) and incubated at 37°C in a humidified 5% CO<sub>2</sub> incubator. The medium was made complete following supplementation with 10% FCS and 1× Penicillin-Streptomycin-Glutamine.

#### *Infection of differentiated THP-1 macrophages with GFP expressing Leishmania*

Experimental infection with *L. aethiopica* and *L. mexicana* was carried out as previously described (Patel *et al.*, 2014). Briefly, peanut lectin agglutination was used to isolate the metacyclic promastigotes from stationary phase cultures. This infectious form of the parasite was used to infect differentiated THP-1 macrophages at a ratio of 10:1 (parasites to cell ratio). This differentiation was achieved by treating THP-1 monocytes with 1 µM of retinoic acid (Sigma-Aldrich) for 72 hours prior to infection (Ogunkolade *et al.*, 1990). The infected cells was seeded on a 24 well culture plate and incubated at 37°C in a humidified 5% CO<sub>2</sub> incubator. The percentage of infection for *L. aethiopica* and *L. mexicana* was determined through flow cytometry (BD Bioscience, UK) at 24, 48, 72 and 96 hours.

### *Induction of apoptosis*

Apoptotic induction was carried out by treatment with 3  $\mu\text{M}$  of camptothecin (Sigma-Aldrich) at 0 hours after infection. Excess camptothecin from *L. mexicana* and *L. aethiopica* infected cells were washed away with 1 $\times$  PBS after 5 and 24 hours after infection, respectively. Following the completion of washing step, the infected cells were re-suspended in complete RPMI medium and incubated at 37°C in a humidified 5% CO<sub>2</sub> incubator for 72 hours.

### *Detection of cellular apoptosis*

PE Annexin V assay kit I (BD Biosciences) was used to determine apoptosis. The assay is based on Annexin V protein binding onto the externalized phosphatidyl serine (PS), which is an early marker for apoptotic cells. The kit also contains 7-AAD dye, which intercalates with DNA if the cell membrane is damaged, hence measuring necrosis. Briefly, the harvested cells were centrifuged at 500 g for 10 minutes and washed twice with ice-cold 1 $\times$  PBS. After re-suspension in 1 $\times$  binding buffer at a cell concentration of 1 $\times 10^6$  per ml, 100  $\mu\text{l}$  of cells were then stained with 5  $\mu\text{l}$  of PE Annexin V and 5  $\mu\text{l}$  of 7AAD and incubated for 15 minutes in the dark at room temperature. Finally, 400  $\mu\text{l}$  of 1 $\times$  binding buffer was added to the stained cells and the percentage of Annexin V positive / 7AAD negative (apoptosis) cells were analysed through flow cytometry within 1 hour of staining.

### *Infection of differentiated THP-1 macrophages with Leishmania infected cells*

After 72 hours incubation of camptothecin treated *L. aethiopica* and *L. mexicana* infected cells, the concentration of viable cells was determined via trypan blue stain (Sigma-Aldrich). The infected cells were then centrifuged at 500 g for 10 minutes and the supernatant removed before co-culturing with a freshly prepared batch of differentiated THP-1 macrophages. Three different infection ratios were tested including 10:1, 5:1 and 1:1

(infected cell per differentiated THP-1 macrophage ratio). The percentage of infection was determined after 0, 12, 24 and 48 hours of co-culture through flow cytometry. For this experiment, 72 hours incubated non-treated *L. aethiopica* and *L. mexicana* infected cells co-cultured with differentiated THP-1 macrophages were taken as a negative control.

### *Fluorescent microscopy*

THP-1 macrophage from 48 and 96 hours incubation were centrifuged at 500 g for 5 minutes before washing twice with 1× PBS. Equal volume (20 µl) of suspended cells was added onto microscopic slides before applying a drop of Prolong gold anti-fade mount with DAPI (Thermo Fisher Scientific). The resulting samples were then visualised through a fluorescent microscope (Nikon, UK).

### *Real time lapse microscopy*

A fresh population of differentiated THP-1 macrophages at a concentration of  $1 \times 10^5$  per ml was labelled with 0.5× CellMask™ orange plasma membrane stain (Thermo Fischer Scientific) for 15 minutes at 37°C. After washing three times with 1× PBS, the stained cells were re-suspended in a previously described cell imaging buffer solution (Richardson *et al.*, 2008). After seeding in a 6 well culture plate containing  $1 \times 10^5$  of apoptotic induced *L. aethiopica* and/or *L. mexicana* infected cells, the co-culture sample was then mounted on 37°C heated stage connected to a fluorescent microscope. Using an in built digital camera (Nikon), live images were acquired every 5 minutes with bright field and TRITC filters for a maximum period of 12 hours under 20× objectives (Nikon). After acquisition, images were processed by NIS Elements version 4.2 software, which allowed for the construction of multidimensional time-lapse videos.

### *Flow cytometry*

To determine the percentage of infection, the infected cells were extensively washed with 1x PBS before analysis with flow cytometry. Following an initial gating with Forward scatter (FSC) and Side scatter (SSC) parameters, the THP-1 population was further gated using a histogram plot to separate GFP positive (infected cell) from GFP negative (uninfected cell) population.

### *Statistical Analysis*

Data shown are representative of at least three independent experiments performed in three biological replicates. Data represents mean values with statistical error mean (SEM). Statistical analysis was performed using student *t* test provided by GraphPad Prism (CA, USA) software and p values less than 0.05 ( $P < 0.05$ ) were considered significant differences between two independent samples.

## RESULTS

### *Infection of THP-1 macrophage with Leishmania promastigotes*

Current *in vitro* models of *Leishmania* infection in human cells are based on promastigotes infection of macrophage cell lines such as THP-1 (Getti *et al.*, 2008, Ogunkolade *et al.*, 1990, Hsiao *et al.*, 2011). In order to identify whether such model is suitable to represent parasite spread, differentiated THP-1 macrophages were infected with *L. aethiopica* (L8G) and *L. mexicana* (M5G) promastigotes. The percentage of cells harbouring GFP expressing parasites during early (24 and 48 hours) and late (72 and 96 hours) stages of infection was quantified via flow cytometry.

After 24 hours,  $57.1 \pm 0.4\%$  and  $57 \pm 9.1\%$  of cells were infected with *L. aethiopica* and *L. mexicana*, respectively (Fig. 1A). This level of infection for both species was maintained

till 48 hours of continuous incubation. However, no spreading was observed as, there was a significant decrease ( $P<0.05$ ) in the percentage of *L. aethiopica* and *L. mexicana* infected cells from 48 to 96 hours (Fig. 1A). To clarify the cause of this reduction, cell viability was quantified and showed a two-fold increase in the concentration of viable cells from 48 to 96 hours in both uninfected and infected cells suggesting that the cells had resumed replication (Fig. 1B and C).

#### *Effect of apoptotic induction on the rate of Leishmania infection*

Induction of host cell apoptosis has been reported to promote cell-to-cell dissemination of *L. amazonensis* during mice macrophage infection *in vitro* (Real *et al.*, 2014). Therefore, we investigated whether apoptotic induction could also support *L. aethiopica* and *L. mexicana* spreading within the infection model based on THP-1 macrophage. Camptothecin is a quinoline alkaloid which binds to topoisomerase I and DNA complex, stabilising it and causing DNA damage which results in apoptosis. The optimal concentration and length of camptothecin treatment was identified for each infecting species (data not shown). Based on optimised conditions, *L. aethiopica* and *L. mexicana* infected cells were treated with camptothecin for 24 and 5 hours respectively after infection and the percentage of infected cells was detected over a period of 96 hours.

At 24 hours,  $35.4\pm 2\%$  and  $67.1\pm 5.4\%$  of cells were infected by *L. aethiopica* and *L. mexicana*, respectively, in the presence of camptothecin treatment (Fig. 2A). Interestingly, the treatment caused a significant increase ( $P<0.05$ ) in the percentage of infection for both species, reaching a maximum of  $64.1\pm 3\%$  for *L. aethiopica* and  $92\pm 1.2\%$  for *L. mexicana* within 72 hours before plateauing at 96 hours (Fig. 2A). This indicated that camptothecin treatment did not affect intracellular parasites viability. The concentration of viable cells was similar from 24 to 96 hours infection, but significantly lower than the non-treated samples (Fig. 1B and Fig. 2B).

### *Infection of differentiated THP-1 macrophages with L. aethiopica and L. mexicana infected cells*

Despite the low concentration of viable cells in camptothecin treated *L. aethiopica* and *L. mexicana* infected samples (compared to non-treated), a heavy proportion of infected cells were obtained at 72 and 96 hours. To determine which of the infected culture, 72 or 96 hours, would be suitable as a donor population, the long-term effect of camptothecin treatment on THP-1 cells was determined. Annexin V assay showed no significant difference between the percentage of apoptosis of *L. aethiopica* and *L. mexicana* infected cells with and without 72 hours camptothecin treatment (Supplementary Fig. S1A). In contrast, the treated *L. aethiopica* and *L. mexicana* infected cells displayed a significantly higher percentage of apoptosis than the non-treated culture at 96 hours (Supplementary Fig. S1B). Therefore, to remove any experimental bias due to camptothecin treatment, the 72 hours infected population ( $64.1\pm 3\%$  and  $92\pm 1.2\%$ ) was used as a donor to infect newly differentiated THP-1 macrophages. Different infection ratios were tested (1:1, 5:1 and 10:1 infected to uninfected cell ratio) and the course of infection was detected at 0, 12, 24 and 48 hours after co-culture through flow cytometry.

Immediately (0 hour) after co-culture between infected and uninfected cells, a dilution effect on the percentage of infection for both *L. aethiopica* and *L. mexicana* was detected as expected (Fig. 3A and C). Following 12 hours of co-culture, the percentage of *L. aethiopica* infected cells almost doubled from  $35\pm 2.3$  to  $60\pm 2.1\%$  for 1:1 and significantly increased ( $P < 0.05$ ) for 5:1 ( $54\pm 2.4$  to  $69\pm 3\%$ ) and 10:1 ( $58\pm 4.1$  to  $74\pm 0.4\%$ ) infection ratios (Fig. 3B). This was also observed during co-culture with *L. mexicana* infected cells;  $51\pm 2.3$  to  $76\pm 5.2\%$  for 1:1,  $71\pm 3.3$  to  $88.2\pm 1\%$  for 5:1 and  $80\pm 2.1$  to  $90\pm 2.3\%$  for 10:1 infection ratios (Fig. 3D). However, such effects were not detected in the negative control, comprising of non-treated infected cells co-cultured with uninfected cells. The trypan blue assay showed

no significant difference in the concentration of viable cells between 0 and 12 hours of co-culture (Fig. 4 A and B). This suggested that the infection increase from 0 to 12 hours time frame was due to spreading of *L. aethiopica* and *L. mexicana* amastigotes from the 72 hours donor infected cells to healthy cells and not caused by the reduction of uninfected cells. No further increase in the percentage of *L. aethiopica* and *L. mexicana* infected cells was detected from 12 to 24 hours after co-culture (Fig. 3B and D).

Overall, the above results provided quantitative evidence supporting the hypothesis that *L. aethiopica* and *L. mexicana* amastigotes spreading takes place from 0 to 12 hours after co-culture. This *in vitro* spreading model provides an initial platform to address which mode of exit is utilised by both species of *Leishmania* in order to transfer from cell-to-cell.

#### *Live cell imaging of intracellular L. aethiopica and L. mexicana amastigotes spreading*

During cell-to-cell infection spreading, intracellular pathogens within the donor cells can trigger either a lytic or a non-lytic release (Friedrich *et al.*, 2012). To verify whether such a process might occur during *L. aethiopica* and *L. mexicana* release, time lapse microscopy was performed.

Initially, differentiated THP-1 macrophages (pre-stained with CellMask™ orange plasma membrane stain) were co-cultured with *L. aethiopica* and/or *L. mexicana* infected cells and imaged over 12 hours. From 8 hours, *L. aethiopica* amastigotes started migrating towards the peripheral membrane of the donor cells, before completely transferring into the already infected recipient stained cells within 10 minutes of recording (Fig. 5A, Supplementary Video 1). This phenomenon was also visualized in *L. mexicana* infected co-culture whereby after 3 hours, an amastigote appeared to be extruded into the recipient stained cells (Fig. 5B, Supplementary Video 2). Since donor cell (in both scenarios) remained intact during amastigote transfer, it is evident that *Leishmania* spread can take

place via a non-lytic mode in human cells. It remains unclear whether recipient macrophages selectively engulf *L. aethiopica* and *L. mexicana* via recognition of parasite or host associated markers present in the donor cell.

## DISCUSSION

The development of leishmaniasis and associated pathogenesis is dependent on *Leishmania* amastigotes ability to successfully disseminate to uninfected macrophages. Despite the importance of this virulence mechanism on the outcome of infection, a lack of a reliable infection model representative of parasite-host interaction has led to a major gap in knowledge pertaining to the understanding of *Leishmania* spreading in human macrophages. Here, we have reported the development of a co-culture based experiment, which models cell-to-cell dissemination of *L. aethiopica* and *L. mexicana* parasites. The spreading of parasites was evaluated by monitoring the total percentage of infection as well as cell viability before and after co-culture. This approach, along with live imaging, not only confirmed spreading but also supported a previous observation that some *Leishmania* species can exit host cells in a non-lytic process.

During natural *Leishmania* infection, macrophages are the definitive human cells for parasite growth and replication. To model this host-parasite interaction *in vitro*, many investigators have used either macrophage-like cell lines or human blood derived macrophages (Ruhland *et al.*, 2007, Ogunkolade *et al.*, 1990, Lisi *et al.*, 2005). Since primary blood derived macrophages do not support replication and spreading of amastigotes (Hsiao *et al.*, 2011), cell lines were selected in our *in vitro* model. The choice of human THP-1 is further supported by the fact that they are non-adherent, which makes them ideal for flow cytometry studies. The use of adherent cell lines requires detachment through trypsin treatment that might damage cell membranes causing subsequent parasite release, leading to inconsistent data (Huang *et al.*, 2010). Direct infection of differentiated THP-1

macrophages with promastigotes was not sufficient to detect spreading as the effect of retinoic acid started to subside after 48 hours. At this point, THP-1 replication resumed causing the percentage of infected cells to decrease, eventually making the model unsuitable for further studies on parasitic spreading.

Treatment with an apoptotic inducer immediately after infection was successful in overcoming the above challenge and producing a population containing a high percentage of infected cells. Induction of host cell apoptosis is considered one of the major routes for the spreading of many virulent pathogens including, *Mycobacterium tuberculosis*, *Chlamydia* and adenovirus (Aguilo *et al.*, 2013, Byrne and Ojcius, 2004 and Mi *et al.*, 2001). The link between apoptosis of host macrophages and *Leishmania* infection is complex. Parasites have shown the ability to protect host cells from apoptotic induction at early stages of infection (Moore and Matlashewski, 1994, Akarid *et al.*, 2004, Ruhland *et al.*, 2007, Lisi *et al.*, 2005 and Donovan *et al.*, 2009) but a link between apoptosis and spreading has also been reported (Real *et al.*, 2014, DaMata *et al.*, 2015, Getti *et al.*, 2008). Based on either set of evidence, apoptotic induction was expected to produce a large proportion of infected macrophages, which it did over a period of 72 and 96 hours. Therefore, to mimic a natural form of parasitic dissemination, a high proportion of *L. aethiopica* and *L. mexicana* infected cells from either 72 or 96 hours were chosen as the donor population before co-culturing with uninfected freshly differentiated THP-1 macrophages. This form of cell-to-cell infection has been successfully utilized in studies of other infectious agents such as HIV-1 (Groppelli *et al.*, 2015), *Mycobacterium* (Hagedorn *et al.*, 2009) and *L. amazonensis* (Real *et al.*, 2014). However, prior to carrying out such an experiment, it was very important to select a suitable time point for co-culture. Specifically, a time during which the effect of camptothecin treatment on infected cells viability had subsided. Careful analysis of early apoptotic markers showed that after 72 hours' infection the treated population of infected cells expressed a

similar percentage of apoptosis to the untreated, infected population. The time point of 72 hours was therefore identified as an appropriate point of co-culture.

Results from flow cytometry and trypan blue assay confirmed that the increase in percentage of infection from 1 to 12 hours was due to intracellular amastigote dissemination and not caused by a decrease in viable cell number. Moreover, microscopy clearly showed parasites spreading during which both species were filmed exiting the host cell via non-lytic mode. This corresponds with a previous report by Real et al (2014), where the authors observed that *L. amazonensis* amastigotes were surrounded by a host membrane during their egress from mice macrophages. It cannot be excluded that amastigotes could also cause macrophages to rupture and once released be picked up by neighbouring macrophages as extracellular parasites. Nevertheless, our findings suggest that *L. aethiopica* and *L. mexicana* parasites have developed a mechanism whereby they do not need to burst the host cells to spread to uninfected macrophages. Parasites can be slowly released from the host cells without damaging them. Such mechanism would explain the silent phase of amastigote spreading within the host body, when no inflammatory response is detected (Belkaid *et al.*, 2000). This mechanism in *L. aethiopica* and *L. mexicana* could be regulated via apoptotic pathways, or could involve an alternative form of induction of membrane budding and/or actin based extrusion, which needs to be investigated further.

In summary, for the very first time, we have successfully developed an *in vitro* model representative of *Leishmania* spreading in human macrophages. This was achieved in two steps. First, a population containing a high percentage of infected cells was obtained via camptothecin treatment of infected cells. This population was then used as a donor to infect a new and healthy population of differentiated THP-1 macrophages. This form of co-culture experiment allowed us to identify the timeframe for the inter-cellular dissemination of *L. aethiopica* and *L. mexicana* spreading to uninfected cells. Live microscopy revealed non-lytic release of amastigotes from cell-to-cell and provided confirmation of a strong link with

quantitative flow cytometry data on infection spreading. Following the establishment of the above model, the next step will be to functionally characterise virulence factors associated with amastigote spreading and decipher the cellular and molecular mechanism by which *Leishmania* exit the host cells in non-lytic mode, an issue neglected in the study of these parasites.

#### ACKNOWLEDGEMENTS

We thank Samantha Ingram, Richard Anthony and Rachael Nice for expert technical assistance, and Stuart Connop for proofreading the manuscript. The work was funded by the University of Greenwich VC PhD studentship.

## REFERENCES

- Aguilo, N., Marinova, D., Martin, C., Pardo, J.** (2013). ESX-1 induced apoptosis during mycobacterial infection: to be or not be, that is the question. *Frontiers in Cellular and Infection Microbiology*. **3**, 88 doi: 10.3389/fcimb.2013.00088.
- Akarid, K., Arnoult, D., Micic-Polianski, J., Sif, J., Estaquier, J., Claude Ameisen, J.** (2004). *Leishmania major* mediated prevention of programmed cell death induction in infected macrophages is associated with the repression of mitochondrial release of cytochrome c. *Journal of Leukocyte Biology*. **76**, 95-103.
- Alvar, J., Vélez, I. D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jannin, J. and de Boer, M.** (2012). Leishmaniasis worldwide and global estimates of its incidence. PLoS ONE, **7**(5). doi: 10.1371/journal.pone.0035671.
- Ashida, H., Mimuro, H., Ogawa, M., Kobayashi, T., Sanada, T., Kim, M., Sasakawa, C.** (2011). Cell death and infection: a double-edged sword for host and pathogen survival. *The Journal of Cell Biology*. **195**, 931-942.
- Belkaid, Y., Mendez, S., Lira, R., Kadambi, N., Milon, G., Sacks, D.** (2000). A natural model of *Leishmania major* infection reveals a prolonged silent phase of parasite amplification in the skin before the onset of lesion formation and immunity. *The Journal of Immunology*. **165**, 969-977.
- Bifeld, E. and Clos, J.** (2015). The genetics of *Leishmania* virulence. *Medical Microbiology and Immunology*, **204**(6), 619-634. doi: 10.1007/s00430-015-0422-1.
- Byrne, G.I., Ojcius, D.M.** (2004). *Chlamydia* and apoptosis: life and death decisions of an intracellular pathogen. *Nature Reviews Microbiology*. **2**, 802-808.
- Cameron, P., McGachy, A., Anderson, M., Paul, A., Coombs, G.H., Mottram, J.C., Alexander, J., Plevin, R.** (2004). Inhibition of Lipopolysaccharide induced macrophage IL-12

production by *Leishmania mexicana* amastigotes: the role of cysteine peptidases and the NF- $\kappa$ B signaling pathway. *The Journal of Immunology*. **173**, 3297-3304.

**DaMata, J.P., Mendes, B.P., Maciel-Lima, K., Menezes, C.A., Dutra, W.O., Sousa, L.P., Horta, M.F.** (2015). Distinct macrophages fates after in vitro infection with different species of *Leishmania*: induction of apoptosis of *Leishmania (Leishmania) amazonensis*, but not by *Leishmania (Viannia) guyanensis*. *PLOS ONE*. **10**, doi:10.1371/journal.pone.0141196.

**De Souza, S., Lang, T., Prina, E., Hellio, R.** (1995). Antoine JC. Intracellular *Leishmania amazonensis* amastigotes internalize and degrade MHC class II molecules of their host cells. *Journal of Cell Science*. **108**, 3219-3231.

**Dominguez, M., Moreno, I., Aizpurua, C., Torano, A.** (2003). Early mechanisms of *Leishmania* infection in human blood. *Microbes and Infection*. **5**, 507-513.

**Donovan, M.J., Maciuba, B.Z., Mahan, C.E., McDowell, M.A.** (2009). *Leishmania* infection inhibits cycloheximide induced macrophage apoptosis in a strain dependent manner. *Experimental Parasitology*. **123**, 58-64.

**Friedrich, N., Hagedorn, M., Soldati-Favre, D., Soldati, T.** (2012). Prison break: pathogens' strategies to egress from host cells. *Microbiology and Molecular Biology Reviews*. **76**, 707-720.

**Getti, G., Cheke, R.A., Humber, D.P.** (2008). Induction of apoptosis in host cells: a survival mechanism for *Leishmania* parasites? *Parasitology*. **135**, 1391-1399.

**Gomez, R., Oliveira, F.** (2012). The immune response to sand fly salivary proteins and its influence on *leishmania* immunity. *Frontiers in Immunology*. **3**, doi: 10.3389/fimu.2012.00110.

**Gregory, D.J., Godbout, M., Contreras, I., Forget, G., Olivier, M.** (2008). A novel form of NF- $\kappa$ B is induced by *Leishmania* infection: involvement in macrophage gene expression. *European Journal of Immunology*. **38**, 1071-1081.

- Groppelli, E., Starling, S., Jolly, C.** (2015). Contact-induced mitochondrial polarization supports HIV-1 virological synapse formation. *Journal of Virology*. **89**, 14-24.
- Hagedorn, M., Rohde, K.H., Russell, D.G., Soldati, T.** (2009). Infection by tubercular mycobacteria is spread by nonlytic ejection from their amoeba hosts. *Science*. **323**, 1729-1733.
- Hsiao, C.H.C., Ueno, N., Shao, J.Q., Schroeder, K.R., Moore, K.C., Donelson, J.E., Wilson, M.E.** (2011). The effects of macrophage source on the mechanism of phagocytosis and intracellular survival of *Leishmania*. *Microbes and Infection*. **13**, 1033-1044.
- Huang, H.L., Hsing, H.W., Lai, T.C., Chen, Y.W., Lee, T.R., Chan, H.T., Lyu, P.C., Wu, C.L., Lu, Y.C, Lin, S.T., Lin, C.W., Lai, C.H., Chang, H.T., Chou, H.C., Chan, H.L.** (2010). Trypsin induced proteome alteration during cell subculture in mammalian cells. *Journal of Biomedical Science*. **17**, doi: 10.1186/1423-0127-17-36.
- Kaye, P., Scott, P.** (2011). Leishmaniasis: complexity at the host-pathogen interface. *Nature Reviews Microbiology*. **9**, 604-615
- Kedzierski, L.** (2010). Leishmaniasis Vaccine: Where are we today? *Journal of Global Infectious Diseases* **2**, 177-185.
- Kimblin, N., Peters, N., Debrabant, A., Secundino, N., Egen, J., Lawyer, P., Fay, P., Kamhawi, S., Sacks, D.** (2008). Quantification of the infectious dose of *Leishmania major* transmitted to the skin by single sand flies. *PNAS*. **105**, 10125-10130.
- Lisi, S., Sisto, M., Acquafredda, A., Spinelli, R., Schiavone, M.A., Mitolo, V., Brandonisio, O., Panaro, M.A.** (2005). Infection with *Leishmania infantum* inhibits actinomycin D induced apoptosis of human monocytic cell line U-937. *Journal of Eukaryotic Microbiology* **52**, 211-217.
- Lodge, R., Descoteaux, A.** (2005). Modulation of phagolysosome biogenesis by the lipophosphoglycan of *Leishmania*. *Clinical Immunology*. **114**, 174-181.

- Mi, J., Li, Z., Ni, S., Steinwaerder, D., Liber, A.** (2001). Induced apoptosis supports spread of adenovirus vectors in tumours. *Human Gene Therapy*. **12**, 1343-1352.
- Mißlitz, A., Mottram, J. C., Overath, P. and Aebischer, T.** (2000). Targeted integration into a rRNA locus results in uniform and high level expression of transgenes in *Leishmania* amastigotes. *Molecular and Biochemical Parasitology* **107**, 251–261.
- Moore, K.J., Matlashewski, G.** (1994). Intracellular infection by *Leishmania donovani* inhibits macrophage apoptosis. *Journal of Immunology*. **152**, 2930-2937.
- Ogunkolade, B.W., Colomb-Valet, I., Monjour, L., Rhodes-Feuillette, A., Abita, J.P., Frommel, D.** (1990). Interactions between the human monocytic leukaemia THP-1 cell line and Old and New World species of *Leishmania*. *Acta Tropica*. **47**, 171-176.
- Patel, A.P., Deacon, A., Getti, G.** (2014). Development and validation of four *Leishmania* species constitutively expressing GFP protein. A model for drug discovery and disease pathogenesis studies. *Parasitology*. **141**, 501-510.
- Real, F., Florentina, P.T.V., Reis, L.C., Ramos-Sanchez, E.M., Veras, P.S.T., Goto, H., Mortara, R.A.** (2014). Cell-to-cell transfer of *Leishmania amazonensis* amastigotes is mediated by immunomodulatory LAMP-rich parasitophorous extrusions. *Cellular Microbiology*. **16**, 1549-1564.
- Ogunkolade, B. W., Colomb-Valet, I., Monjour, L., Rhodes-Feuillette, A., Abita, J. P. and Frommel, D.** (1990). Interactions between the human monocytic leukaemia THP-1 cell line and Old and New World species of *Leishmania*. *Acta Trop*, **47**(3), 171-176.
- Richardson, S.C., Wallom, K.L., Ferguson, E.L., Deacon, S.P., Davies, M.W., Powell, A.J., Piper, R.C., Duncan, R.** (2008). The use of fluorescence microscopy to define polymer localisation to the late endocytic compartments in cells that are targets for drug delivery. *Journal of Controlled Release*. **127**, 1-11.

- Rogers, M.E., Ilg, T., Nikolaev, A.V., Ferguson, M.A., Bates, P.A.** (2004). Transmission of cutaneous leishmaniasis by sand flies is enhanced by regurgitation of fPPG. *Nature*. **430**, 463-467.
- Ruhland, A., Leal, N., Kima, P.E.** (2007). *Leishmania* promastigotes activate PI3K/AKT signaling to confer host cell resistance to apoptosis. *Cellular Microbiology*. **9**, 84-96.
- Seifert, K., Escobar, P., Croft, S.L.** (2010). In vitro activity of anti-leishmanial drugs against *Leishmania donovani* is host cell dependent. *Journal of Antimicrobial Chemotherapy*. **65**, 508-511.
- Stamm, L.V.** (2016). Human Migration and Leishmaniasis –On the move. *DAMA Dermatology*. **152**, 373-374.

## FIGURE LEGENDS

Fig. 1. Infection of THP-1 macrophage with Leishmania promastigotes. (A) Differentiated THP-1 macrophages were infected with two promastigote species, *L. AETHIOPICA* and *L. mexicana* at 10:1 ratio (parasites to cell ratio). The percentage of infection was determined through flow cytometry at 24, 48, 72 and 96 hours of incubation. Data indicate mean percentage of infection  $\pm$  SEM from three independent experiments (B) Uninfected, *L. aethiopica* infected and *L. mexicana* infected cells were stained with trypan blue solution at 24, 48, 72 and 96 hours of incubation. Data indicate mean concentration of viable cells  $\pm$  SEM from three independent experiments. (C) Fluorescence microscopic imaging of DAPI positive nuclei of uninfected cells from 48 and 96 hours. Bar = 50  $\mu$ m. L8G, *L. aethiopica*; M5G, *L. mexicana*; \*P<0.05 and \*\*P<0.01.

Fig. 2. Effect of apoptotic induction on the rate of Leishmania infection. (A) Differentiated THP-1 macrophages were infected with *L. aethiopica* and *L. mexicana* promastigotes at 10:1 ratio (parasite to cell ratio) and treated with Campt to induce apoptosis. Excess Campt was washed away from culture medium at 24 hours after *L. aethiopica* and 5 hours after *L. mexicana* infection. The percentage of infection was determined through flow cytometry at 24, 48, 72 and 96 hours of incubation. Data indicates mean percentage of infection  $\pm$  SEM from three independent experiments. (B) Campt treated uninfected or infected cells by *L. aethiopica* and *L. mexicana* were stained with trypan blue solution at 24, 48, 72 and 96 hours of incubation. Data indicates mean concentration of viable cells  $\pm$  SEM from three independent experiments. L8G, *L. aethiopica*; M5G, *L. mexicana*; Campt, Camptothecin; \*P<0.05

Fig. 3. Infection of THP-1 macrophages with Leishmania infected cells. (A-D) After 72 hours of infection, apoptotic induced *L. aethiopica* and *L. mexicana* infected cells were co-cultured with newly differentiated THP-1 macrophage at different infection ratios, 1:1, 5:1 and 10:1 (infected to uninfected cell ratio). (A) Flow cytometry histogram plot representative of *L. aethiopica* infected cells 'M1' at 0 and 12 hours after co-culture. (B) Data indicate mean percentage of *L. aethiopica* infection  $\pm$  SEM from three independent experiments at 0, 12, 24 and 48 hours after co-culture. (C) Flow cytometry histogram plot representative of *L. mexicana* infected cells 'M1' at 0 and 12 hours after co-culture. (D) Data indicate mean percentage of *L. mexicana* infection  $\pm$  SEM from three independent experiments at 0, 12, 24 and 48 hours after co-culture. L8G, *L. aethiopica*; M5G, *L. mexicana*; \*P<0.05

Fig. 4. Measurement of viable cell concentration during infection with Leishmania infected cells. *L. aethiopica* infected co-culture (A) and *L. mexicana* infected co-culture (B) cells at different infection ratios were stained with trypan blue solution at 0, 12, 24 and 48 hours after co-culture. Data indicates mean concentration of viable cells  $\pm$  SEM from three independent experiments. L8G, *L. aethiopica*; M5G, *L. mexicana*

Fig. 5. Cell-to-cell spreading of Leishmania amastigotes. Freshly differentiated THP-1 cells acting as recipient uninfected cells (labelled as R), were pre-stained with CellMask™ orange plasma membrane dye. The recipient cells were then co-cultured with *L. aethiopica* (A) and *L. mexicana* (B) infected cells, which act as donor cells (labelled as D). The time of acquisition is represented by hours (h) and minutes (min). Image acquisition started at 1 hour after co-culture. Bar = 25  $\mu$ m.

Video 1. Live cell imaging of *L. aethiopica* amastigotes spread from cell-to-cell. Freshly differentiated THP-1 cells acting as recipient uninfected cells, were pre-stained with CellMask™ orange plasma membrane dye. The recipient cells were then co-cultured with *L. aethiopica* (A) and *L. mexicana* (B) infected cells, which act as donor cells. Image acquisition started at 1 hour after co-culture.

Video 2. Live cell imaging of *L. mexicana* amastigotes spread from cell-to-cell. Differentiated THP-1 macrophage, which is the recipient uninfected cells, was pre-stained with CellMask™ orange plasma membrane dye. The recipient cells were then co-cultured with *L. mexicana* infected cells, which are the donor cells. Image acquisition started at 1 hour after co-culture.

Supplementary Fig. S1. Effect of Leishmania infection on the induction of apoptosis in the presence and/or absence of apoptotic inducer. Differentiated THP-1 macrophages were infected with *L. aethiopica* and *L. mexicana* promastigotes at 10:1 ratio (parasite to cell ratio) and treated with and/or without Campt. Excess Campt was washed away from culture medium at 24 hours after *L. aethiopica* and 5 hours after *L. mexicana* infection. The percentage of apoptosis in *L. aethiopica* and *L. mexicana* infected cells were determined through Annexin V assay at 72 hours (A) and 96 hours (B) of incubation. Data indicate mean percentage of apoptosis  $\pm$  SEM from three independent experiments. L8G, *L. aethiopica*; M5G, *L. mexicana*; Campt, Camptothecin; \*P<0.05 and \*\*P<0.01.