

1 Running head: Plants influence bee disease transmission

2

3 Disease where you dine: Plant species and floral traits associated with pathogen transmission in  
4 bumble bees

5

6 Lynn S. Adler<sup>1\*</sup>, Kristen M. Michaud<sup>1</sup>, Stephen P. Ellner<sup>2</sup>, Scott H. McArt<sup>3</sup>, Philip C.  
7 Stevenson<sup>4,5</sup>, and Rebecca E. Irwin<sup>6</sup>

8 <sup>1</sup> Department of Biology, University of Massachusetts, Amherst, MA 01003, USA

9 <sup>2</sup> Department of Ecology and Evolutionary Biology, Cornell University, Ithaca NY 14853, USA

10 <sup>3</sup> Department of Entomology, Cornell University, Ithaca, NY 14853, USA

11 <sup>4</sup> Royal Botanic Gardens, Kew, Richmond TW9 3AB, Surrey, England

12 <sup>5</sup> Natural Resources Institute, University of Greenwich, Chatham ME4 4TB, Kent, England

13 <sup>6</sup> Department of Applied Ecology, North Carolina State University, Raleigh, NC 27695 USA

14

15 \* corresponding author: [lsadler@bio.umass.edu](mailto:lsadler@bio.umass.edu), ph 413-545-1060, fax 413-545-3243

16

**17 Abstract**

18           Hotspots of disease transmission can strongly influence pathogen spread. Bee pathogens  
19 may be transmitted via shared floral use, but the role of plant species and floral trait variation in  
20 shaping transmission dynamics is almost entirely unexplored. Given the importance of pathogens  
21 for the decline of several bee species, understanding whether and how plant species and floral  
22 traits affect transmission could give us important tools for predicting which plant species may be  
23 hotspots for disease spread. We assessed variation in transmission via susceptibility (probability  
24 of infection) and mean intensity (cell count of infected bees) of the trypanosomatid gut pathogen  
25 *Crithidia bombi* to uninfected *Bombus impatiens* workers foraging on 14 plant species, and  
26 assessed the role of floral traits, bee size and foraging behavior on transmission. We also  
27 conducted a manipulative experiment to determine how the number of open flowers affected  
28 transmission on three plant species, *Penstemon digitalis*, *Monarda didyma*, and *Lythrum*  
29 *salicaria*. Plant species differed fourfold in the overall mean abundance of *Crithidia* in foraging  
30 bumble bees (mean including infected and uninfected bees). Across plant species, bee  
31 susceptibility and mean intensity increased with the number of reproductive structures per  
32 inflorescence (buds, flowers and fruits); smaller bees and those that foraged longer were also  
33 more susceptible. Trait-based models were as good or better than species-based models at  
34 predicting susceptibility and mean intensity based on AIC values. Surprisingly, floral size and  
35 morphology did not significantly predict transmission across species. In the manipulative  
36 experiment, more open flowers increased mean pathogen abundance fourfold in *Monarda*, but  
37 had no effect in the other two plant species. Our results suggest that variation among plant  
38 species, through their influence on pathogen transmission, may shape bee disease dynamics.  
39 Given widespread investment in pollinator-friendly plantings to support pollinators,

40 understanding how plant species affect disease transmission is important for recommending plant  
41 species that optimize pollinator health.

42

43 **Key words:** bee decline, bee parasites, *Bombus impatiens*, *Crithidia*, environmental reservoir,  
44 floral traits, foraging behavior, trait-based, transmission hotspots

45

## 46 **Introduction**

47 Pathogen transmission is mediated by environmental heterogeneity (reviewed in Paull et  
48 al. 2012) and can be influenced by features of the transmission site. ‘Hotspots’ are regions  
49 characterized by particularly high pathogen prevalence or incidence, and can be sources of  
50 transmission to less infected areas (Paull et al. 2012). For example, the bacteria that cause  
51 cholera can concentrate on water hyacinth, which prolongs pathogen longevity (Spira et al.  
52 1981). Thus, the presence of water hyacinth at sites may create a ‘hotspot’ that results in  
53 increased transmission across the landscape. Similarly, species may vary in their ability to  
54 transmit pathogens, even given similar levels of pathogen in the environment. For example, six  
55 grass species varied in host susceptibility, competence, and vector population sizes when  
56 exposed to Barley Yellow Dwarf virus, and several traits associated with life history were  
57 associated with this variation (Cronin et al. 2010). The goal of our study was to assess the extent  
58 to which plant species vary as hotspots for bee disease transmission, and, if so, the potential for  
59 floral traits to explain this variation.

60 Populations of many wild bee species are in decline, and pathogens have been implicated  
61 as one of the likely causal factors (e. g., Goulson et al. 2015). There is increasing evidence that  
62 bees share pathogens within and across species (e. g., Gamboa et al. 2015, McMahon et al.

63 2015), including transmission from widespread managed species such as *Apis mellifera* to wild  
64 *Bombus* hosts (Fürst et al. 2014). *Crithidia bombi* (Zoomastigophora:Trypanosomatidae) is a gut  
65 trypanosome that infects a wide range of bumble bee species with infection rates up to 80%  
66 (Shykoff and Schmid-Hempel 1991a, Gillespie 2010). *Crithidia* can impair learning, reduce  
67 colony reproduction under food limitation, reduce a queen's ability to found new colonies, and is  
68 associated with decreased reproduction in wild colonies (e. g., Shykoff and Schmid-Hempel  
69 1991b, Brown et al. 2003, Gegear et al. 2006, Goulson et al. 2017). This pathogen is transmitted  
70 when feces from an infected individual are consumed by an uninfected bee (Durrer and Schmid-  
71 Hempel 1994). While there are obvious routes for transmission within colonies, the  
72 environmental factors that contribute to horizontal transmission are largely unknown.

73         Flowers can be visited by a wide range of pollinators and other species (McArt et al.  
74 2014) and are logical suspects as hotspots of pathogen transmission among bees, but very little  
75 empirical work has addressed this (reviewed in Koch et al. 2017). More than 20 years ago, a  
76 landmark study showed that *Bombus terrestris* could become infected with *Crithidia* by foraging  
77 on inflorescences previously foraged on by infected bees (Durrer and Schmid-Hempel 1994).  
78 More recent work has demonstrated that *Crithidia* was shared among three *Bombus* species, and  
79 that potential to transmit the pathogen varied among plant species (Ruiz-Gonzalez et al. 2012).  
80 Furthermore, multiple pathogens, including *Crithidia* spp., *Nosema* spp. and viruses, can be  
81 transmitted among bee species via shared flower use (Ruiz-Gonzalez and Brown 2006, Singh et  
82 al. 2010, Graystock et al. 2015).

83         Plant species may differ in their potential to transmit pathogens, and such variation could  
84 be mediated by floral traits. Pathogen transmission among bees via shared flower use was  
85 different on two different plant species in each of two studies (Durrer and Schmid-Hempel 1994,

86 Graystock et al. 2015). This suggests that plant community composition can affect transmission,  
87 but consideration of so few plant species limits our ability to generalize. Furthermore, since each  
88 study used only two plant species that differ in many ways, it is not possible to determine which  
89 traits might be responsible for differences in transmission. Only one study has manipulated floral  
90 traits to assess their role in transmission; Durrer and Schmid-Hempel (1994) manipulated  
91 inflorescence architecture in a single plant species, and found that *B. terrestris* were more likely  
92 to become infected after foraging on inflorescences with a 'linear' rather than 'spiral'  
93 arrangement of flowers. This suggests that floral traits can affect disease transmission in foraging  
94 bees, but more work is needed across a broader range of plant species to evaluate this hypothesis.  
95 More generally, both floral and pollinator morphology can be important for efficient pollen  
96 transfer (e. g., Montgomery and Rathcke 2012). Thus, it is a logical extension to hypothesize that  
97 floral morphology could also influence pathogen transmission. Given that pathogen transmission  
98 among bees may be widespread (Fürst et al. 2014, Graystock et al. 2016), it is important to  
99 understand whether and how plant species mediate transmission.

100         While documenting plant species variation in transmission would provide an important  
101 first step to understanding how plant community composition influences pollinator-disease  
102 dynamics, a trait-based approach (Westoby and Wright 2006, Webb et al. 2010) to understanding  
103 disease transmission has several potential advantages over species-by-species approaches. If  
104 traits alone can predict transmission as well as models incorporating species identity, the effort  
105 required to parameterize transmission rate models for complex communities may be greatly  
106 reduced, because many relevant traits (e.g., measures of individual size and life history) are  
107 easily obtained from publicly available databases. For example, the probability that rodent  
108 species were zoonotic reservoirs could be predicted with approximately 75% accuracy based on

109 only five host traits; considering 11 host traits improved predictions to >90% (Han et al. 2015).  
110 Trait-based analyses are also potentially generalizable between taxonomically distinct  
111 communities, while a species-based approach requires a new study for each new species.

112 To evaluate whether variation among plant species can shape pathogen transmission to  
113 foraging bees and to assess the role of floral traits in mediating these dynamics, we used 14 bee-  
114 pollinated plant species from eight different families, encompassing a range of floral trait  
115 variation. We allowed uninfected, individual *Bombus impatiens* workers to forage on  
116 inflorescences provided with *Crithidia* inoculum, and then reared these bees and compared  
117 transmission across species, measured as susceptibility (probability of becoming infected) and  
118 mean intensity (cell counts of infected bees). We also measured floral traits for each species, and  
119 evaluated the effect of these traits, bee foraging behavior, and bee size on susceptibility and  
120 mean intensity across plant species. We then compared how well trait-based models and species-  
121 based models explained variation in susceptibility and mean intensity. We hypothesized that  
122 traits that increase encounter rate with pathogens, such as small flowers and wide or nonexistent  
123 corolla tubes, and bee behavior, such as number of flowers visited or total time foraging, would  
124 increase susceptibility or intensity of infection. Floral traits could also affect how much pathogen  
125 inoculum is consumed per encounter, which positively relates to infection intensity (Otterstatter  
126 and Thomson 2006). Finally, nectar production or floral morphology could affect desiccation,  
127 which is important for viability of some pathogens such as *Crithidia* (Figueroa et. al.,  
128 unpublished manuscript). Because our results suggested that reproductive structures per  
129 inflorescence predicted transmission, we then conducted transmission trials on three plant  
130 species in which we experimentally manipulated open flowers per inflorescence. Ultimately, our  
131 goal was to elucidate the role of flowering species and floral traits in bee disease dynamics.

132

133 **Materials and Methods**134 Transmission trials across 14 plant species

135 *Study site and species.* This research was conducted at the University of Massachusetts Center  
136 for Agriculture (South Deerfield, MA, U.S.A., 42° 28.6' N, 72° 34.8' W) in 2014. The 14 plant  
137 species included in the study were *Antirrhinum majus* (Plantaginaceae), *Asclepias incarnata*  
138 (Asclepiadaceae), *Digitalis purpurea* (Plantaginaceae), *Eupatorium perfoliatum* (Asteraceae),  
139 *Helianthus annuus* (Asteraceae), *Impatiens capensis* (Balsaminaceae), *Linaria vulgaris*  
140 (Plantaginaceae), *Lobelia siphilitica* (Campanulaceae), *Lythrum salicaria* (Lythraceae), *Monarda*  
141 *didyma* (Lamiaceae), *Penstemon digitalis* (Plantaginaceae), *Solidago canadensis* (Asteraceae),  
142 *Thymus vulgaris* (Lamiaceae), and *Verbascum thapsus* (Scrophulariaceae); for simplicity we  
143 refer to all species by genus hereafter. Many of these species were selected from 'bee friendly'  
144 suggested planting lists (e.g. Xerces society; [http://www.xerces.org/pollinator-](http://www.xerces.org/pollinator-conservation/plant-lists/)  
145 [conservation/plant-lists/](http://www.xerces.org/pollinator-conservation/plant-lists/)), and others were chosen for particular interest as invasive, common  
146 horticultural species. Overall, we chose species representing a wide range of variation in traits  
147 including flower size, number, and morphology. Some species were obtained from local  
148 nurseries or grown from seed and transplanted to the field site; others were collected from  
149 naturally-growing areas nearby (Appendix S1).

150 We used commercial colonies of *Bombus impatiens* (Biobest, Leamington, Ontario,  
151 Canada), the common eastern bumble bee, which is the most prevalent wild bumble bee species  
152 in our region (Gillespie 2010). Because *B. impatiens* is widely distributed commercially,  
153 understanding how plants mediate transmission in this species is particularly important due to the  
154 potential for spread from commercial to wild bees (Colla et al. 2006). Colonies were confirmed

155 to be *Crithidia*-free with weekly dissections of five bees per colony. *Crithidia* was maintained in  
156 a ‘source’ colony that was originally infected from wild *B. impatiens* workers collected from two  
157 sites in Amherst, MA, U.S.A. (42°24'32.47"N 72°31'39.57"W; 42°23'20"N 72°31'21"W) and  
158 then transferred to new source colonies as needed. Four source colonies were used over the  
159 course of the experiment; usually only one source was used per day, but on four dates two  
160 sources were used to produce enough inoculum. Colonies were provided *ad libitum* with 30%  
161 sucrose solution replaced weekly, and approximately 10 g pollen loaves made of 30% sucrose  
162 mixed with multifloral honey bee-collected pollen (Koppert Biological Systems, Howell,  
163 Michigan) added every other day.

164 *Inoculum preparation.* To create *Crithidia* inoculum for use in transmission trials, we  
165 dissected up to 10 bees per day from a single source colony. Guts were ground in 300  $\mu\text{L}$   
166 deionized water in microcentrifuge tubes and left to sit for 4 h. Moving *Crithidia* cells were  
167 counted on a Neubauer hemacytometer in a 0.02  $\mu\text{L}$  subsample from a 10  $\mu\text{L}$  sample per bee  
168 using a light microscope at 40x magnification. Because one of the three life stages of *Crithidia* is  
169 non-motile, we note that by counting only moving *Crithidia* cells we introduced some variation  
170 in the number of infective cells in inoculum made each day; this value also varies daily because  
171 we estimate concentration from a small subsample of gut solution. Thus, random variation in  
172 daily inoculum concentration makes our results a more conservative test of plant species  
173 differences. After counting *Crithidia* cells, we then combined 150  $\mu\text{L}$  of gut solution from up to  
174 five bees each day, diluted this with deionized water, and then mixed with an equal volume of  
175 50% sucrose to create a final solution of 25% sucrose with 600 cells  $\mu\text{L}^{-1}$ . Thus, our inoculum  
176 had a *Crithidia* cell concentration within the natural concentration occurring in feces (Otterstatter  
177 and Thomson 2006) and also a sugar concentration within the range of nectar; the average



178 concentration from species we were able to measure in this study was 30% (range: 11.5-55%;  
179 data not shown). We recorded the time at which inoculum preparation was completed each day,  
180 and transported inoculum to the field site in a cooler with ice packs to minimize loss of viability.

181 *Transmission trials.* During natural foraging to wild plants, floral traits could influence  
182 transmission of bee pathogens at flowers by altering 1) the likelihood of depositing pathogens on  
183 flowers, 2) pathogen viability in flowers, 3) the likelihood of encountering flowers that contain  
184 pathogens, and 4) pathogen acquisition and establishment in hosts upon visiting flowers that  
185 contain pathogens (reviewed in McArt et al. 2014). Although we would have ideally assessed all  
186 four of these mechanisms by allowing uninfected bees to forage on plants previously visited by  
187 infected bees (as in Durrer and Schmid Hempel 1994, which used *B. terrestris*), we were unable  
188 to replicate results of that study, suggesting that natural transmission rates in *B. impatiens* are too  
189 low to be detected by this approach. Instead, we compared transmission potential between plant  
190 species by adding controlled amounts of inoculum to *Crithidia*-free inflorescences, allowing a  
191 single uninfected bee to forage, rearing the bee for 7 days and then assessing susceptibility  
192 (presence/absence of pathogens following exposure) and mean intensity (cell counts in infected  
193 bees). This methodology evaluates processes three and four – likelihood of encountering flowers  
194 that contain pathogens (via foraging behavior upon encountering an infected plant), and  
195 pathogen acquisition and establishment in hosts upon visiting flowers that contain pathogens.  
196 Each of these are major unexplored components of pathogen transmission by bees at flowers  
197 (McArt et al. 2014) that could be affected by floral number, size, shape, or nectar production, as  
198 well as (or in addition to) bee foraging behavior. This methodology does not account for  
199 likelihood of depositing pathogen cells on flowers (process one), or variation in viability in  
200 flowers (process two), which are beyond the scope of this paper but are being explored in a

201 forthcoming manuscript (Figueroa et al., unpublished manuscript). In a subsequent large-scale  
202 experiment, we ranked plant species as ‘high’ or ‘low’ transmission based on the trials reported  
203 here, and conducted an experiment with infected bee microcolonies foraging on uninoculated  
204 ‘high’ or ‘low’ transmission plants. Average colony-level infection after two weeks was  
205 approximately twice as high when foraging on ‘high’ compared to ‘low’ transmission plants  
206 (Adler et al., unpublished data), suggesting that processes of transmission we tested in this study  
207 explain substantial variation in longer-term transmission dynamics.

208         Plants were grown and trials were conducted in the field at the University of  
209 Massachusetts Center for Agriculture (South Deerfield, MA, U.S.A., 42° 28.6’ N, 72° 34.8’ W)  
210 from June 24 through August 28, 2014. Whenever possible, we used multiple plant species on  
211 each trial date, but each species was only used on a subset of all possible dates due to phenology.  
212 To conduct transmission trials, inflorescences of all plant species were covered with organza  
213 bags (ULine, Pleasant Prairie, Wisconsin) before flowers opened to prevent wild bee visitation  
214 and potential pathogen deposition. To conduct a trial, an inflorescence with at least five open  
215 flowers was clipped with scissors and immediately placed in a florist’s water tube. We counted  
216 open flowers, placed four 10  $\mu$ L inoculum drops within four separate flowers (one drop per  
217 flower) using a pipette, and marked these flowers at the outside base, calyx or stem with paint  
218 pens (Craftsmart® Fine Line 6 Count, Basic, Michaels Stores, Inc., Irving, Texas). For  
219 *Eupatorium* and *Solidago*, capitula were considered ‘flowers,’ while in *Helianthus* we used a  
220 single capitulum and individual florets were counted as flowers. We chose 10  $\mu$ L to simulate the  
221 volume of feces from a single defecation event. Four drops were used to facilitate encounters  
222 during foraging; by having a minimum of five open flowers we ensured there was at least one  
223 un-inoculated flower to visit. Bees almost always consumed inoculum upon first contact. We

224 placed inoculum in contact with reproductive structures whenever possible; this was typically  
225 inside tubular flowers (e.g., *Lobelia*, *Penstemon*) or on top of open flowers (e.g., *Lythrum*). In  
226 some cases, flowers were so small that the drop rested on top of the corolla (e.g., *Solidago*,  
227 *Thymus*, *Eupatorium*) and may not have contacted nectar. We placed drops inside flowers due to  
228 initial findings of *Crithidia* in nectar and bees defecating on flowers (Durrer and Schmid-Hempel  
229 1994, Otterstatter and Thomson 2006). More recent work suggests that feces are more likely to  
230 be deposited on outer floral surfaces and not in nectar (Cisarovsky and Schmid-Hempel 2014),  
231 although data show that bees deposit up to 47% of their feces within flowers on some plant  
232 species (Figueroa et al., unpublished manuscript). Our goal was to standardize the amount and  
233 presentation of inoculum across species so we could assess variation in susceptibility and  
234 intensity of infection given the same starting conditions, after controlling for foraging-induced  
235 differences in exposure.

236 Each inflorescence was individually placed into a small cage (45.7 cm x 71.0 cm x 55.6  
237 cm) constructed of a wood frame with plexiglass or cloth sides with a chilled, uninfected  
238 experimental *B. impatiens* worker initially placed on the inflorescence. For each trial, we  
239 recorded the plant species, experimental bee colony of origin, start and end time, time spent  
240 foraging (i.e., actively probing flowers), and the number of open flowers probed and number of  
241 inoculum drops probed. For the latter two measures, every new entry into a flower in which  
242 reproductive parts were contacted was considered a new flower probe (and if the flower was  
243 inoculated, it was also a new drop probed) because we could not ascertain whether bees  
244 consumed all the inoculum drop in a single probe. A trial was concluded after the bee ceased  
245 foraging, if at least one inoculated flower was probed. We did not limit trials to a specified time  
246 period because the rate at which bees probed individual flowers varied widely with species, and

247 so limiting trial time period would create a *de facto* difference in the number of flowers probed  
248 per species. Bees that did not forage on an inoculated flower after 20 min were excluded. After  
249 each trial, inflorescences were disposed of and experimental bees were returned to a cooler on  
250 ice until transport to the laboratory at the end of the field day. We used 6-10 experimental  
251 colonies per plant species and had 11-36 bees with successful trials and pathogen counts per  
252 plant species.

253 *Assessing pathogen infection.* Upon returning to the laboratory, each bee was placed in a  
254 20 mL plastic scintillation vial with a nectar feeder with 500  $\mu$ L of 30% sucrose solution and a  
255 0.1-0.2 g portion of a pollen loaf; all pollen loaves were made from the same pollen source used  
256 to maintain colonies. Bees were housed in a growth chamber at 27°C in darkness, and placed in  
257 new vials with fresh nectar and pollen daily. After 7 days, bees were dissected and *Crithidia* cells  
258 were counted as in ‘*Inoculum preparation*’ above, except that guts were left for 5 h instead of 4 h  
259 before counting (the shorter time for inoculation preparation allowed us to begin field trials  
260 sooner). We collected the right forewing of each bee to measure radial cell length as an estimate  
261 of bee size (Harder 1982); we refer to this as ‘bee size’ in analysis.

262 *Measuring floral traits.* To understand the role of floral traits in mediating bee disease  
263 transmission, we measured reproductive structures per inflorescence, floral size and morphology,  
264 nectar production, and nectar secondary chemistry. We measured these floral traits on single  
265 inflorescences from 22-38 (median 30) individuals of each plant species that were not used in  
266 transmission trials (sample sizes are provided in Appendix S1, Table S1). While it would have  
267 been ideal to measure floral traits on the inflorescences used in trials, this would have been  
268 prohibitively time consuming and, in the case of nectar measurements, potentially damaging to  
269 flowers. However, we included the number of open flowers for each trial in analysis. For all

270 other floral traits, we used separate plants to measure species-level values for use in analyses  
271 relating traits to transmission. We measured corolla length and width using digital calipers to the  
272 nearest 0.01 mm (Appendix S2), and used these traits in a principal components analysis to  
273 generate a first component that reflected floral size ( $PC1 = 0.87 \times \text{corolla length} + 0.50 \times \text{corolla}$   
274  $\text{width}$ , accounting for 91% of total variance), and a second component representing floral shape  
275 ( $PC2 = 0.5 \times \text{corolla length} - 0.87 \times \text{corolla width}$ , accounting for 9% of total variance), which  
276 correlated strongly ( $r = 0.88$ ) with the ratio of corolla length:width. We counted reproductive  
277 structures per inflorescence (including buds, flowers and fruits), and measured the height of the  
278 tallest flower. We measured nectar volume after 24 h of bagging using glass microcapillary  
279 tubes; we did not remove nectar prior to bagging flowers to avoid damaging nectaries. We did  
280 not include sugar content in this study since nectar production was too low to measure sugar on  
281 several species. For a subset of species, in 2015 we measured floral longevity by marking buds  
282 and noting the date of first opening and senescence. We present means, sample sizes, and  
283 standard deviation for all predictor traits used in analyses in Appendix S3.

284 *Statistical analysis.* Statistical analyses were conducted in R (R Core Team 2017) version  
285 3.3 or higher. We analyzed susceptibility (presence/absence of *Crithidia*) and mean intensity  
286 (mean raw *Crithidia* cell count per 0.02  $\mu\text{l}$  gut sample in infected bees) as two separate  
287 components of pathogen transmission to bees. Although using a single response with negative  
288 binomial regression should be more powerful, this was inappropriate for our data (see Appendix  
289 S4 for justification). However, we summarize patterns using a combined response variable for  
290 the purposes of comparing species only. Susceptibility is a binary (0-1) response and was  
291 therefore analyzed by logistic regression. Mean intensity (values of all non-zero counts) was  
292 highly right-skewed, so our analyses used log-transformed counts; these had a symmetric and

293 approximately Gaussian distribution, and were therefore analyzed by linear regression assuming  
294 Gaussian error distributions. Models with only fixed effects were fitted using glm (for  
295 susceptibility) and lm (for intensity) functions in R. Models that included any random effects  
296 were fitted using the gam function in the mgcv package (Wood 2006). Note that the fitted  
297 models were logistic or linear mixed regression models, not generalized additive models. We  
298 used the gam function because, for the models we consider here (which do not include multiple  
299 random effects with correlations), gam reports the statistical significance ( $P$  value) for random  
300 effect terms specified through the random effect (“re”) basis.

301 Our statistical analyses assessed whether susceptibility and mean intensity were predicted  
302 by plant species and by floral traits, and whether these responses were better predicted by species  
303 identity or by floral traits. We initially fitted generalized additive models for nonlinear effects of  
304 day of year (Julian date), the elapsed time between inoculum preparation and each trial, or both,  
305 on susceptibility and mean intensity using gam in the mgcv package, but these covariates did not  
306 affect susceptibility or mean intensity ( $P > 0.1$  from anova.gam) and were omitted from  
307 subsequent analyses. We could not include the effect of experimental colony in our analyses  
308 because this was confounded with plant species, since we used different colonies over the course  
309 of the summer and plant species bloomed at different times. However, we have no *a priori*  
310 reason to think that experimental colonies, sourced commercially and reared in the lab, would  
311 vary systematically in susceptibility to infection over a three-month period.

312 Plant species and pathogen transmission. To assess covariates for inclusion in models that  
313 used plant species as a predictor of susceptibility and mean intensity, we first fitted a series of  
314 linear (intensity) and generalized linear (susceptibility) models with single predictors, including  
315 plant species, inoculum source colony, bee size (estimated as wing radial cell length), and bee

316 foraging behavior (number of flowers probed, number of inoculum drops probed, and total time  
317 foraging) as fixed covariates, using both susceptibility and mean intensity as responses in  
318 separate analyses. Species identity and inoculum source colony were factor variables; species  
319 identity was fitted as a random effect but inoculum source colony was fitted as fixed since there  
320 were only six levels (four sources plus two combinations used on some days); all other  
321 covariates are numerical and were fitted as fixed effects. Only covariates that were significantly  
322 or marginally significantly related to susceptibility or mean intensity were retained for model  
323 selection (described in Results and Appendix S4).

324 Traits and transmission. For models using floral traits rather than species identity to  
325 predict susceptibility and mean intensity, we again assessed potential covariates by fitting a set  
326 of linear or generalized linear models with potential covariates as single fixed effects. Potential  
327 covariates were floral traits (corolla size, corolla shape, number of open flowers, reproductive  
328 structures per inflorescence, nectar volume and floral longevity), bee size (estimated as wing  
329 radial cell length), and bee foraging behavior (number of flowers probed, number of inoculum  
330 drops probed, and total time foraging). Traits that were significant predictors in these analyses  
331 were used in model selection (described in Results) to produce final trait-based models for  
332 susceptibility and mean intensity. Species identity was not included in trait-based models  
333 because it is confounded with floral traits, which were measured at the species level. *Helianthus*  
334 was an outlier for several floral traits and foraging behavior measures (see Appendix S4), and so  
335 was omitted from analyses of trait-dependent transmission but included in analyses that assessed  
336 species differences in susceptibility or mean intensity without considering floral traits.

337 Species vs. traits as predictors of susceptibility and mean intensity. Trait-based and  
338 species-based models are non-nested, so model comparison was done using AIC omitting

339 *Helianthus*, because comparison is only possible when all models are fitted to the same data. We  
340 selected the lowest AIC models for both species (with bee size and behavior as potential  
341 covariates, but not floral traits) and traits (with bee size, behavior and floral traits as potential  
342 covariates, but not species) to determine which most effectively predicted susceptibility and  
343 mean intensity.

344

#### 345 Transmission trials manipulating flower number

346 *Experimental trials.* In our observational transmission trials, reproductive structures per  
347 inflorescence was the most consistent predictor of susceptibility and mean intensity (see  
348 Results). This was a surprising result, since the number of open flowers did not predict responses  
349 as strongly. This suggests that some unmeasured trait correlated with reproductive structures per  
350 inflorescence affects transmission. Alternatively, because number of open flowers and  
351 reproductive structures per inflorescence were tightly correlated across species (Spearman's  $\rho =$   
352 0.83,  $n = 14$ ,  $P < 0.001$  for all species;  $\rho = 0.79$ ,  $n = 13$ ,  $P < 0.01$  excluding *Helianthus*), this  
353 observational approach may not be able to distinguish whether number of open flowers or some  
354 other correlated unmeasured trait is the underlying cause of altered transmission.

355 To determine whether the number of open flowers or some other correlated trait underlies  
356 the relationship between number of reproductive structures per inflorescence and transmission,  
357 we conducted transmission trials manipulating flower number and comparing transmission  
358 within three plant species. Although intraspecific variation in number of open flowers may affect  
359 susceptibility or mean intensity differently than interspecific variation, manipulating this trait  
360 within species allows us to assess its importance in the absence of confounding species-level  
361 differences in other traits, and using three plant species provides some generality to this



362 assessment. Trials were conducted in 2016 on *Penstemon* (June 13-29), *Monarda* (June 30 – July  
363 15) and *Lythrum* (July 18 – Aug 9) using plants and protocols from the same site and trials  
364 described previously, except that each inflorescence was assigned to a high or low flower  
365 number treatment in alternating sequence, and inoculum was made using ¼ strength Ringer’s  
366 solution ((Sigma-Aldrich, St. Louis, MO, USA) instead of deionized water. The number of open  
367 flowers for each treatment varied with species; the low and high range of open flowers was 5-7  
368 and 11-13 for *Penstemon*, 10-15 and 25-30 for *Monarda*, and 6-10 and 16-20 for *Lythrum*. These  
369 numbers were based on typical flower production for each species, and such that the upper bound  
370 of the low treatment was half of the upper bound of the high treatment, with constant range of  
371 values within treatment. We only selected inflorescences with at least one more open flower than  
372 the maximum ‘high’ treatment value (e.g., 14 for *Penstemon*). We then randomly assigned  
373 inflorescences to treatments and removed open flowers using dissecting scissors or forceps to  
374 achieve the appropriate number; at least one open flower was removed from every inflorescence  
375 to control for damage effects. The number of open flowers removed per inflorescence and the  
376 final number of open flowers were recorded for each trial, along with the bee behavior variables  
377 described for the previous transmission trials. Bees came from six experimental colonies.

378 *Statistical analysis.* Prior to analysis, we discarded six bees due to unusual foraging (e. g,  
379 spending several minutes inside a single flower), death or missing data, and two extreme outliers  
380 (Appendix S4), resulting in final sample sizes of 63, 49 and 65 bees in *Penstemon*, *Monarda* and  
381 *Lythrum* trials respectively. Because this experiment focused on the within-species effect of  
382 number of open flowers, rather than comparing species differences, we analyzed each species  
383 separately. This allowed us to analyze counts for each species as a single response, including  
384 both infected and uninfected bees, using negative binomial regression, because the treatment

385 with higher mean count was also the treatment with a higher frequency of nonzero counts. Using  
386 mean abundance of *Crithidia* as the response (including uninfected bees) combines susceptibility  
387 and mean intensity into one response variable. We used R functions `glm.nb` for models with only  
388 fixed effects and `gam` with `family=nb` for models including random effects, in both cases using  
389 the default log link function.

390 As in the multi-species infection trials, we first assessed whether other potential  
391 covariates (bee size, trial time, time foraging, number of inoculum drops probed, number of  
392 flowers probed, minutes between inoculum preparation and trial, bee dissection time) should be  
393 included in subsequent analyses. Because we analyzed species separately and inoculum strength  
394 can vary daily, we also included trial date as an unordered, categorical random effect. We fitted  
395 negative binomial regression models (Appendix S4) that always included treatment as a fixed  
396 effect, and trial date and bee colony of origin as random effects. In each model, only one  
397 additional covariate was included, whose significance was tested by `anova.gam`. Additional  
398 covariates that were significant predictors in these analyses were used in model selection  
399 (described in Appendix S4) to produce the final model for testing treatment effects.

400

## 401 **Results**

### 402 Transmission trials across 14 plant species

#### 403 *Plant species and pathogen transmission*

404 Tabulating mean pathogen counts per foraging bee showed that plant species varied  
405 fourfold in mean *Crithidia* abundance (mean count including uninfected bees; Fig. 1A). Mean  
406 abundance was highest in *Asclepias*, and high in *Monarda*, *Lythrum*, and *Lobelia*, and lowest in  
407 *Digitalis*, *Antirrhinum*, *Linaria*, and *Thymus*.

408 Plant species and bee size were significant predictors of variation in susceptibility in  
409 single-variable analyses, with larger bees having lower susceptibility (Table 1). Both remained  
410 significant in a generalized linear mixed model including both variables with species as a random  
411 effect (species:  $n = 293$ ,  $df = 4.6$ ,  $\chi^2 = 8.68$ ,  $P = 0.031$ ; bee size:  $n = 293$ ,  $z = -2.096$ ,  $P = 0.036$ ;  
412 Fig. 1B). Total foraging time (marginally significant in the single-variable analysis; Table 1) was  
413 not significant in a generalized linear mixed model that also included species as a random effect  
414 ( $n=298$ ,  $z=1.633$ ,  $P=0.102$ ). Plant species also predicted mean intensity (Table 2, Fig. 1C). No  
415 other bee behavior covariates were significant predictors of mean intensity in the single-variable  
416 analyses. Patterns in mean abundance were largely reflected in the patterns for mean intensity  
417 (Figs 1A vs. 1C). Some species, such as *Antirrhinum*, had high susceptibility but low mean  
418 intensity, leading to low overall mean abundance.

419

#### 420 *Traits and transmission*

421 In our initial analysis testing each potential predictor one at a time on susceptibility,  
422 reproductive structures per inflorescence, bee size, and total time foraging were significant or  
423 marginally significant (Table 1). Bee size and total time foraging were correlated ( $n = 280$ ,  $r = -$   
424  $0.25$ ,  $P < 0.001$ ), so we fitted two GLMs including each of these separately as a covariate in  
425 addition to reproductive structures per inflorescence. In the first model, bees visiting plant  
426 species with more reproductive structures per inflorescence were more likely to acquire *Crithidia*  
427 ( $n = 293$ ,  $\chi^2 = 6.676$ ,  $P = 0.010$ ; Fig. 2A), as were smaller bees ( $n = 293$ ,  $\chi^2 = 5.11$ ,  $P = 0.024$ ;  
428 Fig. 2B). In the second model, bees visiting plant species with more reproductive structures per  
429 inflorescence were again more likely to acquire *Crithidia* ( $n = 298$ ,  $\chi^2 = 8.93$ ,  $P = 0.003$ ), as were  
430 bees with greater total foraging time ( $n = 298$ ,  $\chi^2 = 5.55$ ,  $P = 0.018$ ; Fig. 2C). Of these two

431 models, the one with total time foraging has the lower AIC ( $\Delta\text{AIC}=3.16$ ), and in a GLM with  
432 both foraging time and bee size as covariates, bee size is not significant ( $P = 0.14$ ) while  
433 foraging time ( $n = 280$ ,  $\chi^2 = 5.31$ ,  $P = 0.021$ ) and reproductive structures per inflorescence were  
434 ( $n = 280$ ,  $\chi^2 = 8.02$ ,  $P = 0.004$ ). The relationship between susceptibility and reproductive  
435 structures per inflorescence (for which we have only one value per species) remained even if we  
436 used a grouped response with one susceptibility value per species ( $t = 2.3$ ,  $P = 0.0418$ ).

437 Using a similar approach for mean intensity as the response, four variables (reproductive  
438 structures per inflorescence, nectar production, corolla size, corolla shape) were significant as  
439 individual predictors in separate models (Table 2). In models that included reproductive  
440 structures per inflorescence as a predictor of mean intensity, none of the other variables was  
441 significant as an additional predictor ( $P > 0.3$  for all three), while reproductive structures per  
442 inflorescence was significant in all cases ( $F > 5.2$ ,  $P < 0.025$  for all). The final trait-based model  
443 for mean intensity thus had reproductive structures per inflorescence as the only covariate; bees  
444 that became infected after visiting plant species with more reproductive structures per  
445 inflorescence had higher mean *Crithidia* loads ( $n=194$ ,  $F=11.71$ ,  $P<0.001$ ; Fig. 2D). The  
446 relationship between intensity and reproductive structures per inflorescence remained even if we  
447 used a grouped response with one intensity value per species ( $t = 3.7$ ,  $P = 0.0035$ ).

448

#### 449 *Species vs. traits as predictors of susceptibility and intensity*

450 For susceptibility, the lowest AIC species-based model was a GLMM including species  
451 as a random effect, and bee size and total time foraging as fixed effects. The lowest AIC trait-  
452 based model was a GLM including reproductive structures per inflorescence, bee size and total  
453 time foraging as fixed effects. AIC for the trait-based model was somewhat lower ( $\Delta\text{AIC} =$

454 2.61). The two models made similar predictions ( $r = 0.83$  between the two models' fitted values),  
455 but the species-based model required more parameters (df = 7.12 for the species-based model,  
456 and 4 for the trait-based model). For prediction of mean intensity, the best species-based model  
457 included only species as a random effect, and the best trait-based model included only  
458 reproductive structures per inflorescence as a fixed effect. Comparing these models, AIC for the  
459 trait-based model was substantially lower ( $\Delta\text{AIC} = 5.85$ ), because the predictions were very  
460 similar ( $r = 0.83$  between the two models' fitted values) but the trait-based model had fewer  
461 parameters (df = 5.93 for the species-based model, and 3 for the trait-based model). Thus, for  
462 both susceptibility and intensity, traits and species identity had similar predictive power, and so a  
463 trait-based model is preferred due to greater simplicity. We also found no evidence of bias in the  
464 trait-based predictions (Appendix S4).

465

#### 466 Transmission trials manipulating flower number

467 For *Penstemon*, there were no significant covariates in model selection, and no significant  
468 treatment effect on mean *Crithidia* abundance (mean counts including zero values;  $n = 66$ ,  $\chi^2 =$   
469  $0.867$ ,  $P = 0.352$ ). In *Lythrum*, there was no significant treatment effect in a model with ( $n = 68$ ,  
470  $\chi^2 = 0.005$ ,  $P = 0.943$ ) or without significant covariates ( $n = 71$ ,  $\chi^2 = 0.042$ ,  $P = 0.847$ ). In  
471 *Monarda*, the effect of treatment was tested in a model including flowers probed as a fixed  
472 effect; both treatment and number of flowers probed were significant (treatment:  $n = 51$ ,  $\chi^2 =$   
473  $5.374$ ,  $P = 0.02$ ; number of flowers probed:  $n = 51$ ,  $\chi^2 = 6.24$ ,  $P = 0.01$ ). The estimated  
474 coefficient for the lower flower number treatment (-1.375) corresponds to a roughly 4-fold  
475 reduction in mean pathogen abundance in the low compared to high flower treatments, aligning

476 closely with raw mean abundance per bee in each treatment (mean  $\pm$  se: low:  $7.07 \pm 2.00$ ; high:  
477  $30.91 \pm 12.13$  cells per  $0.02 \mu\text{l}$ ).

478

## 479 **Discussion**

480 Overall, plant species differed fourfold in the mean abundance of pathogen cells  
481 established after a single bee foraging bout (Fig. 1A), with species explaining significant  
482 variation in both susceptibility and mean intensity (Figs 1B, 1C). These results complement  
483 earlier work which reported that the probability of *Crithidia* infection in *B. terrestris* workers  
484 differed on two plant species (Durrer and Schmid-Hempel 1994). Research more than twenty  
485 years later showed that *B. terrestris* and *Apis mellifera* can vector pathogens of both bee species  
486 via shared floral foraging, and the extent of vectoring differed between two plant species  
487 (Graystock et al. 2015). To our knowledge, these are the only previous studies asking whether  
488 plant species modulate pathogen acquisition among bees. Here we greatly extend the evidence of  
489 earlier work and, based upon the considerable variation in the effectiveness of different plant  
490 species to act as transmission hubs, suggest that plant community composition is likely to  
491 mediate bee-pathogen transmission dynamics. Future work should manipulate plant community  
492 composition in structured microcosms including bees and pathogen to assess longer-term effects.

493 In the transmission trials with 14 plant species, we found that models predicting  
494 susceptibility and mean intensity based on floral traits made similar in-sample predictions to  
495 models based on species identity. However, the trait-based models had lower AIC, and are  
496 therefore expected to have better out-of-sample predictive accuracy (i.e., more accurate forecasts  
497 of new observations), because the trait-based models required fewer parameters to fit the data.  
498 Moreover, only the trait-based models have any predictive power for species not represented in

499 the data set. These gains in parsimony and generalization are the potential benefits of trait-based  
500 models, which has inspired trait-based approaches to many different aspects of community  
501 ecology (e.g., Westoby and Wright 2006, Webb et al. 2010). Given enough data on all species in  
502 a community the situation would be reversed, because species always have idiosyncratic  
503 differences that cannot be fully captured by a list of traits. But in species-rich communities,  
504 getting “enough data” on ecological processes in each species may require prohibitive time or  
505 expense. Measuring relevant traits on all species, and using a subset to estimate trait-  
506 transmission relationships, may be far more feasible. If we can identify specific floral traits that  
507 shape pathogen transmission, these could be used to guide recommendations for pollinator-  
508 friendly habitat, within the context of other constraints such as phenology and providing diverse  
509 resources to support specialist as well as generalist pollinator species.

510 No measure of floral morphology significantly predicted transmission, which was  
511 surprising given the importance of floral morphology for pollen transfer by bees (e. g., Costa et  
512 al. 2017). However, we note that whenever possible we added inoculum within the corolla tube.  
513 Naturally foraging infected bees are likely to deposit feces on the corolla lip or outside the  
514 flower, and floral traits may shape risk or exposure by affecting where and how much bees  
515 defecate as well as trypanosome survival, although we note that in our study the number of  
516 inoculum drops probed had surprisingly little relationship with susceptibility or mean intensity of  
517 infection. The ultimate effect of floral traits on transmission will depend on whether their effects  
518 on risk amplify or counter their effects on susceptibility and intensity.

519 In our observational trials, species with more reproductive structures per inflorescence  
520 had greater transmission, measured as both susceptibility and mean intensity (Figure 2A, D).  
521 This was the most consistent floral trait that predicted transmission, more than the number of

522 flowers probed per trial or the amount of nectar each species produced. Reproductive structures  
523 per inflorescence even explained transmission more than the number of open flowers, which was  
524 surprising for two reasons. First, reproductive structures per inflorescence was measured at the  
525 species level (i.e., one value per plant species) while number of open flowers was counted for  
526 each trial. We would expect that a variable that was evaluated specifically for each trial would  
527 have more predictive power than a similar variable at the species level. Secondly, it is difficult to  
528 explain how reproductive structures per inflorescence (including buds and fruits) could mediate  
529 transmission more than the number of open flowers, given that bees only foraged on open  
530 flowers in our trials. If transmission occurs through spreading inoculum across all floral surfaces,  
531 then increased reproductive structures per inflorescence could provide more surface for spread  
532 via contact. Similarly, if more reproductive structures create more inflorescence complexity, this  
533 may affect micro-climates conducive to pathogen viability (such as increased humidity) or alter  
534 bee foraging behavior in ways that increase exposure. It is also possible that some underlying  
535 trait we did not consider is correlated with species-level variation in reproductive structures per  
536 inflorescence. For example, if plants that produced fewer flowers also produced longer-lasting  
537 flowers with more effective antimicrobial defense through nectar proteins or volatiles (e. g.,  
538 Thornburg et al. 2003), this could help explain our result. We measured floral longevity on a  
539 subset of our species and found no relationship with transmission (Tables 1 and 2), but have not  
540 exhaustively tested this hypothesis.

541       Because reproductive structures per inflorescence and number of open flowers are often  
542 tightly correlated, we manipulated the number of open flowers to determine whether this trait  
543 influences variation in bee host susceptibility and infection intensity within species of plants,  
544 although we acknowledge that the same trait may affect pathogen dynamics differently within



545 versus across species. We found only partial support for the hypothesis that the relationship  
546 between reproductive structures per inflorescence and transmission was due to an underlying  
547 correlation with number of open flowers. Flowers increased trypanosome pathogen abundance in  
548 bees nearly fourfold in *Monarda*, but there was no effect in *Lythrum* or *Penstemon*. Thus, the  
549 number of open flowers is unlikely to be the only mechanism explaining the relationship  
550 between reproductive structures per inflorescence and transmission.

551 Bee size and total time foraging were correlated with each other and with susceptibility in  
552 the observational transmission trials; smaller bees foraged for longer total time in the trials, and  
553 were more likely to become infected than larger bees. It is interesting that smaller bees foraged  
554 for longer periods but did not probe more flowers or inoculum drops in that time, suggesting that  
555 other mechanisms underlie the relationship between bee size and susceptibility. Small bees may  
556 be able to access more of the inoculum drops, particularly in plant species with narrow corollas.  
557 Furthermore, consuming the same amount of inoculum provides proportionally more pathogen  
558 cells per gram of bee tissue in a smaller bee, perhaps resulting in higher probability of infection.  
559 Finally, smaller bees could have less ability to resist infection. Although reduced food  
560 availability can decrease bee size (Schmid-Hempel and Schmid-Hempel 1998, Rotheray et al.  
561 2017) and affect immune function (Schmid-Hempel and Schmid-Hempel 1998, Brown et al.  
562 2003, Alaux et al. 2010), our bees were commercially reared and should not have been food  
563 stressed. Regardless of the mechanism, greater infection in smaller bees could have  
564 consequences for within-hive transmission since smaller bees are more often nurse bees while  
565 larger bees are foragers (Goulson 2010).

566 In conclusion, plant species varied widely in the transmission of *Crithidia* to *B.*  
567 *impatiens*, suggesting that plant species, through their influence on pathogen transmission, may

568 play an important role in shaping bee disease dynamics. Surprisingly, reproductive structures per  
569 inflorescence best predicted variation in transmission; floral size and morphology did not play  
570 significant roles. Across species, trait-based models were as good or better at predicting  
571 susceptibility and mean intensity based on AIC values, indicating the potential to use traits to  
572 select plant species that minimize pathogen spread, rather than requiring an evaluation of every  
573 plant species. Our manipulative experiment suggested that, within species, open flowers play a  
574 partial role explaining variation in transmission; such intraspecific variation may play important  
575 roles in plant-pollinator-pathogen dynamics. Given widespread investment in pollinator-friendly  
576 plantings to support pollinators, determining how plant species affect disease transmission is  
577 critical for recommending plant species that optimize pollinator health.

578

579 **Acknowledgements.** We thank two anonymous reviewers for feedback that improved the  
580 manuscript, Biobest for generously donating bee colonies, M. Ha as project manager and P.  
581 Anderson, O. Biller, S. J. Connon, D. Delaney, E. Duvall, S. Fogel, G. Killough-Hill, J.  
582 Giacomini, A. Hogeboom, A. Lee, G. LoCascio, E. Mann, L. Metz, N. Milano, R. Pasquale, K.  
583 Rothchild, D. Shaheen, T. Shaya, C. Shuja, T. Truong, A. Turkle, I. Weston and A. Zhao for  
584 field and lab assistance, C. Joyner for greenhouse support, and N. Woodard for field site  
585 management. This project was supported by NSF-DEB-1258096 (LSA and REI), by the National  
586 Institute of General Medical Sciences of the National Institutes of Health under Award Number  
587 R01GM122062 (LSA, SHM, REI, and SPE), by the National Research Initiative (NRI)  
588 Arthropod and Nematode Biology and Management Program of the USDA Cooperative State  
589 Research, Education, and Extension Service (CSREES) Grant no. USDA-AFRI 2013-02536  
590 (LSA and REI), and by UMass Amherst Biology Department Summer Research Fellowship and

591 the Summer Scholars program at the UMass Amherst Center for Agriculture, Food and the  
592 Environment (LSA and KMM). Any opinions, findings, and conclusions or recommendations  
593 expressed in this material are those of the authors and do not necessarily reflect the views of the  
594 National Science Foundation or the official views of the National Institutes of Health. LSA and  
595 REI conceived the ideas and designed methodology for the transmission trials; SPE and SHM  
596 conceived the comparison of trait-based and species-based models; LSA collected transmission  
597 and floral trait data for the observational study; LSA and KMM designed methodology for the  
598 manipulative experiment and KMM collected those data; PCS collected chemical data that were  
599 ultimately omitted from the manuscript, and contributed related ideas and content; SPE and LSA  
600 analysed the data; LSA led the writing of the manuscript with contributions from all authors. All  
601 authors contributed critically to the drafts and gave final approval for publication.

602

### 603 **Literature Cited**

604 Alaux, C., F. Ducloz, D. Crauser, and Y. Le Conte. 2010. Diet effects on honeybee  
605 immunocompetence. *Biology Letters* **6**:562-565.

606 Brown, M. J. F., R. Schmid-Hempel, and P. Schmid-Hempel. 2003. Strong context-dependent  
607 virulence in a host-parasite system: reconciling genetic evidence with theory. *Journal of*  
608 *Animal Ecology* **72**:994-1002.

609 Cisarovsky, G., and P. Schmid-Hempel. 2014. Combining laboratory and field approaches to  
610 investigate the importance of flower nectar in the horizontal transmission of a bumblebee  
611 parasite. *Entomologia Experimentalis et Applicata* **152**:209-215.

- 612 Colla, S. R., M. C. Otterstatter, R. J. Gegear, and J. D. Thomson. 2006. Plight of the bumble bee:  
613 Pathogen spillover from commercial to wild populations. *Biological Conservation*  
614 **129**:461-467.
- 615 Costa, J., S. Castro, J. Loureiro, and S. C. H. Barrett. 2017. Experimental insights on Darwin's  
616 cross-promotion hypothesis in tristylous purple loosestrife (*Lythrum salicaria*). *American*  
617 *Journal of Botany* **104**:616-626.
- 618 Cronin, J., M. Welsh, M. Dekkers, S. Abercrombie, and C. Mitchell. 2010. Host physiological  
619 phenotype explains pathogen reservoir potential. *Ecology Letters* **13**:1221–1232.
- 620 Durrer, S., and P. Schmid-Hempel. 1994. Shared use of flowers leads to horizontal pathogen  
621 transmission. *Proceedings of the Royal Society of London Series B* **258**:299-302.
- 622 Fürst, M. A., D. P. McMahon, J. L. Osborne, R. J. Paxton, and M. J. F. Brown. 2014. Disease  
623 associations between honeybees and bumblebees as a threat to wild pollinators. *Nature*  
624 **506**:364+.
- 625 Gamboa, V., J. Ravoet, M. Brunain, G. Smaghe, I. Meeus, J. Figueroa, D. Riano, and D. C. de  
626 Graaf. 2015. Bee pathogens found in *Bombus atratus* from Colombia: A case study.  
627 *Journal of Invertebrate Pathology* **129**:36-39.
- 628 Gegear, R. J., M. C. Otterstatter, and J. D. Thomson. 2006. Bumble-bee foragers infected by a  
629 gut parasite have an impaired ability to utilize floral information. *Proceedings of the*  
630 *Royal Society B-Biological Sciences* **273**:1073-1078.
- 631 Gillespie, S. 2010. Factors affecting parasite prevalence among wild bumblebees. *Ecological*  
632 *Entomology* **35**:737-747.
- 633 Goulson, D. 2010. *Bumble bees: behaviour, ecology and conservation*. Oxford University Press,  
634 New York, New York.

- 635 Goulson, D., E. Nicholls, C. Botias, and E. L. Rotheray. 2015. Bee declines driven by combined  
636 stress from parasites, pesticides, and lack of flowers. *Science* **347**:1435-1439.
- 637 Goulson, D., S. O'Connor, and K. J. Park. 2017. The impacts of predators and parasites on wild  
638 bumblebee colonies. *Ecological Entomology*.
- 639 Graystock, P., E. J. Blane, Q. S. McFrederick, D. Goulson, and W. O. H. Hughes. 2016. Do  
640 managed bees drive parasite spread and emergence in wild bees? *International Journal for*  
641 *Parasitology Parasites and Wildlife* **5**:64-75.
- 642 Graystock, P., D. Goulson, and W. O. H. Hughes. 2015. Parasites in bloom: flowers aid dispersal  
643 and transmission of pollinator parasites within and between bee species. *Proceedings of*  
644 *the Royal Society B-Biological Sciences* **282**:20151371.
- 645 Han, B. A., J. P. Schmidt, S. E. Bowden, and J. M. Drake. 2015. Rodent reservoirs of future  
646 zoonotic diseases. *Proceedings of the National Academy of Sciences, USA* **112**:7039-  
647 7044.
- 648 Harder, L. D. 1982. Measurement and estimation of functional proboscis length in bumblebees  
649 (Hymenoptera: Apidae). *Canadian Journal Of Zoology* **60**:1073-1079.
- 650 Koch, H., M. J. F. brown, and P. C. Stevenson. 2017. The role of disease in bee foraging  
651 ecology. *Current Opinion in Insect Science* **21**:60-67.
- 652 McArt, S. H., H. Koch, R. E. Irwin, and L. S. Adler. 2014. Arranging the bouquet of disease:  
653 floral traits and the transmission of plant and animal pathogens. *Ecology Letters* **17**:624-  
654 636.
- 655 McMahon, D. P., M. A. Furst, J. Caspar, P. Theodorou, M. J. F. Brown, and R. J. Paxton. 2015.  
656 A sting in the spit: widespread cross-infection of multiple RNA viruses across wild and  
657 managed bees. *Journal of Animal Ecology* **84**:615-624.

- 658 Montgomery, B. R., and B. J. Rathcke. 2012. Effects of floral restrictiveness and stigma size on  
659 heterospecific pollen receipt in a prairie community. *Oecologia* **168**:449-458.
- 660 Otterstatter, M. C., and J. D. Thomson. 2006. Within-host dynamics of an intestinal pathogen of  
661 bumble bees. *Parasitology* **133**:749-761.
- 662 Paull, S. H., S. Song, K. M. McClure, L. C. Sackett, A. M. Kilpatrick, and P. T. J. Johnson.  
663 2012. From superspreaders to disease hotspots: linking transmission across hosts and  
664 space. *Frontiers in Ecology and the Environment* **10**:75-82.
- 665 R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for  
666 Statistical Computing, Vienna, Austria.
- 667 Rotheray, E. L., J. L. Osborne, and D. Goulson. 2017. Quantifying the food requirements and  
668 effects of food stress on bumble bee colony development. *Journal of Apicultural  
669 Research* **56**:288-299.
- 670 Ruiz-Gonzalez, M. X., and M. J. F. Brown. 2006. Honey bee and bumblebee trypanosomatids:  
671 specificity and potential for transmission. *Ecological Entomology* **31**:616-622.
- 672 Ruiz-Gonzalez, M. X., J. Bryden, Y. Moret, C. Reber-Funk, P. Schmid-Hempel, and M. J. F.  
673 Brown. 2012. Dynamic transmission, host quality, and population structure in a multihost  
674 parasite of bumblebees. *Evolution* **66**:3053-3066.
- 675 Schmid-Hempel, R., and P. Schmid-Hempel. 1998. Colony performance and  
676 immunocompetence of a social insect, *Bombus terrestris*, in poor and variable  
677 environments. *Functional Ecology* **12**:22-30.
- 678 Shykoff, J. A., and P. Schmid-Hempel. 1991a. Incidence and effects of four parasites in natural  
679 populations of bumble bees in Switzerland *Apidologie* **22**:117-125.

- 680 Shykoff, J. A., and P. Schmid-Hempel. 1991b. Parasites delay worker reproduction in  
681 bumblebees: Consequences for eusociality. *Behavioral Ecology* **2**:242-248.
- 682 Singh, R., A. L. Levitt, E. G. Rajotte, E. C. Holmes, N. Ostiguy, D. Vanengelsdorp, W. I. Lipkin,  
683 C. W. Depamphilis, A. L. Toth, and D. L. Cox-Foster. 2010. RNA viruses in  
684 hymenopteran pollinators: Evidence of inter-taxa virus transmission via pollen and  
685 potential impact on non-*Apis* hymenopteran species. *Plos One* **5**.
- 686 Spira, W. M., A. Huq, Q. S. Ahmed, and Y. A. Saeed. 1981. Uptake of *Vibrio cholerae* biotype  
687 feltor from contaminated water by water hyacinth (*Eichornia crassipes*) *Applied and*  
688 *Environmental Microbiology* **42**:550-553.
- 689 Thornburg, R. W., C. Carter, A. Powell, R. Mittler, L. Rizhsky, and H. T. Horner. 2003. A major  
690 function of the tobacco floral nectary is defense against microbial attack. *Plant*  
691 *Systematics and Evolution* **238**:211-218.
- 692 Webb, C. T., J. A. Hoeting, G. M. Ames, M. I. Pyne, and N. L. Poff. 2010. A structured and  
693 dynamic framework to advance traits-based theory and prediction in ecology. *Ecology*  
694 *Letters* **13**:267-283.
- 695 Westoby, M., and I. J. Wright. 2006. Land-plant ecology on the basis of functional traits. *Trends*  
696 *in Ecology & Evolution* **21**:261-268.
- 697 Wood, S. N. 2006. *Generalized additive models: An introduction with R*. Chapman and  
698 Hall/CRC, Boca Raton, FL.

699

700

701 **Table 1.** Analysis of bee susceptibility to infection as a function of species identity, inoculum  
 702 source colony, floral traits, bee traits, and bee foraging behavior, using generalized linear models  
 703 with each focal variable as the one covariate (see text for details). Only the Species model  
 704 includes data on *Helianthus*. Source file: SpeciesTraitsAndSusceptibility.R and scripts that it  
 705 sources; Dryad repository.

Variable	p value <sup>1</sup>	$\chi^2$	n	Coefficient
Species	0.029	10.194	351	---
Inoculum Source	0.675	3.160	222	---
Nectar Volume	0.558	0.343	298	-0.057
Number of Open Flowers	0.177	1.821	300	0.005
Corolla Size	0.475	0.510	311	-0.006
Corolla Shape	0.140	2.179	311	-0.052
Repro. Structures per Infl.	0.008	7.071	311	0.005
Floral Longevity	0.271	1.213	193	0.073
Bee Size	0.034	4.518	293	-0.906
Total Time Foraging (min)	0.077	3.118	298	0.050
Number of Flowers Probed	0.779	0.079	299	0.002
Number of Inoc. Drops Probed	0.777	0.081	298	0.008

706 <sup>1</sup> *p*-values were obtained from `summary.gam` (for Species, fitted as a random effect) or `drop1`  
 707 with `test= "Chisq"` (all others, fitted as fixed effects),  $\chi^2$  is the value of the test statistic which has  
 708 an approximately chi-square distribution, *n* is the sample size and coefficient is the coefficient of  
 709 the focal variable in the linear predictor.



710 **Table 2.** Analysis of mean intensity (log of positive pathogen counts) as a function of species  
 711 identity, inoculum source colony, floral traits, bee traits, and bee foraging behavior, using linear  
 712 models with each focal variable as the one covariate (see text for details). Only the Species  
 713 model includes data on *Helianthus*. Source file: SpeciesTraitsAndIntensity.R and scripts that it  
 714 sources; Dryad repository.

Variable	p value <sup>1</sup>	F	n	Coefficient
Species	0.048	0.574	215	---
Inoculum source	0.682	0.624	142	---
Nectar Production	0.021	5.432	185	-0.181
Number of Open Flowers	0.201	1.648	189	0.003
Corolla Size	0.013	6.270	194	-0.017
Corolla Shape	0.031	4.739	194	-0.056
Repro. Structures per Infl.	0.001	11.709	194	0.004
Floral Longevity	0.318	1.006	119	-0.049
Bee size	0.701	0.148	183	-0.121
Total Time Foraging (min)	0.158	2.011	188	-0.028
Number of Flowers Probed	0.412	0.676	189	0.003
Number of Inoc. Drops Probed	0.214	1.555	189	0.023

715 <sup>1</sup> Table entries are as in Table 1, except that *p*-values are based on an F statistic.

716 **Figure Legends**

717

718 **Fig. 1.** A) Mean *Crithidia* cell count (in a 2  $\mu$ l sample) of bees foraging at different plant species  
719 provided with the same inoculum, including both zero and positive counts (mean abundance).

720 Error bars are  $\pm 1$  s.e.m.; numbers in bars are the sample size. Species acronyms begin with the

721 first three letters of the genus. B) Susceptibility, the fraction of trials on each plant species for

722 which the pathogen count was positive. Error bars are binomial standard errors on the fraction of

723 positive counts. C) Mean intensity, the mean of positive cell counts. Error bars are  $\pm 1$  s.e.m.

724 Figure generated by R script `SusceptibilityAndIntensityPlots.R`; Dryad repository.

725

726 **Fig. 2.** Relationships between traits and components of pathogen transmission that were

727 statistically significant in the transmission trials using 14 plant species. A) Mean susceptibility

728 (over all trials using a particular species) versus reproductive structures per inflorescence

729 (estimated mean for the species). B), C) Susceptibility in each trial irrespective of flower species

730 (0=not infected, 1=infected; values jittered to separate points) as a function of B) bee size,

731 estimated by the length of the wing radial cell, and C) total time foraging by the bee. D) Mean

732 intensity (mean of all log-transformed positive pathogen cell counts for each species) as a

733 function of reproductive structures per inflorescence (estimated mean for the species). The


734 dashed lines in each panel are regressions fitted to the plotted points (linear regression in panels

735 A and D, logistic regression in B and C). Figure generated by R script

736 `SusceptibilityAndIntensityPlots.R`; Dryad repository.

737


738 Figure 1



739

740

741 Figure 2



742

Disease where you dine: Plant species and floral traits associated with pathogen transmission in bumble bees

Lynn S. Adler<sup>1\*</sup>, Kristen M. Michaud<sup>1</sup>, Stephen P. Ellner<sup>2</sup>, Scott H. McArt<sup>3</sup>, Philip C. Stevenson<sup>4,5</sup>, and Rebecca E. Irwin

**Appendix S1. Plant source and propagation methods**

In general, plants were grown from seed, purchased from nurseries and transplanted to the field site, or collected from wild field sites. All dates are in 2014. In all cases we used Fafard Growing Mix #2 (Fafard, Inc., Anderson, SC, USA) except that sunflower (*Helianthus annuus* Incredible) and *Echium vulgare* were germinated in Fafard Super-fine germinating plug mix. Plants grown from seed were germinated in a propagation room with natural light (24°C day & night, with misting from dawn to dusk every 12 minutes for 15 seconds) and then transferred to a greenhouse with natural light (26.7°C day and 18.3°C night; late July and August : 22.8°C day and 19.4°C night) before transplanting to the field site.

Table S1. Plant source and propagation methods.

Species	Abbreviation	Common name	Source	Propagation Method	Date in field	Sample size for trait measures
<i>Antirrhinum majus</i> 'Twinny Peach' and 'La Belle White'	ANT	snapdragon	Andrews Greenhouse; Hadley Garden Center	purchased from nurseries and directly transplanted	June 4	25
<i>Asclepias incarnata</i>	ASC	milkweed	Collected from the field: 42°23'0.04"N	inflorescences were bagged and collected from field sites	N/A	30

			72°32'16.93"W			
<i>Digitalis purpurea</i> ; 'Foxy' and lavender 'Camelot'	DIG	foxglove	Andrews Greenhouse; Lowes Home Improvement	purchased from nurseries and directly transplanted	June 4 and 9	22
<i>Eupatorium perfoliatum</i>	EUP	common boneset	New England Wetland Plants	purchased from nursery and directly transplanted	June 4	30
<i>Helianthus annuus</i> 'Incredible'	HEL	sunflower	Burpee, from Home Depot	plants seeds in 50-plug trays on June 3, transplanted to 1 gal pots for the field	June 16	31
<i>Impatiens capensis</i>	IMP	jewelweed	Collected from the field: 42°35'18.3"N 72°52'39.5" W	Inflorescences were bagged and collected directly from the field for use in trials	N/A	30
<i>Linaria vulgaris</i>	LIN	butter-and-eggs	seeds collected from the field: 42°17'3.79"N 72°31'55.78"W	seeds planted directly in pots using native soil buried at the field site to prevent spread	June 20	30
<i>Lobelia siphilitica</i>	LOB	blue lobelia	New England Wetland Plants; Ion Exchange	purchased from nurseries; planted in 4" pots in greenhouse or else directly transplanted	June 4	32
<i>Lythrum salicaria</i>	LYT	purple loosestrife	collected from the field: 42°38'53.8" N, 72°53'76.5" W	dug from the field and transplanted, plants were buried in 5 gallon pots to prevent spread	June 12	30
<i>Monarda didyma</i> 'Pink Lace'	MON	bee balm	Pioneer Gardens	purchased from nursery and directly transplanted	June 10, 11 and 12	23
<i>Penstemon digitalis</i>	PEN	penstemon	New England Wetland Plants; Prairie Moon Nursery	purchased from nurseries; planted in 4" pots in greenhouse or directly transplanted	June 4	38
<i>Solidago canadensis</i>	SOLD	goldenrod	collected from	dug from the field and	June 9	30

			the field: 42°47'51.8" N 72°58'23.2" W	transplanted		
<i>Thymus vulgaris</i>	THYG	thyme (German variety)	Lowes Home Improvement	purchased from nursery and directly transplanted	June 16	30
<i>Verbascum thapsus</i>	VER	common mullein	Pioneer Gardens or collected from the field: 42°44'52.1"N 72°47'30.5" W and 42°24'51.03"N 72°30'30.77"W and 42°28'45.53"N 72°34'46.06"W	purchased from nursery and directly transplanted; additional inflorescences were bagged and collected from field sites	June 10, 13 and 16	26

## Disease where you dine: Plant species and floral traits associated with pathogen transmission in bumble bees

Lynn S. Adler <sup>1\*</sup>, Kristen M. Michaud <sup>1</sup>, Stephen P. Ellner <sup>2</sup>, Scott H. McArt <sup>3</sup>, Philip C. Stevenson <sup>4,5</sup>, and Rebecca E. Irwin


### **Appendix S2. Floral trait measurements**

The following images depict what we measured as corolla length and width for each species included in transmission trials. We provide these images because the choice of what to use for these traits wasn't always clear, particularly for species with highly zygomorphic flowers and Asteraceae species.


In all figures below, 'CL' indicates corolla length and 'CW' indicates corolla width. We also measured petal length and width on most species, indicated by 'PL' and 'PW' respectively, but ultimately did not include these measures in analyses because they were not relevant for all species.

Photo credits: Melissa Ha







*Antirrhinum majus*



*Asclepias  
incarnata*




*Digitalis purpurea*





*Eupatorium perfoliatum*


DD (disk diameter)




CW



*Helianthus annuus*





CL and CL2 were added to  
calculate final CL




*Impatiens capensis*

CL and CL2 were added to calculate final CL


*Linaria vulgaris*






*Lobelia siphilitica*








*Lythrum salicaria*



*Monarda didyma*




*Penstemon digitalis*



*Solidago canadensis*


CW



CL



*Thymus vulgaris*



*Verbascum thapsus*

1 Disease where you dine: Plant species and floral traits associated with pathogen transmission in  
2 bumble bees

3 Lynn S. Adler<sup>1\*</sup>, Kristen M. Michaud<sup>1</sup>, Stephen P. Ellner<sup>2</sup>, Scott H. McArt<sup>3</sup>, Philip C.  
4 Stevenson<sup>4,5</sup>, and Rebecca E. Irwin

## 5 **Appendix S4. Statistical analysis details**

### 6 Transmission trials across 14 plant species

7 *Justification of analyzing susceptibility and intensity separately, instead of one response*  
8 *with negative binomial regression.* The counts for each flower species were highly zero-inflated  
9 relative to a Poisson distribution with equal mean (17% to 55% zero counts, vs. <1% zero counts  
10 expected from Poisson distributions). Negative binomial regression is often used for such zero-  
11 inflated count data, but was not suitable for our data because the degree of zero-inflation varied  
12 substantially among species. In a negative binomial regression model, flower species with a  
13 higher mean intensity (i.e., higher mean of positive counts) would also have higher susceptibility  
14 (i.e., higher fraction of non-zero counts), but susceptibility and mean intensity were only weakly  
15 correlated (Pearson correlation coefficient  $r = 0.39$ ,  $p = 0.18$ ; Fig. S1). We therefore analyzed  
16 susceptibility and mean intensity as two separate components of pathogen transmission to bees.

17 *Helianthus as an outlier for floral traits and foraging behavior.* The distinctive floral  
18 architecture of *Helianthus* made it an outlier with respect to several floral traits, and resulted in  
19 very different foraging behavior. Bees on *Helianthus* probed over 400 disc flowers during a trial,  
20 more than twice the maximum number on any other species, and probed disc flowers and  
21 inoculum drops more than 5 and 10 times faster than the maximum rate of any other species,  
22 respectively. Several statistically significant apparent associations between traits and  
23 susceptibility or mean intensity were driven by a few trials with exceptionally active bees

24 foraging on *Helianthus*. Our analyses of trait-dependent transmission (susceptibility or mean  
25 intensity) therefore omitted *Helianthus*, but *Helianthus* was included in analyses that assessed  
26 species differences in transmission without considering floral traits.

27 *Evaluating bias in predictions of trait-based models.* AIC evaluates only the magnitude  
28 of prediction errors. To assess whether predictions of traits-based models might be biased, we  
29 computed the predicted susceptibility and mean intensity for each trial using the final trait-based  
30 models. We averaged those predictions to obtain predicted susceptibility (Fig. S2A) and mean  
31 intensity (Fig. S2B) for each species, which can be compared to the observed susceptibility and  
32 mean intensity. Linear regressions through the plots of observed vs. predicted values (solid black  
33 line), were nearly identical to the 1:1 lines (dashed red line), so there is no evidence of bias in the  
34 traits-based predictions, either upward, downward, or towards the mean for all species.

#### 35 36 Transmission trials manipulating flower number

37 *Discarded data.* Bees were discarded if they exhibited unusual foraging behavior (e. g.,  
38 difficulty flying) or died prior to dissection. Two bees with abnormally high *Crithidia* counts  
39 (>200 cells/ 0.02  $\mu$ L; one bee each from the *Lythrum* low flower and *Monarda* high flower  
40 treatments) were considered outliers and discarded prior to analysis. In total, 1, 2 and 5 bees  
41 were discarded from *Penstemon*, *Monarda* and *Lythrum* trials.

42 *Adequacy of negative binomial model.* The adequacy of the negative binomial model  
43 including all significant covariates for each species was tested by computing the Kolmogorov-  
44 Smirnov distance between the experimental data and the fitted negative binomial distributions  
45 for counts (using the R function `ks.test`), and doing the same for 500 artificial data sets simulated  
46 from the fitted distributions, identical in size and structure to the experimental data (generated by




47 the simulate function for glm.nb fits). For each species, the K-S distance of the experimental data  
48 was below the median K-S distance for the 500 artificial data sets, hence there is no evidence  
49 that the data for any species depart from the fitted negative binomial model.

50 *Model selection.* For *Penstemon*, no additional covariates predicted pathogen count ( $p >$   
51  $0.35$ ,  $\chi^2 < 0.67$ ,  $n = 65$  or  $66$ ). For *Lythrum*, trial time ( $p = 0.035$ ,  $\chi^2 = 4.46$ ,  $n = 68$ ) and minutes  
52 to trial ( $p = 0.044$ ,  $\chi^2 = 4.061$ ,  $n = 69$ ) were significant additional covariates. A second screening  
53 including those covariates as fixed effects and adding other covariates one at a time found that no  
54 other covariates were significant ( $p > 0.3$ ,  $\chi^2 < 0.88$ ,  $n = 67$  or  $68$ ). The presence of a treatment  
55 effect was therefore tested in a model with trial time and minutes to trial. For *Monarda*,  
56 treatment ( $p = 0.014$ ,  $\chi^2 = 6.029$ ,  $n = 51$ ) and number of flowers probed ( $p = 0.002$ ,  $\chi^2 = 9.841$ ,  $n$   
57  $= 51$ ) were significant as predictors so a second screening was done with those as fixed effects  
58 and other covariates added one at a time; none of the other covariates were significant predictors  
59 ( $p > 0.2$ ,  $\chi^2 < 1.6$ ,  $n = 50$  or  $51$ ). The effect of treatment was therefore tested in a model including  
60 flowers probed as a fixed effect.

61


62 **Figure S1.** Association between susceptibility (fraction of bees infected) and mean intensity  
63 (counts in infected bees) across flower species. Species acronyms begin with the first 3 letters of  
64 the genus. Solid circles are at the point estimates of susceptibility and intensity for each species.  
65 Cross-species Pearson correlation between susceptibility and intensity was  $r = 0.41$  ( $p = 0.15$ );  
66 with *Helianthus* removed this becomes  $r = 0.39$  ( $p = 0.18$ ). Source file:  
67 TransmissionAndIntensityPlots.R



68

69

70 **Fig. S2.** Plots of A) observed susceptibility (fraction of bees infected) and B) observed mean  
 71 intensity (counts in infected bees), for each species, versus predictions from the final traits-based  
 72 model. In each panel, the solid black line is the regression line fitted to the plotted species-  
 73 specific values (for observed and predicted susceptibility and mean intensity, respectively, in  
 74 panels A and B) and the dashed red line is the 1:1 line). The regression line would coincide with  
 75 the 1:1 line if predictions are unbiased. Source files: SpeciesTraitsAndSusceptibility.R,  
 76 SpeciesTraitsAndIntensity.R



77

78

79

80