

1        **Comparison of *in vitro* antibacterial activity of streptomycin-diclofenac loaded**  
2        **composite biomaterial dressings with commercial silver based antimicrobial wound**  
3        **dressings**

4        Harshavardhan V. Pawar<sup>1</sup>, John Tetteh<sup>1</sup>, Philip Debrah<sup>2</sup>, Joshua. S. Boateng<sup>1\*</sup>

5        <sup>1</sup>Department of Pharmaceutical, Chemical & Environmental Sciences, School of Science,  
6        University of Greenwich at Medway, Central Avenue, Chatham Maritime, ME4 4TB,  
7        Kent, UK.

8        <sup>2</sup>Department of Pharmaceutics and Microbiology, School of Pharmacy, College of Health  
9        Sciences, University of Ghana, Legon, P.O. Box LG43, Legon, Accra, Ghana.

10  
11        \* Corresponding author: Joshua S. Boateng

12        Tel: + 44 (0) 208 331 8980,

13        Fax: +44 (0) 208 331 9805

14        E-mail: J.S.Boateng@gre.ac.uk; joshboat40@gmail.com

15  
16        **Declarations of interest:** none

20 **Abstract**

21 Infected chronic wounds heal slowly, exhibiting prolonged inflammation, biofilm formation,  
22 bacterial resistance, high exudate and ineffectiveness of systemic antimicrobials. Composite  
23 dressings (films and wafers) comprising polyox/carrageenan (POL-CAR) and  
24 polyox/sodium alginate (POL-SA), loaded with diclofenac (DLF) and streptomycin (STP)  
25 were formulated and tested for antibacterial activity against  $2 \times 10^5$  CFU/mL of *Escherichia*  
26 *coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* representing infected chronic  
27 wounds and compared with marketed silver dressings. Minimum inhibitory concentration  
28 (MIC) showed higher values for DLF than STP due to non-conventional antibacterial  
29 activity of DLF. The DLF and STP loaded dressings were highly effective against *E. coli*,  
30 *P. aeruginosa* and *S. aureus*. POL-SA dressings were more effective against the three types  
31 of bacteria compared to POL-CAR formulations, whilst the DLF and STP loaded dressings  
32 showed greater antibacterial activity than the silver-based dressings. The films, showed  
33 greater antibacterial efficacy than both wafers and silver dressings. STP and DLF can act  
34 synergistically not only to kill the bacteria but also prevent their resistance and biofilm  
35 formation compared to silver dressings, whilst reducing chronic inflammation associated  
36 with infection.

37

38 **Keywords:** Antimicrobial dressing; bacterial infection; carrageenan; chronic wound;  
39 diclofenac; silver dressing; sodium alginate; streptomycin.

40

## 41 **1 Introduction**

42 A wound is an interruption in the defensive role of the skin in protecting against  
43 harmful environmental agents [1]. Injury evokes wound healing comprising distinct phases  
44 (haemostasis, inflammation, proliferation, migration and maturation) involving biochemical,  
45 and molecular events that work sequentially towards tissue regeneration [2]. However,  
46 wounds can get contaminated by microorganisms, especially during the proliferation stage  
47 leading to infection. Persistent infection impairs wound healing causing repeating  
48 inflammatory cycle, resulting in chronic wounds [3,4]. Prevention and control of infection  
49 have been identified as essential aspects of wound management [5]. Effective management  
50 requires reducing exogenous microbial contamination, debridement, using appropriate  
51 dressing(s) and administration of topical and systemic broad-spectrum antimicrobial agents  
52 [6]. Topical agents such as povidone iodine and chlorhexidine acid are commonly employed,  
53 though their use is currently restricted to wound cleansing and skin swabs before surgical  
54 incisions [1]. However, antibiotics have high specificity against infection and ultimately  
55 improve wound healing at low concentrations [1,7]. Various commercial dressings have  
56 been developed that release silver to prevent wound infections both *in vitro* [8] and *in vivo*  
57 [9]. The emergence of microbial resistance has resulted in the need for more effective  
58 treatments for wound infections [1]. Further, systemic antibiotic treatment is difficult in  
59 chronic wounds such as diabetic foot ulcers due to poor blood circulation at the extremities  
60 of diabetics [6].

61 Chronic wound infection also causes pain, excessive exudation and patient  
62 discomfort and is a major source of cross-infection particularly antibiotic-resistant species.  
63 Burns for example provide a protein-rich environment, favourable for microbial colonization  
64 [10]. Most infected wounds involve *Staphylococcus aureus*, *Pseudomonas aeruginosa*,

65 *Streptococci* and *Escherichia coli*. *S. aureus* is considered a challenging microorganism in  
66 wound infections [6] due to its ability to develop resistance against first line antibiotics.

67 Streptomycin (STP) has been used to treat wound infections [11] and for reducing  
68 infection before skin grafting [12]. It's reported that diclofenac (DLF) has antibacterial  
69 activity and acts synergistically with STP against *Mycobacterium tuberculosis* after systemic  
70 administration [13]. Systemic STP in combination with DLF demonstrated synergistic  
71 activity against 45 different strains of mycobacteria [14,15].

72 This paper reports on the evaluation of antibacterial activity of STP and DLF loaded film  
73 and wafer dressings against *S. aureus*, *E. coli* and *P. aeruginosa*. Minimum inhibitory  
74 concentration (MIC) of STP and DLF in the dressings and *in vitro* antibacterial efficacy  
75 (zone of inhibition) against the three microorganisms were evaluated using disk diffusion  
76 assay and compared with three commercial silver containing dressings. To the best of our  
77 knowledge, this is the first study comparing the antibacterial performance of streptomycin-  
78 diclofenac loaded medicated POL-CAR and POL-SA dressings with commercial silver  
79 loaded antimicrobial dressings for their antibacterial performance.

80

## 81 **2 Methods**

82

### 83 **2.1 Materials**

84 (Polyox™ WSR 301 ≈4000 kDa) was a gift from Colorcon Ltd (Dartford, UK), κ-  
85 carrageenan (Gelcarin GP 812) was from IMCD Ltd (Sutton, UK), Aquacel® Ag  
86 (ConvaTech, Ltd.), Melgisorb® Ag (Mölnlycke Health Care, Ltd.) were gifted by the  
87 manufacturers and Allevyn® Ag (Smith and Nephew, Ltd) obtained from a local pharmacy.  
88 Nutrient agar and nutrient broth were purchased from Oxoid, UK. Diclofenac sodium,  
89 streptomycin sulphate, glycerol, phosphate buffered saline (PBS) tablets, were purchased

90 from Sigma-Aldrich, (Gillingham, UK). Sodium alginate was purchased from Fisher  
91 Scientific (Loughborough, UK). National Collection of Type Culture (NCTC) strains of *S.*  
92 *aureus* (A 29213), *E. coli* (DTCC 25922) and *P. aeruginosa* (A 10145), were used for  
93 microbiological assays.

94

## 95 **2.2 Preparation of composite polymer based dressings**

96 Composite films and wafers (Table 1) were prepared as previously reported [16,17].  
97 Polymeric gels of POL-CAR and POL-SA gels were prepared as previously reported  
98 [16,17]. In brief, blends of POL with CAR and POL with SA (weight ratio of 75/25 and  
99 50/50 respectively) yielding 1% w/w of total polymer weight were prepared by stirring on a  
100 magnetic stirrer at 70°C to form a uniform gel (POL-CAR-BLK and POL-SA-BLK). The  
101 composition of the polymers, drugs used for the preparation of gels are summarised in Tables  
102 2 and 3. DL gels of POL-SA and POL-CAR were prepared with 4 ml ethanolic solution of  
103 DLF containing 100 mg and 250 mg of the drug to achieve 10% w/w for POL-SA gel and  
104 for POL-CAR to achieve 25% w/w of DLF in the polymeric gel respectively. These gels  
105 were subsequently cooled to 40°C with constant stirring. Similarly, a 4 ml aqueous solution  
106 containing 250 mg and 300 mg of STP was subsequently added to achieve a final STP  
107 concentration of 25% w/w (POL-SA) and 30% w/w (POL-CAR) in the DL gels.

108 To obtain films the solutions (25g) were poured into Petri dishes (diameter 90 mm)  
109 and dried in an oven at 40°C for 18h, to obtain the films, while unplasticised polymeric  
110 solutions (10g) were freeze-dried to obtain wafers. To obtain the wafers, 10 gm of each  
111 homogeneous gel was transferred into 6 well moulds (diameter 35 mm) (Thermo-Fisher  
112 Scientific Nunc, Leicestershire UK), placed in a Virtis Advantage XL 70 freeze dryer  
113 (Biopharma Process Systems, Winchester, UK) and lyophilised using the automated  
114 lyophilisation cycle. This involved initially cooling samples from room temperature to -5°C

115 and then -50°C over a period of 10 h (at 200 mTorr). An annealing step at -25°C for 2 h was  
116 applied based on the preliminary DSC studies and its effect on the different formulations  
117 was investigated. The frozen samples were then heated in a series of thermal steps to -25°C  
118 under vacuum (20-50 mTorr) over a 24 h period. Secondary drying of the wafers was carried  
119 out at 20°C (10 mTorr) for 7 h.

120

### 121 **2.3 Bacterial sample preparation**

122 Fresh broth cultures were prepared as reported by Labovitiadi *et al.*, [18] by  
123 transferring a single bead unit into 10mL of nutrient broth and incubating for 24h. A loop  
124 full of bacterial culture was streaked onto nutrient agar plate and incubated at 37°C for 24h  
125 to yield separate colonies. Overnight bacterial cultures were centrifuged at 4000 rpm for 10  
126 min in an Accuspin 1 centrifuge (Fisher Scientific, UK), supernatant discarded and pellets  
127 suspended in 20mL of simulated wound fluid (SWF) [16]. This process was repeated twice  
128 and final pellets re-suspended in 5mL SWF, followed by two fold dilutions in SWF.  
129 Bacterial density was determined by measuring the dilute suspension at 500nm to yield the  
130 required density of  $2 \times 10^5$ CFU/mL [18].

131

### 132 **2.4 Minimum inhibitory concentration (MIC) of STP and DLF**

133 The MIC for STP and DLF was evaluated as previously reported [19]. Briefly, three  
134 different stock solutions for each drug were prepared (Table 4) and STP required to obtain  
135 10,000mg/L was calculated using equation 1. Antimicrobial susceptibilities of *S. aureus*, *E.*  
136 *coli* and *P. aeruginosa* were determined by establishing the MIC using a standard agar  
137 dilution method and 0.25-512mg/L calibration solutions of DLF and STP dilutions also  
138 prepared. 200µL of stock and diluted solutions (10,000mg/L, 1,000mg/L and 100mg/L  
139 respectively) were transferred into a Petri plate and 20mL of nutrient agar (stabilized at

140 45°C) added and mixed. The agar was allowed to set at room temperature and 0.1mL of  
141  $1 \times 10^5$  CFU/mL of *S. aureus*, *E. coli* and *P. aeruginosa* were spread on separate Petri plates.  
142 These plates were incubated at 37°C for 24h and ensuring that all microorganisms had grown  
143 on drug free control plate. MIC is the lowest concentration of antimicrobial at which there  
144 was no visible growth of organisms. Growth of one or two colonies or a fine film of growth  
145 was disregarded.

$$W = \frac{1000}{P} \times V \times C \text{ Eq.1}$$

146  
147  $W$  is the weight of actives (mg) dissolved in volume  $V$  (mL),  $C$  is final concentration  
148 of solution (multiples of 1,000mg/L),  $P$  (785 $\mu$ g/mg) is the potency provided by the  
149 manufacturer.

150

## 151 **2.5 *In vitro* antibacterial activity of antimicrobial films, wafers and marketed silver** 152 ***dressings***

153 The disk diffusion method was used for the assessment of the antibacterial activity  
154 of the DL films, wafers and commercial silver dressings. Solutions ( $2 \times 10^5$  CFU/mL) of each  
155 bacterial strain (*S. aureus*, *E. coli* and *P. aeruginosa*) was prepared as specified above  
156 (section 2.3) and 0.1mL of each strain spread separately on set nutrient agar media. The  
157 inoculated microorganisms were incubated at  $37 \pm 1^\circ\text{C}$  for 4h to initiate growth of  
158 microorganisms on the inoculated culture medium before placing the films, wafers and  
159 marketed dressings. The films and marketed silver dressings were cut into 2cm diameter disc  
160 shapes. However, due to difficulty of cutting thicker wafers into smaller discs, DL gels (2g)  
161 were free-dried in 2cm diameter containers to obtain the same diameter as the cut film discs.  
162 Further, circular Whatmann<sup>®</sup> paper discs (2cm diameter), each wetted with reference  
163 solutions (80 $\mu$ L) of STP and DLF at concentrations of 6mg/mL and 5mg/mL respectively  
164 were used as positive controls. Negative controls were BLK films and wafers (2cm diameter)

165 without any STP or DLF. The plates were then incubated at  $37\pm 1^{\circ}\text{C}$  for 24 h after which the  
166 end zones of inhibition (ZOI) in millimetres, formed on the medium ( $n = 3$ ), were measured.

167

## 168 **2.6 Statistical analysis**

169 Statistical data evaluation was performed using two tailed student t-test at 95%  
170 confidence interval (Graph Pad Prism 4 software) with  $p$  value  $< 0.05$  as the minimal level  
171 of significance.

172

## 173 **3 Results**

### 174 **3.1 MIC of STP and DLF**

175 The MIC of STP and DLF was determined for known densities ( $2 \times 10^5$  CFU/mL) of  
176 *S. aureus* *P. aeruginosa* *E. coli* commonly associated with infected chronic wounds. The  
177 MICs of STP for *S. aureus* and *E. coli* ranged from 4 - 8mg/L but ranged from 8 - 16mg/L  
178 for *P. aeruginosa*. MIC for DLF against *P. aeruginosa* was greater than 512mg/L and 256 -  
179 512mg/L for *E. coli* and *S. aureus* respectively.

180

### 181 **3.2 Antimicrobial activity of pure STP and DLF controls**

182 The ZOI of the STP and DLF positive controls for *S. aureus*, *P. aeruginosa* and *E.*  
183 *coli* are shown in Figure 1 (N and O). STP showed significantly ( $p<0.05$ ) lower ZOI  
184 ( $3.2\pm 0.1\text{mm}$ ) for *S. aureus* compared to *P. aeruginosa* and *E. coli*. The maximum ZOI of *P.*  
185 *aeruginosa* was  $4.1\pm 0.1\text{mm}$  which was lower compared to *E. coli* ( $4.6\pm 0.1\text{mm}$ ) and was  
186 statistically significant ( $p<0.05$ ). DLF did not show ZOI for *S. aureus*, *P. aeruginosa* and *E.*  
187 *coli* though there was no bacteria growing directly under the DLF disc (Figure 1, *E. coli*  
188 plate O) implying that their effectiveness alone as antibacterial may be limited application  
189 to infected wounds.



190

191 **3.3 Antibacterial activity of POL-CAR films ( $2 \times 10^5$  CFU/mL)**

192 Figures 2 (A) and 3(A, B and C) show ZOI of POL-CAR-DL and POL-CAR-DL-  
193 20% GLY films against *S. aureus*, *P. aeruginosa* and *E. coli*. There was a significant  
194 difference observed for all POL-CAR-DL films against strains of bacteria (compared to  
195 wafers and marketed dressing and DLF, STP discs. POL-CAR-DL and POL-CAR-DL-20%  
196 GLY films showed a smaller ZOI for *S. aureus* but increased for *P. aeruginosa* and *E. coli*.  
197 For *S. aureus* the ZOI for POL-CAR-DL and POL-CAR-DL-20%GLY films was  
198  $3.6 \pm 0.1$ mm and  $3.5 \pm 0.1$ mm respectively which was significantly ( $p < 0.05$ ) higher than pure  
199 STP ( $3.2 \pm 0.1$ mm). For *P. aeruginosa*, the observed ZOI was higher than *S. aureus* but less  
200 than *E. coli*. POL-CAR-DL and POL-CAR-DL-20%GLY films showed similar ZOI  
201 ( $4.3 \pm 0.1$ mm) for *P. aeruginosa* which was higher than the control STP ( $4.1 \pm 0.1$ mm),  
202 however, the difference was not statistically significant ( $p > 0.05$ ). The maximum ZOI of  
203 POL-CAR-DL and POL-CAR-DL-20%GLY films was  $4.8 \pm 0.2$ mm and  $4.7 \pm 0.1$ mm  
204 respectively, for *E. coli* which though higher than  $4.6 \pm 0.2$ mm for the control STP were not  
205 statistically significant ( $p > 0.05$ ).

206

207 **3.4 Antibacterial activity of POL-SA films ( $2 \times 10^5$  CFU/mL)**

208 Figures 2 (B) and 4 (D, E and F) show the ZOI of POL-SA-BLK, POL-SA-DL and  
209 POL-SA-DL-9% GLY) films for *S. aureus*, *P. aeruginosa* and *E. coli*. For *S. aureus*, the  
210 observed ZOI for POL-SA-DL and POL-SA-DL-9%GLY films was  $4.6 \pm 0.2$ mm and  
211  $4.1 \pm 0.2$ mm respectively which was significantly ( $p < 0.05$ ) higher compared to the STP  
212 ( $3.2 \pm 0.1$ mm) control. The ZOI increased from  $4.6 \pm 0.2$ mm (*S. aureus*) to  $4.8 \pm 0.2$ mm (*P.*  
213 *aeruginosa*) and  $5.0 \pm 0.2$ mm (*E. coli*) for POL-SA-DL films while for POL-SA-9%GLY

214 films it increased from  $4.1\pm 0.2\text{mm}$  (*S. aureus*) to  $5.1\pm 0.2\text{mm}$  (*P. aeruginosa*) and  
215  $5.5\pm 0.2\text{mm}$  (*E. coli*) respectively.

216

### 217 **3.5 Antibacterial activity of POL-CAR and POL-SA wafers ( $2 \times 10^5$ CFU/mL)**

218 Figures 2 (C) and 5 (G, H, I and J) show the ZOI of POL-CAR and POL-SA (BLK  
219 and DL) wafers for *S. aureus*, *P. aeruginosa* and *E. coli* bacterial strains. As was observed  
220 for the films, the BLK (no drug) wafers did not show any ZOIs against all three  
221 microorganisms (Figure 5, G and I). The ZOI of POL-CAR for *S. aureus* was  $3.1\pm 0.1\text{mm}$   
222 which increased to  $3.3\pm 0.1\text{mm}$  for POL-SA whereas STP had a value of  $3.2\pm 0.1\text{mm}$  which  
223 was not statistically significant ( $p>0.05$ ). For *P. aeruginosa*, the ZOI was higher than *S.*  
224 *aureus* but less than *E. coli*. POL-SA-DL and STP showed similar ZOI of  $4.1\pm 0.2\text{mm}$  which  
225 subsequently decreased for POL-CAR-DL ( $3.9\pm 0.1\text{mm}$ ). The maximum ZOI of POL-CAR-  
226 DL and POL-SA-DL wafers was respectively  $4.5\pm 0.1\text{mm}$  and  $4.6\pm 0.3\text{mm}$  for *E. coli*.

227

### 228 **3.6 Antimicrobial efficacy of marketed wound dressings ( $2 \times 10^5$ CFU/mL)**

229 Figures 2 (D) and 6 (K, L and M) show the ZOI of silver loaded marketed dressings  
230 (Table 5) (Aquacel<sup>®</sup> Ag, Melgisorb<sup>®</sup> Ag and Allevyn<sup>®</sup> Ag) for *S. aureus*, *P. aeruginosa* and  
231 *E. coli*. There were very small ZOIs observed for all three different strains of microorganisms  
232 in the presence of these marketed silver based dressings, though these bacteria were  
233 completely absent in the area directly underneath the dressing as shown in figure 6 inset (M,  
234 *S. aureus*). The ZOI for *S. aureus* was increased for Allevyn<sup>®</sup> Ag foam dressing ( $2.3 \pm$   
235  $0.1\text{mm}$ ) while all three marketed dressings showed a ZOI of  $2.0\pm 0.1\text{mm}$  for *P. aeruginosa*.  
236 The ZOI for *E. coli* was higher for Allevyn<sup>®</sup> Ag foam dressing ( $2.9\pm 0.0\text{mm}$ ) compared to  
237 Aquacel<sup>®</sup> Ag and Melgisorb<sup>®</sup> Ag ( $2.0\pm 0.0\text{mm}$ ).

## 238 **4. Discussion**

239 One of the overall objectives of the broad study was to compare the properties of dense  
240 dressings such as films to corresponding porous formulations such as freeze-dried wafers  
241 relative to commercial silver based dressings. Drying in an oven only yields non porous  
242 films and therefore it was important to freeze-dry other gels in a freeze-dryer. The reason  
243 for plasticising the films, was purely to improve the flexibility and ease of handling, to  
244 fulfil one of the key functional performance requirements for film dressings. The  
245 hypothesis for the comparison, was that the differences in physical properties (porosity),  
246 which are known to significantly affect rate of hydration and swelling, will also  
247 significantly affect the rate of drug diffusion out of the swollen gels and subsequently  
248 affect the degree of antibacterial efficacy. The hypothesis for the comparison, was that the  
249 differences in physical properties (porosity), which are known to significantly affect rate of  
250 hydration and swelling, will also significantly affect the rate of drug diffusion out of the  
251 swollen gels and subsequently affect the degree of antibacterial efficacy.

252 Ineffective control of wound infections caused by antibiotic resistant strains of  
253 pathogens has intensified the need to consider modifying current approaches including use  
254 of medicated dressings which can overcome resistance and reduce bacterial biofilm  
255 formation. This study assessed the *in vitro* antibacterial activity of composite films and  
256 wafers combining antibacterial (STP) and anti-inflammatory (DLF) drugs for targeting two  
257 phases of wound healing. The two drugs were also selected based on their reported  
258 synergistic antibacterial effect when administered systemically [14]. Many texts refer to  
259 bacterial bio-burden greater than  $10^5$  CFU/mL organisms per gram of tissue as a criterion  
260 for infection [3,6]. In this study we used  $2 \times 10^5$  CFU/mL of *S. aureus*, *P. aeruginosa* and *E.*  
261 *coli* to evaluate antimicrobial efficacy of DL film and wafer dressings and compared their  
262 performance against marketed silver dressings.

263 POL-CAR-BLK films did not show any zone against all three different  
264 microorganisms (Figure 3 A) implying that the observed antibacterial effect was solely due  
265 to the presence of STP and DLF. The formulated films, wafers and marketed dressings  
266 showed antibacterial efficacy against bacterial bio-burden  $2 \times 10^5$  CFU/mL of *S. aureus*, *P.*  
267 *aeruginosa* and *E. coli*. Both *P. aeruginosa* and *E. coli* are Gram-negative microorganisms  
268 and required a higher MIC of STP compared to *S. aureus*. This means STP is more effective  
269 against the Gram-positive microorganism *S. aureus* than the Gram-negative *E. coli* and *P.*  
270 *aeruginosa* which is interesting, given the fact the *S. aureus* and related species are a major  
271 cause of antibiotic resistance [14].

272 During the antibacterial study, the films and wafers swelled when placed on the  
273 highly water saturated agar gel under incubation, simulating a broken skin (wound) surface  
274 and this is to be expected. The swelling of the drug loaded polymeric dressings is an  
275 important characteristic as that is important to ensure ease of drug dissolution, diffusion out  
276 of the swollen gel and eventually release to reach the target bacterial organisms.

277 To kill the bacteria, STP and DLF must interact with the binding site, occupy a  
278 critical number of sites of the bacteria and remain there long enough to inhibit normal  
279 biochemical reactions [20]. It's been reported that antimicrobial activity is either  
280 concentration or time dependent [21,22]. Concentration dependent drugs include  
281 aminoglycosides (e.g. STP), whose ability to kill bacteria is dependent on the presence of  
282 high concentrations at the site of infection. At least a ratio of 10:1 is required for such  
283 concentration dependent antibiotics to effectively kill bacteria and prevent development of  
284 resistance [21,23,24,25]. On the other hand, drug concentrations above the MIC should  
285 remain for long periods of time at the site of infection in order to achieve antibacterial  
286 action [21,22].

287 In previous studies [13,14,15], it has been demonstrated that the concentrations  
288 required to kill *S. aureus* is higher than *P. aeruginosa* which is time dependent. DLF required  
289 higher concentrations to kill the bacteria that are beyond those clinically achievable with  
290 antibiotics, implying that DLF on its own could not effectively inhibit *P. aeruginosa* based  
291 infections. Dutta *et al.*, [14] previously demonstrated that when DLF is used *in vitro*, it  
292 showed higher MIC values compared to conventional antibiotic drugs such as STP but *in*  
293 *vivo*, the amount of DLF required to protect an animal from *Mycobacterium spp* was much  
294 lower. This suggests that DLF might be used as adjuvant to current to manage bacterial  
295 infections [13,14,15] as was done in this study.

296 For *S. aureus*, different ZOI were observed attributed to the rate of diffusion of STP  
297 and DLF (films and wafers) and silver (commercial) from the dressings. Both POL-CAR;  
298 POL-SA films had significantly higher ZOI suggesting a synergistic action between both  
299 drugs compared to each individual drug (refer to figure1). ZOI was ellipsoidal for POL-SA  
300 films due to the rapid initial swelling and disintegration of the polymer matrix and rapid  
301 diffusion of STP and DLF through the free flowing swollen gels (figure 4 E&F). Bajpai &  
302 Sharma [26] explained that the more rapid swelling of SA is due to the mannuronate block  
303 where  $\text{Ca}^{2+}$  binds to the poly gluconate units which starts to disintegrate the swollen matrix  
304 [26]. Differences in the ZOI of POL-CAR-DL and POL-SA-DL formulation could be related  
305 to the two different polymers (CAR and SA), their percentage ratios used and their different  
306 swelling mechanisms (surface wetting, hydration, hydrogel formation and erosion) [17]  
307 which subsequently affects rate of drug diffusion through the matrix and onto the bacterial  
308 colonies.

309 Maximum ZOI was observed for POL-SA-DL and POL-SA-DL-9%GLY films due  
310 to rapid swelling and subsequently rapid diffusion of both STP and DLF from the swollen  
311 matrix. This supports the swelling and drug release data from previous studies [16,17].

312 All the DL films showed greater antibacterial activity compared to wafers which was  
313 interesting. Wafers generally have a higher loading capacity, faster hydration and cumulative  
314 percent drug release compared to films due to their generally more porous nature [27].  
315 However, it was observed that higher drug loading in the wafers resulted in the formation of  
316 greater amounts of sodium sulphate which decreased the hydration capacity [16,17] of DL  
317 wafers subsequently affecting drug diffusion with a consequent decrease in ZOI compared  
318 to films but greater than the marketed dressings.

319 From a pharmaceutical perspective, these differences could be associated with the  
320 total amounts of polymer present in films and wafers which resulted in the different  
321 hydration rates and eventually different ZOIs. For example, the weights ranged from 22.1  
322 mg and 30.3 mg for POL-CAR DL films and wafers respectively. This was also true for the  
323 POL-SA DL films and wafers (17.9 mg and 24.6 mg for films and wafers respectively). It  
324 should be noted that though both formulations had similar diameters, their contents were  
325 different as the films were cut out directly from a bigger sheet due to difficulty of removing  
326 a film with small diameter whilst the wafers were cast directly into 2 cm diameter moulds  
327 due to ease of removal. It is very difficult to effectively cut a relatively thick wafer into  
328 circular discs without damaging the structure due to their soft and porous nature.

329 More interestingly, the formulated film and wafer dressings, showed greater  
330 antibacterial efficacy than marketed silver based antibacterial dressing which showed either  
331 lower or absence of ZOIs for all three different microorganisms even though the area directly  
332 under the discs showed no microbial growth. This may be due to two reasons: (i) the lower  
333 amounts of silver present in these dressings (Figure 6) relative to the combined  
334 concentrations of STP and DLF present in the composite films and wafers and (ii) most  
335 likely due to STP and DLF present in both films and wafers acting synergistically to kill the  
336 bacteria and potentially inhibiting biofilm formation and resistance of the bacteria. DLF

337 consists of a secondary amino group and a phenyl ring, both ortho positions of which are  
338 occupied by chlorine atoms. This causes an angle of torsion between the two aromatic rings,  
339 which presents structural similarities with phenothiazine and this is responsible for its  
340 antibacterial activity against microorganisms such as *E. coli*, *S. aureus* and *P. aeruginosa*  
341 [14,15]. DLF's antibacterial activity involves the inhibition of bacterial DNA synthesis  
342 whereas STP acts by binding to 30S ribosomal subunits in the microorganisms and  
343 disrupting the initiation and elongation steps in protein synthesis. On the other hand, silver  
344 in the presence of moisture, such as wound exudate, readily ionises to release silver ions  
345 ( $\text{Ag}^+$ ) which is involved in oxidation reactions by catalysing reactions between oxygen  
346 present in the cell and hydrogen from thiol groups. This results in disulphide bond formation,  
347 ultimately inhibiting cell function due to changes in protein structure, resulting in protein  
348 denaturation and enzyme inhibition [28]. The increased antibacterial activity of the film and  
349 wafer dressings suggests a potential application in chronic wound management.

350 Formulations administered for systemic use usually have to overcome the challenges  
351 to drug absorption, metabolism, distribution and elimination before the drug reaches the  
352 target sites for activity, hence such systemic formulations tend not to always have direct *in*  
353 *vitro-in vivo* correlations. For formulations such as wound dressings, intended for direct  
354 application, where the drug(s) are in direct contact with the target tissues, a high positive *in*  
355 *vitro – in vivo* correlation tends to exist due to minimal pharmacokinetic barriers.

356 Silver is a widely used anti-microbial agent effective against infection causative  
357 wound pathogens which are responsible for delayed wound healing and can be added to a  
358 range of composite dressings [29]. Silver containing wound dressings release silver ions  
359 which vary due to the different forms (silver sulfadiazine, ionic silver nanoparticles  
360 containing scaffolds, nanofiber containing silver nanoparticles, silver-containing activated  
361 carbon and fibres) and the amount of the silver present [1,30]. Although there are important

362 questions raised by Modak *et al* [31] in regards to the use of silver in infected wounds and  
363 formation of biofilms by the microorganisms, the versatile effect of silver carries a low risk  
364 of resistance even though some studies in burn wounds have shown bacterial resistance to  
365 silver sulfadiazine and silver nitrate by *Pseudomonas spp* [31]. Moreover, the antimicrobial  
366 effect of silver incorporated in a number of dressings depends on the release rate of silver  
367 ions which influences the overall antimicrobial effect [32]. In comparative antimicrobial  
368 efficacy studies, it was reported that certain types of methicillin resistant strains among *S.*  
369 *aureus*, *P. aeruginosa*, and *E. coli* were less sensitive to Urgotul SSD<sup>®</sup>, Bactigras<sup>®</sup>,  
370 Acticoat<sup>®</sup>, Askina Calgitrol Ag<sup>®</sup> and Aquacel Ag<sup>®</sup> antimicrobial dressings [33].  
371 Furthermore, *in vivo* silver can bind to proteins present in biofilms instead of binding to the  
372 bacterial cell walls, resulting in reduced antimicrobial effect against the bacteria [34].  
373 Another potential concern is that silver does not act specifically against bacteria but also acts  
374 on any host proteins. Therefore, if very few bacteria counts are present at the wound site,  
375 then the effect on host tissue is greater which could slow down healing [35]. Concentrations  
376 above 1mg/L (1 part per million) of silver reacts with wound exudate and could cause  
377 transient skin staining [36]. Li and co-authors suggested that bacterial resistance could be  
378 induced when low concentrations of silver were used [37]. There is therefore the possibility  
379 of these silver containing dressings inducing resistance from *S. aureus* and *P. aeruginosa*  
380 which are known to be able to form biofilms in an infected chronic wound environment [38].  
381 However, because there was absence of bacteria in the immediate application area beneath  
382 the marketed dressings, it implies the silvers dressing were effective to kill the bacteria in  
383 only the applied area of a wound and could also potentially limit or completely prevent  
384 infection from external sources.

385



386 **5 Conclusion**

387 Composite polymer based dressings containing STP and DLF appear to show  
388 significantly higher inhibition of the three bacterial strains compared to silver containing  
389 commercial dressings. STP can help to reduce bacterial infection by its known antimicrobial  
390 action and potentially in synergy with DLF while the latter can also help to reduce the  
391 swelling and pain associated with injury due to its anti-inflammatory action. However, these  
392 will, require further investigations in an *in vitro* cell culture (for cell viability and cell  
393 migration/proliferation) and *in vivo* animal study.

394

395 **Funding**

396 This research did not receive any specific grant from funding agencies in the public,  
397 commercial, or not-for-profit sectors.

398 **6**      **References**

- 399 [1] Boateng, J.S., Catanzano, O. 2015. Advanced Therapeutic Dressings for Effective  
400 Wound Healing--A Review. *Journal of Pharmaceutical Sciences*; 104, 3653-3680.
- 401 [2] Beldon, P. 2010. Basic science of wound healing. *Surgery (Oxford)*; 28, 409-412.
- 402 [3] White, R.J., Cutting, K., Kingsley, A. 2006. Topical antimicrobials in the control of  
403 wound bioburden. *Ostomy Wound Management*; 52, 26-30.
- 404 [4] Lipsky, B.A., Hoey, C. 2009. Topical Antimicrobial Therapy for Treating Chronic  
405 Wounds. *Clinical Infectious Diseases*; 49, 1541-1549.
- 406 [5] Kim, H., Makin, I., Skiba, J., Ho, A., Housler, G., Stojadinovic, A., Izadjoo, M.  
407 2014. Antibacterial efficacy testing of a bioelectric wound dressing against clinical  
408 wound pathogens. *Open Microbiology Journal*; 8, 15-21.
- 409 [6] Bowler, P.G., Duerden, B.I., Armstrong, D.G. 2001. Wound microbiology and  
410 associated approaches to wound management. *Clinical Microbiology Reviews*; 14,  
411 244-269.
- 412 [7] Ahmed, A., Getti, G., Boateng, JS. 2017. Ciprofloxacin-loaded calcium alginate  
413 wafers prepared by freeze-drying technique for potential healing of chronic diabetic  
414 foot ulcers. *Drug Delivery and Translational Research*; doi: 10.1007/s13346-017-  
415 0445-9.
- 416 [8] Boateng, J.S, Burgos-Amador, R., Okeke, O., Pawar, H. 2015. Composite alginate  
417 and gelatin based bio-polymeric wafers containing silver sulfadiazine for wound  
418 healing. *International Journal of Biological Macromolecules*; 79, 63-71.
- 419 [9] Ong, S.Y., Wu, J., Moochhala, S.M., Tan, M.-H., Lu, J. 2008. Development of a  
420 chitosan-based wound dressing with improved hemostatic and antimicrobial  
421 properties. *Biomaterials*; 29, 4323-4332.

- 422 [10] Church, D., Elsayed, S., Reid, O., Winston, B., Lindsay, R. 2006. Burn  
423 wound infections. *Clinical Microbiology Reviews*; 19, 403-434.
- 424 [11] Howes, E.L. 1947. Topical use of streptomycin in wounds. *The American*  
425 *Journal of Medicine*; 2, 449-456.
- 426 [12] Kilner, T.P., Peet, E., Calnan, J.S., Kernahan, D.A. 1951. Streptomycin as a  
427 local application to control wound infection prior to skin grafting: preliminary  
428 report. *British Journal of Plastic Surgery*; 4, 74-78.
- 429 [13] Dutta, N.K., Mazumdar, K., Dastidar, S.G., Park, J.-H. 2007. Activity of  
430 diclofenac used alone and in combination with streptomycin against  
431 *Mycobacterium tuberculosis* in mice. *International Journal of Antimicrobial*  
432 *Agents*; 30, 336-340.
- 433 [14] Dutta, N.K., Dastidar, S.G., Kumar, A., Mazumdar, K., Ray, R., Chakrabarty,  
434 A.N. 2004. Antimycobacterial activity of the antiinflammatory agent diclofenac  
435 sodium, and its synergism with streptomycin. *Brazilian Journal of Microbiology*;  
436 35, 316-323.
- 437 [15] Mazumdar, K., Dastidar, S., Park, J., Dutta, N. 2009. The anti-inflammatory  
438 non-antibiotic helper compound diclofenac: an antibacterial drug target. *European*  
439 *Journal of Clinical Microbiology & Infectious Diseases*; 28, 881-891.
- 440 [16] Boateng, J.S., Pawar, H.V., Tetteh, J. 2013. Polyox and carrageenan based  
441 composite film dressing containing anti-microbial and anti-inflammatory drugs for  
442 effective wound healing. *International Journal of Pharmaceutics*; 441, 181-191  
443 (2013).
- 444 [17] Pawar, H.V., Tetteh, J., Boateng, J.S. 2013. Preparation, optimisation and  
445 characterisation of novel wound healing film dressings loaded with streptomycin  
446 and diclofenac. *Colloids and Surfaces B: Biointerfaces*; 102, 102-110.

- 447 [18] Labovitiadi, O., Lamb, A.J., Matthews, K.H. 2012. In vitro efficacy of  
448 antimicrobial wafers against methicillin-resistant *Staphylococcus aureus*.  
449 *Therapeutic Delivery*; 3, 443-455.
- 450 [19] Andrews JM. 2001. Determination of minimum inhibitory concentrations.  
451 *Journal of Antimicrobial Chemotherapy*; 48 (suppl 1): 5-16.
- 452 [20] Zhanel, G.G., Shroeder, C., Vercaigne, L., Gin, A.S., Embil, J., Hoban, D.J.  
453 2001. A critical review of oxazolidinones: an alternative or replacement for  
454 glycopeptides and streptogramins? *Canadian Journal of Infectious Diseases*; 12,  
455 379-390.
- 456 [21] Craig, W. 1998. Choosing an antibiotic on the basis of pharmacodynamics.  
457 *ENT Journal*; 77 (Suppl 6):7-11.
- 458 [22] Nightingale, C. Pharmacokinetics and pharmacodynamics of newer  
459 macrolides. *Pediatric Infectious Diseases*, 1997; 16:438-43.
- 460 [23] Zhanel, G.G., Craig, W.A. 1994. Pharmacokinetic contributions to post-  
461 antibiotic effects: focus on aminoglycosides. *Clinical Pharmacokinetics*; 27:377-  
462 92.
- 463 [24] Zhanel, G.G., Waltky, A., Vercaigne, L., Karlowsky, J.A. Embil, J., Gin,  
464 A.S., Hoban, D.J. 1999. The new fluoroquinolones: a critical review. *Canadian*  
465 *Journal of Infectious Diseases*; 10:207-38.
- 466 [25] Thomas, J.K., Forrest, A., Bhaunani, S.M., Hyatt, J.M., Cheng, A., Ballow, C.H.,  
467 Schentag, J.J. 1998. Pharmacodynamic evaluation of factors associated with the  
468 development of bacterial resistance in acutely ill patients during therapy.  
469 *Antimicrobial Agents in Chemotherapy*; 42, 521-527.

- 470 [26] Bajpai, S.K., Sharma, S. 2004. Investigation of swelling/degradation  
471 behaviour of alginate beads crosslinked with Ca<sup>2+</sup> and Ba<sup>2+</sup> ions. *Reactive and*  
472 *Functional Polymers*; 59, 129-140.
- 473 [27] Boateng, J.S., Matthew, K.H., Auffret, A.D., Humphrey, M.J., Stevens,  
474 H.N., Eccleston, G. M. 2009. In vitro drug release studies of polymeric freeze-dried  
475 wafers and solvent-cast films using paracetamol as a model soluble drug.  
476 *International Journal of Pharmaceutics*; 378, 66-72.
- 477 [28] Jung, W.K., Koo, H.C., Kim, K.W., Shin, S., Kim, S.H., Park, Y.H. 2008.  
478 Antibacterial activity and mechanism of action of the silver ion in *Staphylococcus*  
479 *aureus* and *Escherichia coli*. *Applied and Environmental Microbiology*; 74, 2171-  
480 2178.
- 481 [29] Vermeulen, H., Van Hattem, J.M., Storm-Versloot, M.N., Ubbink, D.T.  
482 2007. Topical silver for treating infected wounds. *Cochrane Database of*  
483 *Systematic Reviews*; DOI: 10.1002/14651858.CD005486.pub2.
- 484 [30] Dowsett, C. 2004. The use of silver-based dressings in wound care. *Nursing*  
485 *Standard*; 19, 56-60.
- 486 [31] Modak, S.M., Fox, C.L. 1981. Sulfadiazine silver-resistant pseudomonas in  
487 burns - new topical agents. *Archives of Surgery*; 116, 854-857.
- 488 [32] Ovington, L.G. 2004. The truth about silver. *Ostomy/wound management*;  
489 50, 1S-10S.
- 490 [33] Aramwit, P., Muangman, P., Namviriyachote, N., Srichana T. 2010. In vitro  
491 evaluation of the antimicrobial effectiveness and moisture binding properties of  
492 wound dressings. *International Journal of Molecular Science*. 11:2864-2874.
- 493 [34] Mertz, P.M. 2003. Cutaneous biofilms: Friend or foe? *Wounds-a*  
494 *Compendium of Clinical Research and Practice*; 15, 129-132.

- 495 [35] Innes, M.E., Umraw, N., Fish, J.S., Gomez, M., Cartotto, R.C. 2001. The  
496 use of silver coated dressings on donor site wounds: a prospective, controlled  
497 matched pair study. *Burns*; 27, 621-627.
- 498 [36] Percival, S.L., Bowler, P.G., Russell, D. 2005. Bacterial resistance to silver  
499 in wound care. *Journal of Hospital Infection*; 60, 1-7.
- 500 [37] Li, X.Z., Nikaido, H., Williams, K.E. 1997. Silver-resistant mutants of  
501 *Escherichia coli* display active efflux of Ag<sup>+</sup> and are deficient in porins. *Journal of*  
502 *Bacteriology*; 179, 6127-6132.
- 503 [38] Braff, M.H., Jones, A.L., Skerrett, S.J., Rubens, C.E. 2007. *Staphylococcus*  
504 *aureus* exploits cathelicidin antimicrobial peptides produced during early  
505 pneumonia to promote staphylokinase-dependent fibrinolysis. *Journal of Infectious*  
506 *Diseases*; 195, 1365-1372.
- 507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520

521 **FIGURE LEGENDS**

522 **Figure 1:** ZOI of control STP (N) and control DLF (O) for *S. aureus*, *P. aeruginosa* and *E.*  
523 *coli*. The inset of control DLF shows the absence of bacteria around the applied area of the  
524 disk (mean  $\pm$  SD,  $n = 3$ ).

525 **Figure 2:** Extracted data comparing the measured ZOI data (mm) of *S. aureus*, *P.*  
526 *aeruginosa* and *E. coli* for the various formulations and marketed dressings tested.(A)  
527 POL-CAR (DL and DL-20%GLY) films and STP and DLF (mean  $\pm$  SD,  $n = 3$ ). (B) POL-  
528 SA-DL and POL-SA-DL-9%GLY films and control STP and DLF (mean  $\pm$  SD,  $n = 3$ ). (C)  
529 POL-CAR-DL-An and POL-SA-DL-An wafers and control STP and DLF (mean  $\pm$  SD,  $n =$   
530 3). (D) The marketed dressings (Aquacel<sup>®</sup> Ag, Melgisorb<sup>®</sup> Ag, Allevyn<sup>®</sup> Ag (mean  $\pm$  SD,  
531  $n=3$ ).

532 **Figure 3:** The digital images of ZOI of (A) POL-CAR-BLK, (B) POL-CAR-DL, (C) POL-  
533 CAR-DL-20%GLY films observed for *S. aureus*, *P. aeruginosa* and *E. coli*(mean  $\pm$  SD,  
534  $n=3$ ).

535 **Figure 4:** The digital images of ZOI of (D) POL-SA-BLK, (B) POL-SA-DL, (C) POL-SA-  
536 DL-9%GLY observed for *S. aureus*, *P. aeruginosa* and *E. coli*(mean  $\pm$  SD,  $n = 3$ ).

537 **Figure 5:** Digital images of ZOI of (G) POL-CAR-BLK-An, (H) POL-CAR-DL-An, (I)  
538 POL-SA-BLK-An, (J) POL-SA-DL-An, observed for *S. aureus*, *P. aeruginosa* and *E. coli*  
539 (mean  $\pm$  SD,  $n = 3$ ).

540 **Figure 6:** Digital images of ZOI observed for *S. aureus*, *P. aeruginosa* and *E. coli* by (K)  
541 Aquacel<sup>®</sup> Ag; (L) Melgisorb<sup>®</sup> Ag; and (M) Allevyn<sup>®</sup> Ag. Inset shows the absence of bacteria  
542 in the immediate applied area of the dressing (mean  $\pm$  SD,  $n = 3$ ).

543

544  
545  
546  
547  
548  
549  
550

**Table 1:** Formulations used to evaluate antimicrobial efficacy against *S. aureus*, *P. aeruginosa* and *E. coli*.

| <b>Formulation</b> | <b>CODE</b> |
|--------------------|-------------|
| POL-CAR-BLK        | A           |
| POL-CAR-DL         | B           |
| POL-CAR-DL-20%GLY  | C           |
| POL-SA-BLK         | D           |
| POL-SA-DL          | E           |
| POL-SA-DL-9%GLY    | F           |
| POL-CAR-BLK-An     | G           |
| POL-CAR-DL-An      | H           |
| POL-SA-BLK-An      | I           |
| POL-SA-DL-An       | J           |
| Aquacel® Ag        | K           |
| Melgisorb® Ag      | L           |
| Allevyn® Ag        | M           |
| STP                | N           |
| DLF                | O           |

551  
552



553

554 Table 2: Quantities of the polymers, drugs and GLY (varying amounts based on total solid  
 555 weight) within composite polymer gels used for formulation of POL-CAR and POL-SA  
 556 (BLK and DL) films.

| <b>Formulation</b> | <b>POL<br/>(gm)</b> | <b>CAR<br/>(gm)</b> | <b>SA<br/>(gm)</b> | <b>GLY<br/>(gm)</b> | <b>DLF<br/>(gm)</b> | <b>STP<br/>(gm)</b> | <b>Total<br/>weight<br/>(gm)</b> | <b>% GLY<br/>Content</b> |
|--------------------|---------------------|---------------------|--------------------|---------------------|---------------------|---------------------|----------------------------------|--------------------------|
| POL-CAR-BLK        | 0.75                | 0.25                | -                  | 0.00                | -                   | -                   | 1.00                             | 0.00                     |
| POL-CAR-BLK        | 0.75                | 0.25                | -                  | 0.10                | -                   | -                   | 1.10                             | 9.09                     |
| POL-CAR-BLK        | 0.75                | 0.25                | -                  | 0.25                | -                   | -                   | 1.25                             | 20.00                    |
| POL-CAR-BLK        | 0.75                | 0.25                | -                  | 0.50                | -                   | -                   | 1.50                             | 33.33                    |
| POL-CAR-BLK        | 0.75                | 0.25                | -                  | 0.75                | -                   | -                   | 1.75                             | 42.86                    |
| POL-CAR-BLK        | 0.75                | 0.25                | -                  | 1.00                | -                   | -                   | 2.00                             | 50.00                    |
| POL-CAR-DL         | 0.75                | 0.25                | -                  | 0.00                | 0.10                | 0.30                | 1.40                             | 0.00                     |
| POL-CAR-DL         | 0.75                | 0.25                | -                  | 0.10                | 0.10                | 0.30                | 1.50                             | 6.67                     |
| POL-CAR-DL         | 0.75                | 0.25                | -                  | 0.25                | 0.10                | 0.30                | 1.65                             | 15.15                    |
| POL-CAR-DL         | 0.75                | 0.25                | -                  | 0.50                | 0.10                | 0.30                | 1.90                             | 26.32                    |
| POL-CAR-DL         | 0.75                | 0.25                | -                  | 0.75                | 0.10                | 0.30                | 2.15                             | 34.88                    |
| POL-CAR-DL         | 0.75                | 0.25                | -                  | 1.00                | 0.10                | 0.30                | 2.40                             | 41.67                    |
| POL-SA-BLK         | 0.50                | -                   | 0.50               | 0.00                | -                   | -                   | 1.00                             | 0.00                     |
| POL-SA-BLK         | 0.50                | -                   | 0.50               | 0.10                | -                   | -                   | 1.10                             | 9.09                     |
| POL-SA-BLK         | 0.50                | -                   | 0.50               | 0.25                | -                   | -                   | 1.25                             | 20.00                    |
| POL-SA-BLK         | 0.50                | -                   | 0.50               | 0.50                | -                   | -                   | 1.50                             | 33.33                    |
| POL-SA-DL          | 0.50                | -                   | 0.50               | 0.00                | 0.05                | 0.15                | 1.20                             | 0.00                     |
| POL-SA-DL          | 0.50                | -                   | 0.50               | 0.10                | 0.05                | 0.15                | 1.30                             | 7.69                     |
| POL-SA-DL          | 0.50                | -                   | 0.50               | 0.25                | 0.05                | 0.15                | 1.45                             | 17.24                    |
| POL-SA-DL          | 0.50                | -                   | 0.50               | 0.50                | 0.05                | 0.15                | 1.70                             | 34.48                    |

557

558

559

560

561

562

563

564

565

566

567 Table 3: Composition of polymers and drugs (varying quantity) present in composite  
568 polymer gels used to produce composite freeze dried POL-CAR and POL-SA (BLK and  
569 DL) wafers.

| <b>Pure material</b> | <b>POL-CAR-BLK</b>    | <b>POL-CAR-DL</b> | <b>POL-SA-BLK</b> | <b>POL-SA-DL</b> |
|----------------------|-----------------------|-------------------|-------------------|------------------|
|                      | <b>(weight in gm)</b> |                   |                   |                  |
| POL                  | 0.75                  | 0.75              | 0.50              | 0.50             |
| CAR                  | 0.25                  | 0.25              | -                 | -                |
| SA                   | -                     | -                 | 0.50              | 0.50             |
| STP                  | -                     | 0.30              | -                 | 0.25             |
| DLF                  | -                     | 0.25              | -                 | 0.10             |
| <b>Total weight</b>  | <b>1.00</b>           | <b>1.55</b>       | <b>1.00</b>       | <b>1.35</b>      |

570

571

572

573

574

575

576

577 **Table 4:** Stock solutions of STP and DLF used to evaluate MIC of *S. aureus*, *E. coli* and *P.*  
578 *aeruginosa* (mean  $\pm$  SD,  $n = 3$ ).

579

|     | Stock solution 1                                      | Stock solution 2   | Stock solution 3  |
|-----|---|--|---|
| STP | 10000 mg/L (254 mg of STP + 20 ml of distilled water) | 1000 mg/L (1 ml of stock solution 1 + 9 ml of distilled water) | 100 mg/L (1 ml of stock solution 2 + 9 ml of distilled water) |
| DLF | 10000 mg/L (200 mg of DLF + 20 ml of distilled water) | 1000 mg/L (1 ml of stock solution 1 + 9 ml of distilled water) | 100 mg/L (1 ml of stock solution 2 + 9 ml of distilled water) |


580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
  
591  
592  
593  
594

**Table 5:** Description of the silver containing dressings used for antimicrobial study (Hamberg *et al.*, 2012) (mean  $\pm$  SD,  $n = 3$ ).

| Product                   | Formulation details  | Silver content (mg/cm <sup>2</sup> ) |
|---------------------------|--|--------------------------------------|
| Aquacel <sup>®</sup> Ag   | Sodium carboxymethylcellulose with ionic silver                            | 0.08-0.09                            |
| Melgisorb <sup>®</sup> Ag | Alginate dressing with silver sodium hydrogen zirconium phosphate          | 0.08                                 |
| Allevyn <sup>®</sup> Ag   | Polyurethane foam dressing with soft gel adhesive and silver sulphadiazine | 0.90                                 |

595

596




597

598


599 **Figure 1**

600



601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615

**Figure 2**




616

617

618 **Figure 3**

619



620

621

622 **Figure 4**

623

624

625

626

627

628

629

630

631

632


633

634

635



636




637

638

639 **Figure 5**

640

641



642

643

644

645

646

647

648

649

650

**Figure 6**