

4
5 **Frailty, Swallowing and Dysphagia**

6 **David G. Smithard^{1,2} · M. Shazra³ · D. Hansjee¹ · I. Swaine²**

7
8
9 © Springer Science+Business Media, LLC, part of Springer Nature 2018

10 **Abstract**

11 **Purpose of Review** This paper is a brief overview of the relationship between frailty, swallowing and dysphagia. Its goal is to
 12 explore the interplay between age and sarcopenia in the development of dysphagia, which is known to be linked to aspiration
 13 pneumonia. It is postulated that there is growing justification for routine screening for dysphagia in older frail people, to enable
 14 rehabilitation of swallowing through exercise and nutritional intervention, after a hospital stay.

15 **Recent Findings** The global population is ageing, with a particular increase in the very old and frail. Frail people have a limited
 16 functional and physiological reserve and often have sarcopenia. Any subsequent insult (trauma, illness, medication change)
 17 frequently results in decompensation and the need for a hospital stay. Often, in these patients, there are changes in the biome-
 18 chanics of swallowing that can cause impairment and dysphagia. But, many patients adapt the way they eat with subtle
 19 compensatory techniques, to bypass this difficulty. It is possible that many more people, than is currently evident, have undiag-
 20 nosed dysphagia. Pneumonia and respiratory disease are common reasons for hospital admission in the frail elderly population.
 21 Dysphagia with aspiration is an important aetiological factor in pneumonia, which is a serious health concern with increasing age.
 22 Dysphagia may simply be a consequence of physiological decompensation, related to age, frailty and sarcopenia. Dysphagia is
 23 not systematically screened for and may not be identified in many older frail people who have adapted their swallowing, to
 24 accommodate their dysphagia. This may be a significant factor in pneumonia-related hospital admissions. Swallow rehabilitation,
 25 after such admission to hospital, is also rarely offered in the acute medical setting. This needs to change to reduce recurrent
 26 admission, morbidity and mortality.

27 **Summary** The population is ageing. Sarcopenia, frailty and dysphagia are common with increasing age. Pneumonia is a common
 28 admission to hospital and often, aspiration secondary to dysphagia is a common cause. Proactive identification and intervention
 29 has the potential to reduce morbidity, hospital admission, length of hospital stay and mortality.

30 **Keywords** Frailty · Dysphagia · Exercise · Nutrition · Inflammation

31
32 **Introduction**

33 The global population is expanding in an asymmetric way,
 34 with an increase in older and in particular very old people.
 35 This is driven, in part by the improvements in public health
 36 and acute medical care. As the population ages, so the number

of people with long-term conditions increases. This ultimately 37
 results in more people becoming frail, such that 50% of those 38
 over the age of 80 years are categorised as frail [1]. Frailty is 39
 described as a general decline in physiological function asso- 40
 ciated with loss of muscle bulk, weakness, fatigue and slow- 41
 ness. Sarcopenia frequently accompanies frailty and is also 42
 associated with muscle weakness, loss of skeletal muscle mass 43
 and poor quality of muscle fibres. 44

Frailty is a complex syndrome associated with a progres- 45
 sive decline in physical, mental and social functions. There is 46
 increased vulnerability to deterioration and reduced potential 47
 for recovery [1, 2] associated with sarcopenia, undernutrition, 48
 slowness and disability. Frail older people have little or no 49
 physiological reserve, such that whenever they are exposed 50
 to an insult (medication change, constipation, trauma and 51
 acute illness) physiological and functional decompensation 52
 results. The insult may be minor, but due to the lack of reserve, 53

This article is part of the Topical Collection on *Swallowing Disorders*

✉ David G. Smithard
 david.smithard@nhs.net

1 Queen Elizabeth Hospital, Woolwich, London, UK
 2 University of Greenwich, London, UK
 3 Princess Royal University Hospital, Farnborough, Kent, Orpington, UK

54 it will result in reduced mobility, confusion/delirium and fre- 104
 55 quently, dysphagia. Frail older people also have multiple co- 105
 56 morbidities, increased mortality [3] and have a limited phys- 106
 57 iological reserve. Any illness will result in decompensation [4, 107
 58 5], which may include the development of difficulties in 108
 59 swallowing (dysphagia).

60 The International Classification of Functioning, Disability 109
 61 and Health [6] classifies normal swallowing as ‘functions of 110
 62 clearing the food and drink through the oral cavity, pharynx 111
 63 and oesophagus into the stomach at an appropriate rate and 112
 64 speed’. As the bolus (food or liquid) passes through the mouth 113
 65 and pharynx, it is subject to multiple pressures, which are 114
 66 dependent on the function of the tongue and upper- 115
 67 oesophageal sphincter (UES), pharynx and influenced by the 116
 68 inherent properties of the bolus.

69 **Frailty-Related Dysphagia**

70 Frail older people have muscle loss and sarcopenia; the loss of 117
 71 appendicular skeletal muscle is easy to recognise. The supra- 118
 72 hyoid muscles, which are involved in swallowing (assisting in 119
 73 tongue movement, laryngeal elevation and anterior movement 120
 74 and UES opening [7]), are also skeletal muscles and will be 121
 75 affected by sarcopenia. This is often forgotten, or medical staff 122
 76 is not aware of the fact. The consequences of this include re- 123
 77 duced elevation and anterior movement of the larynx and re- 124
 78 duced opening aperture of the UES, which results in a poten- 125
 79 tially less safe swallow than in younger adults [8]. A smaller 126
 80 UES aperture and reduced pharyngeal stripping wave pressures 127
 81 result in food residue remaining in the pharynx after the swal- 128
 82 low, increasing the risk of post-swallow aspiration [4, 7, 9, 10]. 129

83 Frail older adults have complex medical needs. Frequently 130
 84 people have multiple, co-existing, long-term medical condi- 131
 85 tions that will impinge on the eating, drinking and swallowing. 132
 86 For example, rheumatoid arthritis can affect the arytenoid 133
 87 joints, thereby affecting closing of the vocal cords [11]. 134
 88 Swallowing requires a short period of breath holding; in the 135
 89 context of cardio-respiratory disease can lead to hypoxia and 136
 90 secondary dysphagia; head and neck cancer; stroke, vascular 137
 91 or Alzheimer’s dementia can all lead to changes in the oral or 138
 92 pharyngeal phases of the swallow. A dry mouth secondary to 139
 93 medication, mouth breathing, radiotherapy or auto immune 140
 94 disease can all affect the consistency of food eaten. 141

95 The aetiology of sarcopenia and frailty is complex often 142
 96 involving neurohumoral pathways (suggesting inflammatory 143
 97 pathway involvement), poor diet and lack of exercise. 144
 98 Therefore, it should be feasible to break the downward 145
 99 spiral/cycle which will ultimately result in death, through life- 146
 100 style change. This could help to avoid the need for a hospital 147
 101 stay every time a frail person becomes unwell. When this 148
 102 occurs, the functional ability of the older frail person declines 149
 103 and recovery may take many months. Often, the end result, 150

104 physically and cognitively, may be a person who is more 105
 106 dependent than previously. 107

108 To reduce and/or ameliorate the presence of dysphagia and 109
 110 its complications in this older cohort, strategies to combat 111
 112 frailty need to be developed. 113
 114
 115
 116

117 **Exercise**

118 There is increasing evidence to suggest that exercise can coun- 119
 120 teract some of the effects of frailty on physical function. Many 121
 122 expert groups are recommending exercise for frail older peo- 123
 124 ple. Indeed, a recent review suggested that exercise consistent- 125
 126 ly combats the deleterious effects of sarcopenia. Most reviews 127
 128 have also suggested that resistance exercise, for muscle- 129
 130 strengthening, is more efficacious in this regard [12]. 131

132 Muscle mass and strength can be maintained and increased 133
 134 by aerobic and resistance exercise (including isometric exer- 135
 136 cise). For example, taking up dancing can improve balance and 137
 138 coordination, singing can help respiratory muscle strength and 139
 140 walking can improve aerobic capacity and leg muscle strength. 141
 142 Exercise, commenced in or before middle age, has been shown 143
 144 to protect against circulatory disease (stroke and heart). In the 145
 146 case of appendicular sarcopenia, other types of exercise may 147
 148 prevent or improve muscle bulk and strength [1, 13–15]. 149

150 It is therefore plausible that a ‘general’ exercise programme 151
 152 could help to counteract frailty-related dysphagia. Improvement 153
 154 in cardiorespiratory fitness and general muscle strength could 154
 155 have an indirect effect on the aerobic function and strength of 155
 156 the swallowing musculature. However, more-targeted, specific 156
 157 exercise programmes have been developed to improve 157
 158 swallowing. These exercise programmes target the swallowing 158
 159 muscles (e.g. hyoid muscles) usually by requiring people to 159
 160 perform chin tuck movements against some type of resistance. 160
 161 Because this type of exercise usually involves a ‘static phase’ 161
 162 (where the movement is held stationary against the resistance, 162
 163 for example, for 30 s, or 1–2 min), these exercises are some- 163
 164 times described as ‘isometric resistance’. 164

165 Indeed, for dysphagia, Shaker and colleagues developed a 165
 166 system of exercises to train the hyoid group of muscles and neck 166
 167 muscles [16]. Shaker exercise resulted in an increase in 167
 168 thyrohyoid shortening after 6 weeks compared to tongue exer- 168
 169 cises and swallowing manoeuvres [17]; increases in contraction 169
 170 pressures in the pharynx, increased opening of the UES, [18, 19] 170
 171 occurs. However, the studies are small, and more data is required 171
 172 to determine how chin tuck affects the physiology of swallowing. 172

173 Therefore, it may be entirely possible to significantly affect 173
 174 dysphagia in the frail older person through general and targeted 174
 175 exercise programmes. The mechanism whereby this occurs 175
 176 needs to be further elucidated, but would probably involve im- 176
 177 proved oxygen supply, muscle strength, endurance, coordina- 177
 178 tion and neurohumoral control of movement. The most appro- 178
 179 priate way to achieve these effects, in relation to frail older 179
 180

154 people with dysphagia, is not yet known. This is largely because
 155 so few studies have explored this idea in this group of people.

156 **Nutrition**

157 Malnutrition is associated with increased infection risk and
 158 poor wound healing. Malnutrition and dehydration will result
 159 in muscle weakness and hypotension and an increased risk of
 160 falls. Falls in frail older people, place them at risk of subdural
 161 haematomas and fractured neck of femurs, both of which carry
 162 a significant morbidity and mortality.

163 Improvement in nutrition and calorie intake will help in the
 164 anabolic effect provided by exercise. Older adults are ineffi-
 165 cient in the utilisation of amino-acids from protein and are also
 166 inefficient in energy usage and hence older people require a
 167 balanced diet to maintain weight, muscle mass, strength, and
 168 to improve immunity.

169 **Inflammation**

170 More recently there has been interest shown in the interruption
 171 of the inflammatory process (raised CRP, interleukins and
 172 cytokines) present in sarcopenia and frailty, to arrest the pro-
 173 cess. Some work has shown a benefit in the use of anti-
 174 inflammatory agents (Ibuprofen). However, there is limited
 175 evidence of clinical benefit [20].

176 **Dysphagia**

177 Various studies have noted a frequency of up to 30% of older
 178 people living at home may have dysphagia [21] with up to
 179 28% of older people noted to be aspirating on instrumentation
 180 [22, 23]. The frequency of swallowing problems in older peo-
 181 ple is uncertain as many older people do not report problems
 182 [4] or have accepted them as a fact of life [24] and have often
 183 subtly and unknowingly compensated for motor changes [4].

184 Many older frail people admitted to hospital (55%) will
 185 have difficulties with swallowing (dysphagia) [25]. Up to
 186 90% of those admitted with a diagnosis of pneumonia, may
 187 have inhaled saliva or food into the lungs (aspiration).
 188 Periodontal disease is not uncommon, with rotten teeth and
 189 gum infection, which will increase the risk of pneumonia sec-
 190 ondary to the inhalation of infected saliva. Recurrent infection
 191 will result in a decreased lung function and also a worsening
 192 of functional state following each infection.

193 Swallowing problems/dysphagia, when present in older
 194 people, is poorly managed in many institutionalised settings
 195 including the acute hospital. In many European countries, the
 196 presence of dysphagia (or difficulties with eating and drink-
 197 ing) in frail people is not routinely sought. There is no

requirement to systematically screen frail patients for the pres- 198
 ence of dysphagia as there is in stroke patients [26, 27]. This is 199
 a missed opportunity, by policy makers, to potentially im- 200
 prove outcome and reduce hospital stay and re-admissions. 201

Routine Screening for Dysphagia in Frail Older People 202
 203

If frail patients are to be screened for dysphagia, on admission 204
 to hospital, which swallow screen should be used? There are 205
 many swallowing screens available to clinicians (Medical, 206
 SLP, AHP, Nursing), many which are validated, but many 207
 staff continue to use locally developed, non-validated tools 208
 [28]. These tools have been validated in the stroke population 209
 and recently, evidence has been published supporting the use 210
 of TOR-BSST in care homes [29]. 211

The swallow screens that are available are generally very 212
 similar. They all have the same aim, of a clinical/anatomical 213
 assessment, a trial of small volumes of water followed by a 214
 larger volume. Some are simple, with others being more com- 215
 plex. Recent work that we have conducted has shown that the 216
 most characteristic identifiers are coughing and choking when 217
 swallowing, taking longer to drink, a change in diet and a 218
 change in voice quality. 219

Proactive Intervention 220

Implementing a policy of screening will not improve patient 221
 outcomes unless there is a positive proactive intervention. 222
 Standard hospital rehabilitation, for dysphagia, includes pos- 223
 tural manoeuvres (e.g. Chin-Tuck). It has been accepted that 224
 the Chin Tuck manoeuvre pulls the larynx up and forwards 225
 and at the same time opens the UES [17, 30]. Welch et al. [30] 226
 also claimed that there was posterior movement of the tongue 227
 base towards the posterior pharyngeal wall. Momasaki et al. 228
 [31] using a large Japanese database demonstrated that those 229
 patients with dysphagia, who were offered appropriate reha- 230
 bilitation, were more likely to have a total oral intake com- 231
 pared to those not offered oral-pharyngeal rehabilitation (OR 232
 1.2 $P < 0.001$). Pogus-Pulia et al. [32] demonstrated, using 233
 isometric progressive resistance oropharyngeal therapy [33], 234
 improved FOIS (Functional Oral Intake Score- a measure of 235
 food intake [34] effect estimate = 0.4, $p < 0.02$), reduced inci- 236Q2
 dence of pneumonia and reduced number of hospital admis- 237
 sions. Their cohort was mixed in aetiology of dysphagia. 238

Mouth Care 239

Dentition and mouth care are important factors to consider. 240
 Lack of teeth will make chewing more difficult, resulting in 241

242 modifications to the food choices eaten. This may mean
 243 blending what is usually eaten, or switching to softer and
 244 possibly nutritionally poorer foods. For those who are eden-
 245 tulous, dentures are often provided (where available), but in
 246 many cases are not worn due to discomfort, or are not kept
 247 clean. Unhygienic dentures could result in candidiasis and in
 248 dysphagia due to pain.

249 **End of Life**

250 Frailty carries a significant mortality, with those with extreme
 251 frailty [35] having a short life span. The management of dys-
 252 phagia in this group will move fairly rapidly from active man-
 253 agement to a palliative care approach. There will be a move
 254 from ensuring adequate calories, to that of offering food/liquid
 255 for comfort as and when the person is able and in volumes that
 256 are safe. There is little consensus on terminology with respect
 257 to this approach and terms include ‘finger feeding’ and ‘risk
 258 feeding’. How this is approached will vary depending on so-
 259 cietal, cultural, religious sensibilities and expectations.

260 **Conclusions**

261 Worldwide, by 2050, 34% of the world’s 9.7 billion popula-
 262 tion there will be > 60 years, with 125 million > 80 years [36].
 263 Fifty percent of older adults > 80 years older adults will be
 264 frail and prone to deteriorating health and dependence. It is
 265 possible to slow down the march to frailty and dependence,
 266 particularly in the pre-frail phase, by implementing a pro-
 267 gramme of exercise and good nutrition. The presence of dys-
 268 phagia is likely to be high in this population; routine screening
 269 for problems, eating, drinking and swallowing should be un-
 270 dertaken in primary care and when an older frail person is
 271 admitted to hospital.

272 **Compliance with Ethical Standards**

273 **Conflict of Interest** The authors declare that they have no competing
 274 interests.

275 **Human and Animal Rights and Informed Consent** This article does not
 276 contain any studies with human or animal subjects performed by any of
 277 the authors.

278 **References**

279 1. Phu S, Boersma D, Duque G. Densitometry: assessment and man-
 280 agement of musculoskeletal health. *J Clin.* 2015;18:1–5.
 281 2. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C,
 282 Gottdiener J, et al. Frailty in older adults: evidence for a phenotype.
 283 *J Gerontol.* 2001;56:M146–57.

3. Stanaway FF, Gnjjidic D, Blyth FM, Le Couteur DG, Naganathan V, 284
 Waite L, et al. How fast does the grim reaper walk? Receiver oper- 285
 ating characteristics curve analysis in healthy men aged 70 and over. 286
BMJ. 2011;343:d7679. 287
 4. Omari TI, Kritas S, Cock C, Besanko L, Burgstad C, Thompson A, 288
 et al. Swallowing dysfunction in healthy older people using pha- 289
 ryngeal pressure-flow analysis. *Neurogastroenterol Motil.* 2014;1: 290
 59–68. 291
 5. Nicosia MA, Hind JA, Roecker EB, Carnes M, Doyle J, Dengel 292
 GA, et al. Age effects on the temporal evolution of isometric and 293
 swallowing pressure. *J Gerontol: Medi Sci.* 2000;55A:M634–40. 294
 6. The International Classification of Functioning, Disability and 295
 Health (ICF). 2001 WHO Geneva [http://apps.who.int/iris/](http://apps.who.int/iris/bitstream/10665/42407/7/9241545429_tha%2Beng.pdf) 296
[bitstream/10665/42407/7/9241545429_tha%2Beng.pdf](http://apps.who.int/iris/bitstream/10665/42407/7/9241545429_tha%2Beng.pdf) 297
 7. Curtis JA, Laus J, Yung KC, Courey MS. Static endoscopic evalu- 298
 ation of swallowing: transoral endoscopy during clinical swallow 299
 evaluations. *Laryngoscope.* 2016;126:2291–4. 300
 8. Dejaeger M, Liesenborghs C and Dejaeger E Presbyphagia. In 301
Seminars in Dysphagia. InTech; 2015. 302
 9. Curtis J, Langenstein J, Scheider S. Superior and anterior displace- 303
 ment during swallowing in non-dysphagic individuals. *Dysphagia.* 304
 2018; <https://doi.org/10.1007/s00455-018-9878-7.0123456789>. 305
 10. Sze WP, Yoon WL, Escoffier N. Liow evaluating the training ef- 306
 fects of two swallowing rehabilitation therapies using surface elec- 307
 tro-myography—chin tuck against resistance (CTAR) exercise and 308
 the shaker exercise. *Dysphagia.* 2016;31:195–205. 309
 11. Gleason JB, Hadeh A. Vocal hoarseness in rheumatoid arthritis: 310
 early recognition is critical. *J Clin Diagn Res.* 2017;11(4):OJ03. 311
 12. Theou O, Stathokostas L, Roland KP, Jakobi JM, Patterson C, 312
 Vandervoort AA, et al. The effectiveness of exercise interventions 313
 for the management of frailty: a systematic review. *J Aging Res.* 314
 2011, Article ID 569194, 19 pages;2011:1. 315
 13. Otsuka R, Matsui C, Tange C, Nishita Tomida M, Fujiko A, 316
 Shimokata H, et al. What is the best adjustment of appendicular 317
 mass for predicting mortality or disability among Japanese commu- 318
 nity dwellers? *BMC Geriatr.* 2018;18:8. [https://doi.org/10.1186/](https://doi.org/10.1186/s12877-017-0699-6) 319
[s12877-017-0699-6](https://doi.org/10.1186/s12877-017-0699-6). 320
 14. Komi PV, Viitasalo JT, Rauramaa R, et al. *Europ. J Appl Physiol.* 321
 1978;40:45. 322
 15. Anwer S, Alghadir A. Effect of isometric quadriceps exercise on 323
 muscle strength, pain, and function in patients with knee osteoar- 324
 thritis: a randomized controlled study. *J Phys Ther Sci.* 2014;26(5): 325
 745–8. 326
 16. Shaker R, Eastreling C, Kern M, Nitschke T, Massey B, Daniels 327
 SK, et al. Rehabilitaion of swallowing by exercises in tube-fed 328
 patients with pharyngeal dysphagia secondary to abnormal UES 329
 opening. *Gastroenterology.* 2002;122:131401321. 330
 17. Mapani R, Antonik S, Massey B, Kern M, Logemann J, Pauloski B, 331
 et al. Augmentation of deglutitive thyrohyoid muscle shortening by 332
 the Saker exercise. *Dysphagia.* 2009;24:26–31. 333
 18. Balaou M, McCullough GH, Aduli F, Brown D, Stack BC, Snoddy 334
 P, et al. Manometric measure of head rotation and chin tuck in 335
 healthy participants. *Dysphagia.* 2014;29:25–32. 336
 19. McCullough GH, Kamarunas E, Mann GC, Schmidley JW, 337
 Robbins JA, Crary MA. Effects of Mendelsohn maneuver on mea- 338
 sures of swallow duration post stroke. *Top Stroke Rehabil.* 339
 2012;19:234–43. 340
 20. Li H, Manwani B, Leng SX. Frailty, inflammation, and immunity. 341
Ageing Dis. 2011;2:466–73. 342
 21. Kertscher B, Speyer R, Fong E, Georgiou AM, Smith M. 343
 Prevalence of oropharyngeal dysphagia in the Netherlands: a tele- 344
 phone survey. *Dysphagia.* 2015;30:114–20. 345
 22. Butler SG, Stuart A, Leng X, Rees C, Williamson J, Kritchevsky 346
 SB. Factors influencing aspiration dur- ing swallowing in healthy 347
 older adults. *Laryngoscope.* 2010;120:2147–52. 348

- 349 23. Almirall J, Rofes L, Serra-Prat M, Icart R, Palomera E, Arreola V, et al. Oropharyngeal dysphagia is a risk factor for community-acquired pneumonia in the elderly. *Eur Respir J*. 2013;41:923–8. 376
- 350 24. Chen P, Golub J, Hapner E, Johns MM. Prevalence of perceived 377
- 351 dysphagia and quality of life impairment in a geriatric population. 378Q3
- 352 *Dysphagia*. 2009;24:1–6. 379
- 353 25. Cabre M, Serra-Pratt S, Palomera E, Almirall J, Pallares R, Clavé. 380
- 354 Prevalence and prognostic implications of dysphagia in elderly pa- 381
- 355 tients with pneumonia. *Age Ageing*. 2010;39:39–45. 382
- 356 26. National Clinical Guideline for Stroke. Intercollegiate stroke work- 383
- 357 ing party. 5th ed. London: RCP; 2016. 384
- 358 27. Powers WJ, Rabinstein A, Ackerson T, Adeoye OM, Bambakidis 385
- 359 NC, Becker K, et al. 2018 guidelines for the early management of 386
- 360 patients with acute ischaemic stroke: a guideline for healthcare 387
- 361 providers from the American Heart Association/American Stroke 388
- 362 Association. *Stroke*. 2018;49:e46–99. <https://doi.org/10.1161/STR.000000000000158>. 389
- 363 28. Fairfield C, Smithard D. Management of dysphagia in the acute 390
- 364 phase: a consideration of international practice. *Dysphagia across 391*
- 365 ages. 4th Congress of European Society of Swallowing Disorders. 392
- 366 Congress Abstracts. *Dysphagia*. 2015;30:213–71. 393
- 367 29. Park YH, Bang HL, Han HR, Chang HK. Dysphagia screening 394
- 368 measures for use in nursing homes: a systematic review. *J Korean 395Q4*
- 369 Acad Nurs. 2015;45:1–13. 396
- 370 30. Welch MV, Logemann JA, Rademaker AW, Kahrilas PJ. Changes 397
- 371 in pharyngeal dimensions effected by chin tuck. *Arch Phys Med 398*
- 372 Rehabil. 1993;74:78–81. 399
- 373 31. Momosaki R, Yasunaga H, Matsui H, Horiguchi H, Fushimi K, 376
- 374 Abo M. Effect of dysphagia rehabilitation on oral intake in elderly 377
- 375 patients with aspiration pneumonia. *Geriatr Gerontol Int* b2014. 378Q3
- 376 32. Rogu-Pulia N, Rusche N, Hind JA, Zielinski J, Gangnon R, Safdar 379
- 377 N, et al. Effects of device-facilitated isometric progressive resis- 380
- 378 tance oropharyngeal therapy on swallowing and health related out- 381
- 379 comes in older adults with dysphagia. *JAGS*. 2016; 382
- 380 33. Robbins J, Kays SA, Gangnon RE, Hind JA, Hewitt AL, Gentry 383
- 381 LR, et al. The effects of lingual exercise in stroke patients with 384
- 382 dysphagia. *Arch Phys Med Rehabil*. 2007;88(2):150–8. 385
- 383 34. Crary MA, Carnaby Mann GD, Groher ME. Initial psychometric 386
- 384 assessment of a functional oral intake scale for dysphagia in stroke 387
- 385 patients. *Arch Phys Med Rehabil*. 2005;86:1516–20. 388
- 386 35. Mitnitski A, Graham J, Mogilner A, Rockwood K. Frailty, fitness 389
- 387 and late-life mortality in relation to chronological and biological 390
- 388 age. *BMC Geriatr*. 2002;2:1–8. 391
- 389 36. United Nations, Department of Economic and Social Affairs, 392
- 390 Population Division (2015). *World Population Ageing 2015 (ST/ 393*
- 391 ESA/SERA/390). 394
- 392 37. Artaza-Artabe I, Saez-Lopez P, Sanchez-Hernandez N, Fernandez- 395Q4
- 393 Gutierrez N, Malafarina V. The relationship between nutrition and 396
- 394 frailty: effects of protein intake, nutritional supplementation, vita- 397
- 395 min D and exercise on muscle metabolism in the elderly: a system- 398
- 396 atic review. *Maturitas*. 2016;93:89–99. 399