

ASSESSING CHEMICAL MECHANISMS UNDERLYING THE EFFECTS OF
SUNFLOWER POLLEN ON A GUT PATHOGEN IN BUMBLE BEES

*LYNN S. ADLER¹, ALISON E. FOWLER¹, ROSEMARY L. MALFI¹, PATRICK R.
ANDERSON¹, LILY M. COPPINGER¹, PHEOBE M. DENEEN^{1,2}, STEPHANIE LOPEZ¹,
REBECCA E. IRWIN³, IAIN W. FARRELL⁴ and PHILIP C. STEVENSON^{4,5}

¹ *Department of Biology, University of Massachusetts at Amherst, Amherst, Massachusetts
01003, United States*

² *current affiliation: Department of Biological Sciences, Binghamton University, Binghamton, NY
13902, United States*

³ *Department of Applied Ecology, North Carolina State University, Raleigh, North Carolina
27695, United States*

⁴ *Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AE, United Kingdom*

⁵ *University of Greenwich, Medway ME4 4TB United Kingdom*

* corresponding author: lsadler@ent.umass.edu, tel 413-545-1060, fax 413-545-3243

ORCIDs:

Adler 0000-0003-2125-5582

Fowler 0000-0002-9263-2253

Malfi 0000-0003-0144-5928

Anderson 0000-0003-1094-2483

Irwin 0000-0002-1394-4946

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

24 Stevenson 0000-0002-0736-3619

25

26 *Author contributions.* LSA, REI and PCS conceived of and designed the study. PCS and IWF
27 conducted chemical analysis of pollen and synthesized spermidines. AEF, RLM, PRA, LMC,
28 PMD, and SL carried out bioassay experiments. AEF analyzed data and prepared figures. LSA
29 wrote the manuscript with substantial contributions from PCS. All co-authors read and provided
30 feedback on the manuscript.

31

Abstract- Many pollinator species are declining due to a variety of interacting stressors including pathogens, sparking interest in understanding factors that could mitigate these outcomes. Diet can affect host-pathogen interactions by changing nutritional reserves or providing bioactive secondary chemicals. Recent work found that sunflower pollen (*Helianthus annuus*) dramatically reduced cell counts of the gut pathogen *Crithidia bombi* in bumble bee workers (*Bombus impatiens*), but the mechanism underlying this effect is unknown. Here we analyzed methanolic extracts of sunflower pollen by LC-MS and identified triscoumaroyl spermidines as the major secondary metabolite components, along with a flavonoid quercetin-3-*O*-hexoside and a quercetin-3-*O*-(6-*O*-malonyl)-hexoside. We then tested the effect of triscoumaroyl spermidine and rutin (as a proxy for quercetin glycosides) on *Crithidia* infection in *B. impatiens*, compared to buckwheat pollen (*Fagopyrum esculentum*) as a negative control and sunflower pollen as a positive control. In addition, we tested the effect of nine fatty acids from sunflower pollen individually and in combination using similar methods. Although sunflower pollen consistently reduced *Crithidia* relative to control pollen, none of the compounds we tested had significant effects. In addition, diet treatments did not affect mortality, or sucrose or pollen consumption. Thus, the mechanisms underlying the medicinal effect of sunflower are still unknown; future work could use bioactivity-guided fractionation to more efficiently target compounds of interest, and explore non-chemical mechanisms. Ultimately, identifying the mechanism underlying the effect of sunflower pollen on pathogens will open up new avenues for managing bee health.

Key Words-Bee pathogens, *Bombus impatiens*, *Crithidia bombi*, *Helianthus annuus*, pollen chemistry, pollinator health.

INTRODUCTION

Parasites and pathogens have been implicated in the declines and even disappearance of bee species, including bumble bees (Cameron et al. 2016; Cameron et al. 2011; Schmid-Hempel et al. 2014). Pollinators provide billions of dollars in crop pollination annually, and are critical for preserving plant biodiversity (reviewed in Potts et al. 2016). Species declines have been attributed to a wide range of stressors, including loss of habitat, pesticide exposure, nutritional deficits, and parasites that may all interact synergistically (Goulson et al. 2015; Vanbergen et al. 2013). Diet may play a central role in mediating how pollinators interact with their pathogens. For example, poor nutrition can reduce bee immunity and increase the detrimental effects of parasites (Alaux et al. 2010; Brown et al. 2003; Roger et al. 2017). Furthermore, quality as well as quantity of floral resources may affect bee health. Consuming a diverse pollen diet increased longevity of honey bees infected with *Nosema* (Di Pasquale et al. 2013), a pathogen that has been implicated in widespread colony losses. The consumption of *p*-coumaric acid, a component of pollen and honey, upregulated honey bee genes involved in pesticide detoxification and antimicrobial function (Mao et al. 2013). Fatty acids are necessary for bee reproduction and development (Manning 2001), including fat bodies that play a role in energy storage and release as well as immune function in insects (Arrese and Soulages 2010). Plant secondary compounds can also influence pollinator-pathogen interactions. For example, multiple nectar secondary compounds reduced intensity of infection with a gut pathogen in bumble bees at natural concentrations (e. g., Baracchi et al. 2015; Koch et al. 2019; Manson et al. 2010; Richardson et al. 2015) although this is not always the case (Biller et al. 2015; Thorburn et al. 2015). Thus,

both nutrition and secondary chemicals may play important roles in mediating pollinator health and interactions with pathogens.

Although previous research asking how floral reward chemistry affects bee pathogens has focused on nectar, pollen can contain greater diversity and orders of magnitude higher concentrations of secondary compounds than nectar (Cook et al. 2013; Palmer-Young et al. 2019b), consistent with optimal defense theory predicting greater investment in tissues more tightly linked with fitness (McKey 1979). Pollen chemistry could therefore have greater efficacy on pollinator pathogens than the less consistent effects of nectar secondary chemicals (Biller et al. 2015; Thorburn et al. 2015). Pollen nutrients including protein, lipids, amino acids and starches vary widely among plant taxa (Roulston and Cane 2000; Roulston et al. 2000; Yang et al. 2013), and could additionally influence bee susceptibility to pathogens.

We recently discovered that sunflower pollen (*Helianthus annuus*) consistently and dramatically reduced infection by *Crithidia bombi* in one species of bumble bee, *Bombus impatiens* (Giacomini et al. 2018). Pathogen cell counts were typically 30- to 50-fold lower in bees fed sunflower pollen than in bees fed buckwheat pollen (*Fagopyrum esculentum*), and at least 4-fold lower than in bees fed mixed wildflower pollen. Furthermore, this effect was consistent across 13 *H. annuus* cultivars and wild accessions, two *Helianthus* congeners, and two *Solidago* species, suggesting a potentially widespread effect in Asteraceae pollen (LoCascio et al. 2019a). These results are consistent with previous work in which the pathogen *Nosema ceranae* was most strongly reduced by sunflower honey compared to three other honeys (Gherman et al. 2014). Furthermore, mason bee (*Osmia*) species that specialize on Asteraceae pollen experienced almost no brood parasitism, whereas *Osmia* generalists had up to 50% parasitism (Spear et al. 2016). All these lines of evidence suggest some mechanism by which

1
2
3
4 101 sunflower and related pollen reduces pathogen and parasite infection.

5
6
7 102 The mechanism by which these pollens reduce pathogens could be pollen chemistry,
8
9 103 including both secondary chemicals and nutritional components. Sunflower pollen is
10
11 104 antimicrobial and inhibited the causal agent of American foulbrood, a honey bee brood pathogen,
12
13 105 *in vitro* (Fatrcova-Sramkova et al. 2016). Sunflower pollen methanolic extracts also have anti-
14
15 106 inflammatory and antitumor-promoting effects in mice (reviewed in Ukiya et al. 2003).
16
17 107 Nutritionally, sunflower pollen is low in protein (Human et al. 2007; Yang et al. 2013) and
18
19 108 lacked two essential amino acids necessary for honey bee development in one study (Nicolson
20
21 109 and Human 2013), although another study did not find the same deficiencies (McAulay and
22
23 110 Forrest 2019). However, sunflower pollen is high in saturated fatty acids such as palmitic, lauric,
24
25 111 stearic and caprylic acids (Kostic et al. 2017) and another study also found high concentrations
26
27 112 of myristic acid (Farag et al. 1978). Palmitic and stearic acids together comprise 20-35% of the
28
29 113 total lipid composition of adult honey bees, suggesting important roles in nutrition (Manning
30
31 114 2001; Robinson and Nation 1970). Lauric acid had antimicrobial properties against the honey
32
33 115 bee pathogen *Paenibacillus larvae larvae* (Feldlaufer et al. 1993), and several analogues of
34
35 116 myristic acid, a common saturated fatty acid of plant oils, have potent anti-trypanosomal activity
36
37 117 (Doering et al. 1994). Thus, fatty acids from sunflower pollen could also play important roles in
38
39 118 bee health.
40
41
42
43
44
45
46
47

48 119 The goal of this study was to assess major chemical components of sunflower pollen to
49
50 120 determine whether they represent the mechanism underlying the dramatic reduction of *Crithidia*
51
52 121 gut pathogen cells. First, we isolated major secondary compounds of sunflower pollen. We then
53
54 122 synthesized one compound and purchased another and tested their effects on *Crithidia* infection
55
56 123 in the bumble bee *B. impatiens* in separate experiments. In addition, we assessed the effects of an
57
58
59
60
61
62
63
64
65

array of fatty acids, singly and in combination, to determine their effects on *Crithidia*. If we can discover a chemical mechanism underlying the effect of sunflower pollen on pathogen infection, this could lead to breeding programs selecting for these traits in agricultural crops or choosing wild plants for pollinator habitat partially based on pollen composition.

METHODS AND MATERIALS

Study System. *Bombus impatiens* (Apidae), or the common eastern bumble bee, is one of the most abundant bumble bee species in the eastern United States (Cordes et al. 2012; Gillespie 2010). *Crithidia bombi* (Trypanosomatidae; *Crithidia* hereafter) is a gut pathogen of bumble bees that can infect up to 80% of *Bombus* at sites in western Massachusetts (Gillespie 2010) and up to 80% of *B. terrestris* workers near Basel, Switzerland (Shykoff and Schmid-Hempel 1991). *Crithidia* can be relatively benign when bees have sufficient resources (Brown et al. 2003), but under food-limited conditions can reduce queen fitness and colony performance (Brown et al. 2003) and worker survival (Brown et al. 2000). In the field, infection with *Crithidia* was associated with reduced reproduction in wild colonies (Goulson et al. 2018).

Helianthus annuus, or sunflower, is native to North America (Reagon and Snow 2006) and is a major oilseed crop worldwide whose yield is improved by bee visitation (reviewed in Nicolson and Human 2013). Buckwheat pollen (*Fagopyrum esculentum*) has protein content comparable to sunflower (Yang et al. 2013) but bees that consume it have much higher *Crithidia* cell counts (Giacomini et al. 2018), and so it can be used as a control pollen treatment to compare diets with comparable protein but different effects on *Crithidia*.

Chemical Analysis. We analyzed *H. annuus* pollen sourced from China (Changge Hauding Wax Industry, China Co. LTD). Pollen samples (50 mg) were sonicated in 100% methanol for 10 min and left to extract at room temperature for 24 hr. Each sample was analyzed by LC–MS on a Waters Alliance LC solvent delivery system with a ZQ MS detector. Compounds were separated over a Phenomenex Luna C18(2) column (150 x 4.0 mm i.d., 5 µm particle size) operating under gradient elution conditions, with A = MeOH, B = H₂O, C = 1 % HCO₂H in MeCN; A = 0 %, B = 90 % at t = 0 min; A = 90 %, B = 0 % at t = 20 min; A = 90 %, B = 0 % at t = 30 min; A = 0 %, B = 90 % at t = 31 min; column temperature 30 °C and flow rate of 0.5 ml min⁻¹. Analysis of our samples by this method identified two major peaks eluting after 10.4 and 11.0 min as a quercetin-3-*O*-hexoside and a quercetin-3-*O*-(6-*O*-malonyl)-hexoside respectively. These compounds were characterised by a UV spectrum that was similar to that of rutin (quercetin-3-*O*-rutinoside) with band maxima at 357 and 260 nm and pseudo molecular ions at $m/z = 463$ and $549 [M-H]^-$ in negative mode and 465 and $551 [M+H]^+$ in positive mode respectively. Fragmentation in the mass spectrometer in positive mode revealed ions at $m/z = 303$ in both compounds corresponding to the aglycone [quercetin+H]⁺. These identifications correspond with those of Kostic et. al. (2019).

Two additional major peaks were recorded in the LC-MS chromatogram of the sunflower pollen methanol extract that eluted between 13 and 15 min and corresponded to two triscoumaroyl spermidines. These compounds were characterised by pseudomolecular ions with $m/z = 584 [M+H]^+$ and a UV spectrum similar to that recorded in our laboratory (Kew) for coumaric acid derivatives (with a band max = 313 nm). The most abundant triscoumaroyl spermidine co-eluted and showed similar fragmentation, including three losses of a $m/z = 146.03$

fragment in the mass spectrometer, to N¹, N⁵, N¹⁰- tri(*E*)-*p*-coumaroyl spermidine which was synthesised in our laboratory (Online Resource 1). Minor components were also recorded in the chromatogram with similar pseudo-molecular ions and fragmentation in the mass spectrometer and were likely to be structural analogues of the main compound, with various E and Z isomerization at the propenyl of the three coumaroyl groups since high levels of variation in isomerization of the cinnamoyls in these compounds are reported (Kite et al. 2013).

Polynomial calibration curves for each compound via quantification of the [M+H]⁺ molecular ion of a commercial standard of rutin (Sigma-Aldrich, Dorset, UK) and a sample of triscoumaroyl spermidine synthesized in our laboratory were used to quantify compounds. The two quercetin glycosides occurred at 0.56 and 1.37 mg/g pollen while the mean concentration of the triscoumaroyl spermidine was 12.16 mg/g pollen.

Bee Bioassay General Approach. The general approach was similar for all assays and followed previously published protocols (Giacomini et al. 2018; LoCascio et al. 2019a; Richardson et al. 2015). Commercial experimental colonies (BioBest, Leamington, Canada) were confirmed to be *Crithidia*-free by biweekly subsampling, and the *Crithidia* strain used in these assays was originally collected in 2015 from infected wild *B. impatiens* at a farm in Hadley, Massachusetts USA (42°21'51.93"N, 72°33'55.88"W) and maintained in commercial colonies thereafter. All colonies were fed mixed wildflower pollen (Koppert Biological Systems, Howell, Michigan, USA or CC Pollen Inc., Phoenix, Arizona, USA). Worker bees were inoculated after a 1-2 hr starving period with either 10 µl (secondary compounds) or 15 µl (fatty acids) of a 25% sucrose solution containing 600 cells/µl, well within the range of natural fecal *Crithidia* concentrations (Otterstatter and Thomson 2006). Upon inoculation, bees were placed in individual 18.5 ml vials

(secondary compounds) or Placon 473 ml cups (fatty acids) and assigned to diet treatments that they received for 7 days along with 30% sucrose solution that was replaced daily (secondary compounds) or every other day (fatty acids). Bees were maintained in darkness at 27°C in an incubator during assays. After 7 days, bees were dissected to remove the gut, which was ground in 300 µl of ¼ strength Ringer's solution (Sigma-Aldrich – Fluka 96724). After 4 hr, a 10 µl subsample was removed and moving *Crithidia* cells were counted on a 0.02 µl field of vision on the grid of a hemacytometer.

For the secondary compound assays we used callows, which are bees that had emerged from pupae within the last 24 hr. We isolated pupal clumps from colonies, collected callows as they emerged and fed them wildflower pollen for two days before inoculating them to enter the experiment. Because callows emerged without nestmates, this was an extra precaution to ensure they were free of *Crithidia* (which infects adults), but these bees also did not receive gut microbiota from their nestmates that could influence interactions with *Crithidia* (Koch and Schmid-Hempel 2011). For the fatty acid assays, we pulled adult workers directly from colonies and inoculated them to enter the experiment. This method does not control for worker age, but allows bees to acquire their colony's gut microbiome. Within each assay, bees with pollen treatments were compared to controls under the same conditions, so while our methods varied, we can still assess diet effects within each experiment. We also included a measure of bee size for all assays since smaller bees often have higher infections (e. g., Richardson et al. 2015). For the secondary compound assays we used callow mass, and for fatty acids we used radial cell length, which is highly correlated with other measures of body size (Nooten and Rehan 2020). In the triscoumaryl spermidine experiment, we measured both callow mass and radial cell length and they were highly correlated (Pearson's $r = 0.85$, $n = 259$).

For all assays we recorded daily mortality, and measured daily pollen and sucrose consumption once per bee, except for omitting sucrose consumption in the triscoumaroyl spermidine assay. Pollen and sucrose rations were typically weighed upon providing them to bees and then approximately 24 hr later (48 hr in the first fatty acid assay).

Secondary Chemical Bioassays. We tested the effects of two major secondary compounds found in sunflower pollen, a triscoumaroyl spermidine and rutin, at four concentrations each. We used rutin as a proxy for the quercetin glycosides identified in sunflower pollen because rutin is also a quercetin glycoside and available commercially. Quercetin glycosides are reported to show equivalent activity across a range of analogues against microorganisms including trypanosomes (da Silva et al. 2019; Marin et al. 2017), suggesting that rutin provides a suitable proxy for other quercetin glycosides.

Tri-*p*-coumaroyl spermidine ('triscoumaryl spermidine' hereafter) was synthesized in the laboratory as described in Online Resource 1, while rutin was purchased commercially (Sigma-Aldrich, R5143-50G). Both were added to buckwheat pollen at 1% (very high), 0.1% (high), 0.01% (medium), or 0.001% (low) concentrations wt/wt. For context, we found 12 mg/g of triscoumaroyl spermidine in pollen (1.2%; comparable with the 'very high' treatment) and 0.2% quercetin glycosides in pollen, similar to the 'high' level. Thus, our concentrations are within natural ranges. Compounds were mixed with buckwheat pollen and compared to pure buckwheat pollen as a negative control and sunflower pollen as a positive control (sunflower and buckwheat pollen came from Changge Huading Wax Industry Co., Ltd., Henan, China; except that buckwheat came from Fuyang Import and Export Ltd for the final fatty acid assay). Compounds were added to dry pollen, which was then mixed with distilled water for all treatments to create a

paste that was frozen at -20°C until use. Details of mixing secondary compounds with pollen and preparing pollen balls are in Online Resource 2.

The triscoumaryl spermidine assay was conducted from July 17 through August 19, 2015, using 108 bees from three experimental colonies, ultimately including 15-20 bees/treatment. The rutin experiment was conducted from November 3, 2015 through April 7, 2016 and used 292 bees from 12 experimental colonies, ultimately including 45-54 bees/treatment.

Fatty Acid Bioassays. We tested the effects of several fatty acids singly and in combination. We chose the most common ones detected in sunflower pollen that were commercially available, and tested each at typical concentrations based on published literature (Nicolson and Human 2013; Yang et al. 2013). We conducted three assays that each tested individual or combinations of fatty acids added directly to control pollen as powders or liquids. Compounds tested in the first assay were linoleic (1.67 mg/g; CAS 60-33-3), lauric (12.39 mg/g; CAS 143-07-7), caprylic (1.46 mg/g; CAS 124-07-2), palmitic (8.46 mg/g; CAS 57-10-3), and decanoic (0.14 mg/g; CA: 334-48-5) acids. The second assay included linolenic (7.61 mg/g; CAS 463-40-1), myristic (1.62 mg/g; CAS 544-63-8), stearic (0.65 mg/g; CAS 57-11-4) and oleic (5.39 mg/g; CAS 112-80-1) acids. The third assay tested all nine fatty acids combined in a single treatment (using the same concentrations as in individual assays) and sunflower oil, which contains a mix of fatty acids (37.07 mg/g; Organic 365 Everyday Value Products, Whole Foods Market, Austin, Texas, USA), to assess whether fatty acids had interactive effects that would not be detected in assays of individual compounds. Details of suppliers and concentration calculations are provided in Online Resource 2. Fatty acids were added to dry pollen with a coffee grinder, and then distilled water was added in a 6:1 or 7:1 (third assay only) ratio to make a paste. As with the secondary

chemical assays, compounds were added to buckwheat pollen and compared to buckwheat and sunflower pollen as negative and positive controls, respectively. However, in the second assay we did not have sufficient buckwheat pollen and instead added fatty acids to mixed wildflower pollen (CC Pollen Inc. Phoenix, Arizona, USA) and used wildflower pollen as the negative control.

We conducted the first assay from September 28, 2018 to February 6, 2019 using 246 bees from 14 experimental colonies, including 32-37 bees per fatty acid treatments and 47 in the buckwheat control. The second was conducted from February 20 to April 12, 2019, using 167 bees from 5 experimental colonies, including 25-31 bees/treatment. The third assay was conducted from July 18 to 26, 2019 using 106 bees from 4 experimental colonies, including 25-27 bees/treatment.

Statistical Analysis. We used the open source software R v3.3.3 (R Core Team 2014). All data CSV files and R scripts are available from the Dryad Digital Repository (www.datadryad.org); [NNNN] (Adler et al. 202X). For all analyses, the significance of model terms was tested with likelihood ratio χ^2 tests, conducted with the Anova function, which compares relative goodness of fit between models with and without each term (car package, Fox and Weisberg 2019). To analyze the number of *Crithidia* cells counted in a 0.02- μ l gut extract as the response, we used generalized linear mixed models (Bates et al. 2015). Cell counts were over-dispersed, and thus we used a negative binomial error distribution and the model function glmer.nb (lme4 package, Bates et al. 2015). Each model included diet treatment and callow mass or wing marginal cell size as fixed effects. Colony of origin was included as a random effect, along with inoculation date to account for the independent preparation of inoculation on each trial date. When diet

treatment was significant, means were compared with post-hoc Tukey's tests using the `glht` function (`multcomp` package, Hothorn et al. 2008). To analyze the amount of pollen and sucrose consumed, we used general linear models with diet and callow mass or wing marginal cell size as fixed effects and colony of origin and inoculation date as random effects. To analyze death hazard rate by pollen treatment, we used a survival analysis with a cox proportional hazards mixed model using the `coxme` function (`coxme` package, Therneau 2019), `survfit`, and `Surv` functions (`survival` package, Therneau 2015; Therneau and Grambsch 2000)) including diet treatment, callow mass or wing marginal cell length, date of inoculation and colony of origin as predictors. In the rutin experiment, inoculation date was not recorded for dead bees and so was not included in the survival analysis. In the two individual fatty acid assays, we did not measure wing cell length on dead bees and so did not include this predictor in the survival analysis. All figures were made using `lsmeans` function (`lsmeans` package, Lenth 2016)) and `ggplot` function (`ggplot2` package, Wickham 2016).

RESULTS

Crithidia cells replicated in hosts; final raw counts in the control treatments across assays ranged from 29.7 – 59.9 cells per 0.02 μ l. On average, this is approximately 100 times more *Crithidia* cells than the initial inoculation.

In every assay, diet significantly affected *Crithidia* cell counts (combined fatty acid assay: $\chi^2 = 16.02$, $P = 0.007$; all other assays: $\chi^2 > 21.3$, $P < 0.001$). However, in every case this effect was driven by the difference between the sunflower pollen and all other treatments (Figures 1 and 2); sunflower pollen reduced *Crithidia* counts by 64-94% per assay compared to

the control treatment. None of the secondary compounds or fatty acids, singly or in combination, significantly reduced *Crithidia* cell counts compared to the control treatments, and all were significantly higher than *Crithidia* counts with sunflower pollen.

Diet did not affect mortality in any assay ($\chi^2 < 4.3$, $P > 0.5$ for all), suggesting that neither sunflower pollen nor any of the compounds tested had notable costs or benefits for bee survival. In the rutin and combined fatty acid assays, smaller bees had higher mortality (rutin: $\chi^2 = 14.39$, $P < 0.001$; combined fatty acid: $\chi^2 = 8.47$, $P = 0.004$) but there was no relationship between bee size and mortality in the triscoumaroyl spermidine assay ($\chi^2 = 1.63$, $P = 0.214$); relationships between bee size and mortality could not be assessed for the other two assays because wings were not collected from dead bees.

Diet treatments also did not affect consumption of pollen ($\chi^2 < 5.3$, $P > 0.38$ for all) or sucrose (rutin assay: $\chi^2 = 9.99$, $P = 0.075$; all others: $\chi^2 < 5.9$, $P > 0.43$). Larger bees consumed more pollen than smaller bees in all experiments ($\chi^2 > 5.8$, $P < 0.017$ for all) and also consumed more sucrose in all assays in which this was measured ($\chi^2 > 4.8$, $P < 0.03$ for all).

DISCUSSION

We isolated major secondary compounds of sunflower and tested their efficacy against a common bumble bee gut pathogen; we also assessed the effects of numerous fatty acids from sunflower pollen, singly and in combination. Although we found that sunflower pollen consistently reduced *Crithidia* compared to control pollens, matching previous work (Giacomini et al. 2018; LoCascio et al. 2019a; LoCascio et al. 2019b), none of the compounds we tested appear to be the sole mechanism driving this result, as none of them significantly reduced

pathogen counts relative to control pollens. Although we did not test protein content in this study, it seems unlikely that protein content is the underlying mechanism since protein concentrations are similarly low in buckwheat and sunflower pollen (Yang et al. 2013) but outcomes of these diets for *Crithidia* infection are very different. Rather than focusing on specific compounds, it may be more efficient in the future to assess potential chemical mechanisms of sunflower pollen by testing sequential extractions to more rapidly facilitate identification of the key biologically active components. Progressing from apolar to more polar fractions (to selectively extract compounds first that can be most easily isolated via evaporation) and adding each to control pollen to assess effects against *Crithidia* may provide a more efficient method to identify the specific bioactive compounds or combinations that make sunflower so consistently effective against infection by this pathogen. Alternatively, the slow digestibility of pollen could result in compounds being naturally released in the hind gut where infection occurs. Mixing commercial compounds with pollen might cause earlier release, which could be less effective. However, some secondary compounds are metabolized in the mid-or hind-gut and early exposure in the crop can have prophylactic effects on infection (Koch et al. 2019). Further work is needed to elucidate whether sunflower pollen digestion and metabolism processes play a role in its effect on infection.

The mechanism underlying the interaction between sunflower pollen and *Crithidia* could be mechanical rather than chemical. For example, the roughness of some plant leaves can rid chimpanzees of intestinal parasites via scouring intestinal walls (Huffman 1997), a mechanical mechanism. Sunflower and other Asteraceae pollen is notable for spines on the outer coat (Blackmore et al. 2009). *Crithidia* is a gut parasite that establishes infection by attaching to the ileum wall, a portion of the hindgut (Gorbunov 1996; Koch et al. 2019). An exciting recent study

showed that a compound from heather nectar (*Calluna vulgaris*) dramatically reduced *Crithidia* infection in *B. terrestris* by removing the flagellum, preventing attachment (Koch et al. 2019). Although we do not know how spines could interfere in attachment, it is possible that spines make it difficult for cells to attach to the gut wall, which we can examine in future work. We note that in Koch et al. (2019), nectar compounds were not effective at reducing infections that were already established, likely because compounds were metabolized before reaching the hindgut where infection occurs. By contrast, sunflower is effective even after infection has established (Giacomini et al. 2018), suggesting a mechanism acting in the hindgut itself. Interestingly, species in the family Malvaceae also have spiny pollen that is hypothesized to be a defense preventing consumption by bees (Lunau et al. 2015). Pollen from hollyhocks, *Alcea rosea*, were not collected by *B. terrestris* unless spines and sticky pollenkitt were removed; extracted chemicals did not affect pollen collection, suggesting that deterrence was due to mechanical rather than chemical mechanisms (Lunau et al. 2015). Future work can use a similar approach or take advantage of sunflower pollen ‘shells’ used for nanodrug delivery (Mundargi et al. 2016) to dissect the role of mechanical and chemical mechanisms underlying the effect on *Crithidia*.

We identified flavonoid glycosides and triscoumaroyl spermidines in high quantities as the major secondary metabolites in the sunflower pollen. The occurrence of flavonoids in pollen and honey is well reported (Ciappini 2019; Palmer-Young et al. 2019a; Palmer-Young et al. 2019b) so their occurrence in sunflower pollen was not unexpected. Indeed, Kostic et al. (2019) recently analyzed bee-collected pollen from sunflower and also identified flavonoid glycosides in methanol extracts of sunflower pollen including a quercetin 3-*O*-(6-*O*-acetyl)-hexoside as the major flavonoid component, mirroring the occurrence of the quercetin-3-*O*-(6-*O*-malonyl)-

hexoside identified here. It is likely that the acetylated quercetin hexoside recorded by Kostic et al. (2019) was in fact a decarboxylated derivative and occurred naturally as a malonylated analogue, as identified in our material and reported here. Facile decarboxylation in solution of malonyl to acetyl has been reported, and naturally occurring acetyl glycosides are rare in nature compared to malonyl derivatives (Agrawal 1992; Stevenson and Veitch 1996).

Triscoumaroyl spermidines have not previously been reported in sunflower pollen extracts but are otherwise known to occur widely although not ubiquitously in the pollen of angiosperms (Kite et al. 2013; Martin-Tanguy et al. 1978; Palmer-Young et al. 2019a; Palmer-Young et al. 2019b). Their role in flower growth and reproduction has been determined using metabolomic and genomic studies (Fellenberg et al. 2009; Hanhineva et al. 2008), while deficiency of spermidine conjugates caused pollen grains to become deformed, indicating a developmental role for these compounds (Grienenberger et al. 2009). Their potential as plant defense compounds against fungi has been demonstrated (Bassard et al. 2010; Martin-Tanguy et al. 1978; Walters et al. 2001), and although some compounds show biological activity against insect neuroreceptors, they have low toxicity in bioassays when ingested by insects (Fixon-Owoo et al. 2003). Mori et al. (2019) suggest their occurrence in apricot pollen (*Prunus mume* L.) was associated with non-fluorescing pollen so they may absorb UV light and protect the male gamete against harmful UV radiation (Gill and Tuteja 2010). Despite the variety of biological activities attributable to acyl spermidines and quercetin glycosides, they do not explain the biological activity of sunflower pollen against *Crithidia*.

Fatty acids are important components of the bee diet (Vaudo et al. 2016). They typically occur in the pollen kit (Chichiricco et al. 2019) and can make up to 10% of the dry weight of pollen (Arien et al. 2015), providing a significant dietary component for larvae and adult bumble

1
2
3
4 400 bees. Overall, the most common free fatty acids in pollen are α -linolenic acid, α -linoleic acid,
5
6 401 palmitic acid, stearic acid, and oleic acid (Manning 2001). However, they differ in sunflower,
7
8
9 402 with myristic acid dominant and including palmitic, lauric, stearic and caprylic acids (Kostic et
10
11 403 al. 2019). The role of lipids in colony growth and individual bee behaviors has been studied in
12
13
14 404 honeybees and commercial bumble bees (Arien et al. 2015; Muth et al. 2018), but how they
15
16 405 influence the development or establishment of bee pathogens such as *Crithidia* is unknown. Fatty
17
18
19 406 acids have antimicrobial properties (Manning 2001) and several analogues of myristic acid, a
20
21 407 common saturated fatty acid of plant oils, have potent anti-trypanosomal activity (Doering et al.
22
23
24 408 1994). Thus, we hypothesized that pollen fatty acids could contribute to the bioactivity of
25
26 409 sunflower pollen against *Crithidia*. However, we found no activity among the fatty acids tested
27
28
29 410 here. The most active myristate derivatives reported by Doering et al. (1994) were oxidized and
30
31 411 sulfonated analogues of myristic acid rather than myristic acid itself, suggesting that the common
32
33
34 412 fatty acids occurring widely in floral pollen as saturated or unsaturated fats may not have the
35
36 413 structural characters to challenge *Crithidia*.

37
38 414 If we can discover the mechanism by which sunflower pollen reduces bee pathogen
39
40
41 415 infection, this can open new avenues for management of bee health. Because sunflower pollen
42
43 416 has low protein, if the mechanism was known we could search for this trait in other pollens to
44
45
46 417 identify potential species that are more nutritionally beneficial. If we discover chemical
47
48 418 components of pollen responsible for reducing pathogen infection, this could lead to breeding
49
50
51 419 programs selecting for these traits in agricultural crops or choosing wild plants for pollinator
52
53 420 habitat partially based on these traits. Other recent work in our labs showed that blueberry
54
55
56 421 domestication reduced levels of a nectar caffeic acid below the threshold that is effective in
57
58 422 reducing a bee pathogen (Egan et al. 2018), suggesting that there is an opportunity to incorporate
59
60
61
62
63
64
65

selective breeding of crop floral traits to improve bee health. If the mechanism is chemical, another possibility would be isolating or synthesizing those compounds for incorporation into commercial bee diet. We note that sunflower pollen often leads to poor performance of generalist bees such as honey bees (Human et al. 2007; Nicolson et al. 2018; Nicolson and Human 2013). and bumble bees (McAulay and Forrest 2019; Tasei and Aupinel 2008), although this is not always the case (Giacomini et al. 2018; Treanore et al. 2019). However, mixing sunflower pollen with other pollens can ameliorate negative effects on worker survival (McAulay and Forrest 2019). Thus, sunflower pollen could be an effective treatment for *Crithidia* infection without sacrificing bee performance if it were mixed with other pollens (Giacomini et al., in preparation), but isolating the underlying mechanism may open up new avenues for development of supplements that could be added to more nutritious pollen diets.

Acknowledgements- We thank E. Amponsah, L. Cleary, J. Cook, L. Gagnon, Z. Henry, K. Michaud, J. Roch, and C. Sylvia for assistance with experiments, and two anonymous reviewers for feedback on the manuscript. Funding came from USDA-AFRI 2013-02536 (LSA, REI and PCS), USDA-NIFA-2016-07962 and USDA-NIFA-2018-08591 (both to LSA and REI), the USDA/CSREES (Multi-state Hatch) MAS00497 (LSA), and the Peter Sowerby Foundation in the UK (PCS).

REFERENCES

Agrawal PK (1992) NMR-spectroscopy in the structural elucidation of oligosaccharides and glycosides *Phytochemistry* 31:3307-3330 doi:10.1016/0031-9422(92)83678-r

- Alaux C, Ducloz F, Crauser D, Le Conte Y (2010) Diet effects on honeybee immunocompetence. *Biol Lett* 6:562-565 doi:10.1098/rsbl.2009.0986
- Arien Y, Dag A, Zarchin S, Masci T, Shafir S (2015) Omega-3 deficiency impairs honey bee learning. *Proc Natl Acad Sci USA* 112:15761-15766 doi:10.1073/pnas.1517375112
- Arrese EL, Soulages JL (2010) Insect fat body: Energy, metabolism, and regulation. *Annu Rev Entomol* 55:207-225 doi:10.1146/annurev-ento-112408-085356
- Baracchi D, Brown MJF, Chittka L (2015) Behavioural evidence for self-medication in bumblebees? [version 3; peer review: 3 approved]. *F1000Research* 4:1-15 doi:<https://doi.org/10.12688/f1000research.6262.3>
- Bassard JE, Ullmann P, Bernier F, Werck-Reichhart D (2010) Phenolamides: Bridging polyamines to the phenolic metabolism. *Phytochemistry* 71:1808-1824 doi:10.1016/j.phytochem.2010.08.003
- Bates D, Maechler M, Bolker BM, Walker S (2015) Fitting linear mixed-effects models Using lme4. *Journal of Statistical Software* 67:1-48 doi:doi:10.18637/jss.v067.i01
- Biller OM, Adler LS, Irwin RE, McAllister C, Palmer-Young EC (2015) Possible synergistic effects of thymol and nicotine against *Crithidia bombi* parasitism in bumble bees. *Plos One* 10 doi:10.1371/journal.pone.0144668
- Blackmore S, Wortley AH, Skvarla JJ, Robinson H (2009) Evolution of pollen in the Compositae. In: Funk VA, Susanna A, Stuessy TF, Bayer RJ (eds) *Systematics, evolution and biogeography of Compositae*. IAPT, Vienna, pp 101–130
- Brown MJF, Loosli R, Schmid-Hempel P (2000) Condition-dependent expression of virulence in a trypanosome infecting bumblebees. *Oikos* 91:421-427 doi:10.1034/j.1600-0706.2000.910302.x

- 469 Brown MJF, Schmid-Hempel R, Schmid-Hempel P (2003) Strong context-dependent virulence
470 in a host-parasite system: reconciling genetic evidence with theory. *J Anim Ecol* 72:994-
471 1002 doi:10.1046/j.1365-2656.2003.00770.x
- 472 Cameron SA, Lim HC, Lozier JD, Duennes MA, Thorp R (2016) Test of the invasive pathogen
473 hypothesis of bumble bee decline in North America. *Proc Natl Acad Sci USA* 113:4386-
474 4391 doi:10.1073/pnas.1525266113
- 475 Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, Griswold TL (2011)
476 Patterns of widespread decline in North American bumble bees. *Proc Natl Acad Sci USA*
477 108:662-667 doi:10.1073/pnas.1014743108
- 478 Chichiricco G, Pacini E, Lanza B (2019) Pollenkitt of some monocotyledons: lipid composition
479 and implications for pollen germination. *Plant Biol* 21:920-926 doi:10.1111/plb.12998
- 480 Ciappini MC (2019) Polyphenolic profile of floral honeys in correlation with their pollen
481 spectrum. *J Apic Res* 58:772-779 doi:<https://doi.org/10.1080/00218839.2019.1654967>
- 482 Cook D, Manson JS, Gardner DR, Welch KD, Irwin RE (2013) Norditerpene alkaloid
483 concentrations in tissues and floral rewards of larkspurs and impacts on pollinators.
484 *Biochem Syst Ecol* 48:123-131
- 485 Cordes N, Huang W-F, Strange JP, Cameron SA, Griswold TL, Lozier JD, Solter LF (2012)
486 Interspecific geographic distribution and variation of the pathogens *Nosema bombi* and
487 *Crithidia* species in United States bumble bee populations. *J Invertebr Pathol* 109:209-
488 216
- 489 da Silva ER, Brogi S, Lucon JF, Campiani G, Gemma S, Maquiaveli CD (2019) Dietary
490 polyphenols rutin, taxifolin and quercetin related compounds target *Leishmania*
491 *amazonensis* arginase. *Food & Function* 10:3172-3180 doi:10.1039/c9fo00265k

- Di Pasquale G et al. (2013) Influence of pollen nutrition on honey bee health: Do pollen quality and diversity matter? Plos One 8 doi:10.1371/journal.pone.0072016
- Doering TL, Lu TB, Werbovetz KA, Gokel GW, Hart GW, Gordon JI, Englund PT (1994) Toxicity of myristic acid analogs toward African trypanosomes Proc Natl Acad Sci USA 91:9735-9739 doi:10.1073/pnas.91.21.9735
- Egan PA, Adler LS, Irwin RE, Farrell IW, Palmer-Young EC, Stevenson PC (2018) Crop domestication alters floral reward chemistry with potential consequences for pollinator health. Frontiers in Plant Science 9 doi:10.3389/fpls.2018.01357
- Farag RS, Youssef AM, Ewies MA, Hallabo SAS (1978) Long-chain fatty acids of 6 pollens collected by honeybees in Egypt J Apic Res 17:100-104 doi:10.1080/00218839.1978.11099911
- Fatrcova-Sramkova K, Nozkova J, Mariassyova M, Kacaniova M (2016) Biologically active antimicrobial and antioxidant substances in the *Helianthus annuus* L. bee pollen. Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes 51:176-181 doi:10.1080/03601234.2015.1108811
- Feldlaufer MF, Knox DA, Lusby WR, Shimanuki H (1993) Antimicrobial activity of fatty acids against *Bacillus larvae*, the causative agent of American foulbrood disease Apidologie 24:95-99 doi:10.1051/apido:19930202
- Fellenberg C, Bottcher C, Vogt T (2009) Phenylpropanoid polyamine conjugate biosynthesis in *Arabidopsis thaliana* flower buds. Phytochemistry 70:1392-1400 doi:10.1016/j.phytochem.2009.08.010
- Fixon-Owoo S et al. (2003) Preparation and biological assessment of hydroxycinnamic acid amides of polyamines. Phytochemistry 63:315-334 doi:10.1016/s0031-9422(03)00133-x

- 1
2
3
4 515 Fox J, Weisberg S (2019) An R Companion to Applied Regression, Third Edition. . Thousand
5
6
7 516 Oaks CA: Sage. ,
8
9 517 Gherman BI et al. (2014) Pathogen-associated self-medication behavior in the honeybee *Apis*
10
11 518 *mellifera*. Behav Ecol Sociobiol 68:1777-1784 doi:10.1007/s00265-014-1786-8
12
13
14 519 Giacomini JJ, Leslie J, Tarpay DR, Palmer-Young EC, Irwin RE, Adler LS (2018) Medicinal
15
16 520 value of sunflower pollen against bee pathogens. Scientific Reports 8:14394
17
18
19 521 doi:10.1038/s41598-018-32681-y
20
21 522 Gill SS, Tuteja N (2010) Polyamines and abiotic stress tolerance in plants. Plant Signaling &
22
23 523 Behaviour 5:26-33 doi:<https://doi.org/10.4161/psb.5.1.10291>
24
25
26 524 Gillespie S (2010) Factors affecting parasite prevalence among wild bumblebees. Ecol Entomol
27
28 525 35:737-747 doi:10.1111/j.1365-2311.2010.01234.x
29
30
31 526 Gorbunov PS (1996) Peculiarities of life cycle in flagellate *Crithidia bombi* (Protozoa,
32
33 527 Trypanosomatidae). Zoologicheskyy Zhurnal 75:803-810
34
35
36 528 Goulson D, Nicholls E, Botías C, Rotheray EL (2015) Bee declines driven by combined stress
37
38 529 from parasites , pesticides , and lack of flowers. Science 2010:1-16
39
40
41 530 Goulson D, O'Connor S, Park KJ (2018) The impacts of predators and parasites on wild
42
43 531 bumblebee colonies. Ecol Entomol 43:168-181 doi:10.1111/een.12482
44
45
46 532 Grienemberger E et al. (2009) A BAHD acyltransferase is expressed in the tapetum of
47
48 533 *Arabidopsis* anthers and is involved in the synthesis of hydroxycinnamoyl spermidines.
49
50 534 Plant J 58:246-259 doi:10.1111/j.1365-313X.2008.03773.x
51
52
53 535 Hanhineva K, Rogachev I, Kokko H, Mintz-Oron S, Venger I, Karenlampi S, Aharoni A (2008)
54
55 536 Non-targeted analysis of spatial metabolite composition in strawberry (*Fragaria x*
56
57 537 *ananassa*) flowers. Phytochemistry 69:2463-2481 doi:10.1016/j.phytochem.2008.07.009
58
59
60
61
62
63
64
65

- Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biometrical Journal* 50:346-363
- Huffman MA (1997) Current evidence for self-medication in primates: A multidisciplinary perspective. In: Steegmann AT (ed) Yearbook of Physical Anthropology, Vol 40 - 1997, vol 40. Yearbook of Physical Anthropology. pp 171-200
- Human H, Nicolson SW, Strauss K, Pirk CWW, Dietemann V (2007) Influence of pollen quality on ovarian development in honeybee workers (*Apis mellifera scutellata*). *J Insect Physiol* 53:649-655 doi:10.1016/j.jinsphys.2007.04.002
- Kite GC, Larsson S, Veitch NC, Porter EA, Ding N, Simmonds MSJ (2013) Acyl spermidines in inflorescence extracts of elder (*Sambucus nigra* L., Adoxaceae) and elderflower drinks. *J Agric Food Chem* 61:3501-3508 doi:10.1021/jf304602q
- Koch H, Schmid-Hempel P (2011) Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proc Natl Acad Sci USA* 108:19288-19292 doi:10.1073/pnas.1110474108
- Koch H, Woodward J, Langat MK, Brown MJF, Stevenson PC (2019) Flagellum removal by a nectar metabolite inhibits infectivity of a bumblebee parasite. *Curr Biol* 29:3494–3500 doi:<https://doi.org/10.1016/j.cub.2019.08.037>
- Kostic AZ, Milincic DD, Gasic UM, Nedic N, Stanojevic SP, Tesic ZL, Pesic MB (2019) Polyphenolic profile and antioxidant properties of bee-collected pollen from sunflower (*Helianthus annuus* L.) plant. *LWT-Food Science and Technology* 112 doi:10.1016/j.lwt.2019.06.011
- Kostic AZ, Pesic MB, Trbovic D, Petronijevic R, Dramicanin AM, Milojkovic-Opsenica DM, Tesic ZL (2017) The fatty acid profile of Serbian bee-collected pollen - a

- chemotaxonomic and nutritional approach. J Apic Res 56:533-542
doi:10.1080/00218839.2017.1356206
- Lenth RV (2016) Least-Squares Means: The R Package lsmeans. Journal of Statistical Software 69 doi:doi: 10.18637/jss.v069.i01
- LoCascio GM, Aguirre L, Irwin RE, Adler LS (2019a) Pollen from multiple sunflower cultivars and species reduces a common bumblebee gut pathogen. Royal Society Open Science 6:190279 doi:<http://dx.doi.org/10.1098/rsos.190279>
- LoCascio GM, Pasquale R, Amponsah E, Irwin RE, Adler LS (2019b) Effect of timing and exposure of sunflower pollen on a common gut pathogen of bumble bees. Ecol Entomol 44:702-710 doi:10.1111/een.12751
- Lunau K, Piorek V, Krohn O, Pacini E (2015) Just spines-mechanical defense of malvaceous pollen against collection by corbiculate bees. Apidologie 46:144-149
doi:10.1007/s13592-014-0310-5
- Manning R (2001) Fatty acids in pollen: a review of their importance for honey bees. Bee World 82:60-75
- Manson JS, Otterstatter MC, Thomson JD (2010) Consumption of a nectar alkaloid reduces pathogen load in bumble bees. Oecologia 162:81-89
- Mao W, Schuler MA, Berenbaum MR (2013) Honey constituents up-regulate detoxification and immunity genes in the western honey bee *Apis mellifera*. Proc Natl Acad Sci USA 110:8842-8846 doi:10.1073/pnas.1303884110
- Marin C et al. (2017) Antitrypanosomatid activity of flavonoid glycosides isolated from *Delphinium gracile*, *D-staphisagria*, *Consolida oliveriana* and from *Aconitum napellus* subsp *Lusitanicum*. Phytochemistry Letters 19:196-209 doi:10.1016/j.phytol.2016.12.010

- 584 Martin-Tanguy J, Cabanne F, Perdrizet E, Martin C (1978) Distribution of hydroxycinnamic
585 acid-amides in flowering plants *Phytochemistry* 17:1927-1928 doi:10.1016/s0031-
586 9422(00)88735-x
- 587 McAulay MK, Forrest JRK (2019) How do sunflower pollen mixtures affect survival of
588 queenless microcolonies of bumblebees (*Bombus impatiens*)? *Arthropod-Plant*
589 *Interactions* 13:517-529 doi:10.1007/s11829-018-9664-3
- 590 McKey D (1979) The distribution of secondary compounds within plants. In: Rosenthal GA,
591 Janzen DH (eds) *Herbivores: their interaction with secondary plant metabolites*.
592 Academic Press, New York, pp 56-134
- 593 Mori S, Akamatsu M, Fukui H, Tsukioka J, Goto K, Hirai N (2019) The unusual conformational
594 preference of N-1,N-5,N-10-tri-p-coumaroylspermidine E-Z isomers from the Japanese
595 apricot tree, *Prunus mume*, for the (ZZZ)-form. *Phytochemistry Letters* 31:131-139
596 doi:10.1016/j.phytol.2019.02.028
- 597 Mundargi RC, Potroz MG, Park S, Shirahama H, Lee JH, Seo J, Cho NJ (2016) Natural
598 sunflower pollen as a drug delivery vehicle. *Small* 12:1167-1173
599 doi:10.1002/smll.201500860
- 600 Muth F, Breslow PR, Masek P, Leonard AS (2018) A pollen fatty acid enhances learning and
601 survival in bumblebees. *Behav Ecol* 29:1371-1379 doi:10.1093/beheco/ary111
- 602 Nicolson SW, Das Neves SD, Human H, Pirk CWW (2018) Digestibility and nutritional value of
603 fresh and stored pollen for honey bees (*Apis mellifera scutellata*). *J Insect Physiol*
604 107:302-308 doi:10.1016/j.jinsphys.2017.12.008

- Nicolson SW, Human H (2013) Chemical composition of the 'low quality' pollen of sunflower
(*Helianthus annuus*, Asteraceae). *Apidologie* 44:144-152 doi:10.1007/s13592-012-0166-
5
- Nooten SS, Rehan SM (2020) Historical changes in bumble bee body size and range shift of
declining species. *Biodivers Conserv* 29:451-467 doi:10.1007/s10531-019-01893-7
- Otterstatter MC, Thomson JD (2006) Within-host dynamics of an intestinal pathogen of bumble
bees. *Parasitology* 133:749-761 doi:10.1017/s003118200600120x
- Palmer-Young EC, Farrell IW, Adler LS, Milano NJ, Egan PA, Irwin RE, Stevenson PC (2019a)
Secondary metabolites from nectar and pollen: a resource for ecological and evolutionary
studies. *Ecology* 100:e02621 doi:10.1002/ecy.2621
- Palmer-Young EC et al. (2019b) Chemistry of floral rewards: intra- and interspecific variability
of nectar and pollen secondary metabolites across taxa. *Ecol Monogr* 89:e01335
doi:10.1002/ecm.1335
- Potts SG et al. (2016) Safeguarding pollinators and their values to human well-being. *Nature*
540:220-229 doi:10.1038/nature20588
- R: A language and environment for statistical computing (2014) R Foundation for Statistical
Computing. Reference Source.
- Reagon M, Snow AA (2006) Cultivated *Helianthus annuus* (Asteraceae) volunteers as a genetic
"bridge" to weedy sunflower populations in North America. *Am J Bot* 93:127-133
doi:10.3732/ajb.93.1.127
- Richardson LL et al. (2015) Secondary metabolites in floral nectar reduce parasite infections in
bumblebees. *Proceedings of the Royal Society B-Biological Sciences* 282
doi:10.1098/rspb.2014.2471


- Robinson FA, Nation JL (1970) Long-chain fatty acids in honeybees in relation to sex, caste, and food during development. *J Apic Res* 9:121-127
- Roger N, Michez D, Wattiez R, Sheridan C, Vanderplanck M (2017) Diet effects on bumblebee health. *J Insect Physiol* 96:128-133 doi:10.1016/j.jinsphys.2016.11.002
- Roulston TH, Cane JH (2000) Pollen nutritional content and digestibility for animals. *Plant Syst Evol* 222:187-209
- Roulston TH, Cane JH, Buchmann SL (2000) What governs protein content of pollen: Pollinator preferences, pollen-pistil interactions, or phylogeny? *Ecol Monogr* 70:617-643
- Schmid-Hempel R et al. (2014) The invasion of southern South America by imported bumblebees and associated parasites. *J Anim Ecol* 83:823-837 doi:10.1111/1365-2656.12185
- Shykoff JA, Schmid-Hempel P (1991) Incidence and effects of four parasites in natural populations of bumble bees in Switzerland *Apidologie* 22:117-125 doi:10.1051/apido:19910204
- Spear DM, Silverman S, Forrest JRK (2016) Asteraceae pollen provisions protect *Osmia* mason bees (Hymenoptera: Megachilidae) from brood parasitism. *Am Nat* 187:797-803 doi:10.1086/686241
- Stevenson PC, Veitch NC (1996) Isoflavones from the roots of *Cicer judaicum*. *Phytochemistry* 43:695-700 doi:10.1016/0031-9422(96)00346-9
- Tasei JN, Aupinel P (2008) Nutritive value of 15 single pollens and pollen mixes tested on larvae produced by bumblebee workers (*Bombus terrestris*, Hymenoptera : Apidae). *Apidologie* 39:397-409 doi:10.1051/apido:2008017
- Therneau TM (2015) A package for survival analysis in S. version 2.38.


- 1
- 2
- 3
- 4 651 Therneau TM (2019) coxme: Mixed effects Cox models. R package version 2.2-14. .
- 5
- 6 652 Therneau TM, Grambsch PM (2000) Modeling survival data: Extending the Cox model.
- 7
- 8
- 9 653 Springer, New York
- 10
- 11 654 Thorburn LP, Adler LS, Irwin RE, Palmer-Young EC (2015) Variable effects of nicotine and
- 12
- 13
- 14 655 anabasine on parasitized bumble bees. F1000Research 4
- 15
- 16 656 Treanore ED, Vaudo AD, Grozinger CM, Fleischer SJ (2019) Examining the nutritional value
- 17
- 18
- 19 657 and effects of different floral resources in pumpkin agroecosystems on *Bombus impatiens*
- 20
- 21 658 worker physiology. Apidologie 50:542-552 doi:10.1007/s13592-019-00668-x
- 22
- 23
- 24 659 Ukiya M et al. (2003) Isolation, structural elucidation, and inhibitory effects of terpenoid and
- 25
- 26 660 lipid constituents from sunflower pollen on Epstein-Barr virus early antigen induced by
- 27
- 28
- 29 661 tumor promoter, TPA. J Agric Food Chem 51:2949-2957 doi:10.1021/jf0211231
- 30
- 31 662 Vanbergen AJ et al. (2013) Threats to an ecosystem service: pressures on pollinators. Front Ecol
- 32
- 33 663 Environ 11:251-259 doi:10.1890/120126
- 34
- 35
- 36 664 Vaudo AD, Patch HM, Mortensen DA, Tooker JF, Grozinger CM (2016) Macronutrient ratios in
- 37
- 38 665 pollen shape bumble bee (*Bombus impatiens*) foraging strategies and floral preferences.
- 39
- 40
- 41 666 Proc Natl Acad Sci USA 113:E4035-E4042 doi:10.1073/pnas.1606101113
- 42
- 43 667 Walters D, Meurer-Grimes B, Rovira I (2001) Antifungal activity of three spermidine
- 44
- 45
- 46 668 conjugates. FEMS Microbiol Lett 201:255-258 doi:10.1111/j.1574-6968.2001.tb10765.x
- 47
- 48 669 Wickham H (2016) ggplot2: Elegant graphics for data analysis. Springer-Verlag, New York
- 49
- 50
- 51 670 Yang K, Wu D, Ye XQ, Liu DH, Chen JC, Sun PL (2013) Characterization of chemical
- 52
- 53 671 composition of bee pollen in China. J Agric Food Chem 61:708-718
- 54
- 55 672 doi:10.1021/jf304056b
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

FIGURE LEGENDS

FIG. 1 Effect of (a) rutin and (b) triscoumaroyl spermidine in pollen on log₁₀-transformed *Crithidia* cell counts in *B. impatiens* workers. In both experiments, compounds were added to buckwheat pollen (*Fagopyrum esculentum*), which was used as a negative control, and compared to sunflower (*Helianthus annuus*) pollen as a positive control. “Low,” “Medium,” “High” and “Very High” refer to 0.001%, 0.01%, 0.1%, or 1% of compound added to pollen (wt/wt) prior to adding water. Different letters above means represent significantly different treatments ($P < 0.05$). Error bars indicate standard error

FIG. 2 Effect of (a, b) individual fatty acids and (c) combined fatty acids and sunflower oil in pollen on log₁₀-transformed *Crithidia* cell counts in *B. impatiens* workers. In (a) and (c), compounds were added to buckwheat pollen (*Fagopyrum esculentum*), which was used as a negative control, and compared to sunflower (*Helianthus annuus*) pollen as a positive control. In (b), compounds were instead added to mixed wildflower pollen, which was used as the control. In (c), “All 9 fatty acids” refers to a treatment in which the nine fatty acids tested individually in (a) and (b) were added in combination to a single pollen treatment. Different letters above means represent significantly different treatments ($P < 0.05$). Error bars indicate standard error







Click here to access/download
Supplementary Material
ESM_1_d2.docx





Click here to access/download
Supplementary Material
ESM_2.docx

