

# EXOTIC SPRINGER FIBERS FOR ORBITS CORRESPONDING TO ONE-ROW BIPARTITIONS

NEIL SAUNDERS AND ARIK WILBERT

ABSTRACT. We study the geometry and topology of exotic Springer fibers for orbits corresponding to one-row bipartitions from an explicit, combinatorial point of view. This includes a detailed analysis of the structure of the irreducible components and their intersections as well as the construction of an explicit affine paving. Moreover, we compute the ring structure of cohomology by constructing a CW-complex homotopy equivalent to the exotic Springer fiber. This homotopy equivalent space admits an action of the type C Weyl group inducing Kato's original exotic Springer representation on cohomology. Our results are described in terms of the diagrammatics of the two-boundary Temperley-Lieb algebra. This provides a first step in generalizing the geometric versions of Khovanov's arc algebra to the exotic setting.

## 1. INTRODUCTION

In [Kat09], Kato introduced the exotic nilpotent cone in a successful attempt to extend the Kazhdan-Lusztig-Ginzburg geometrization of affine Hecke algebras (see [KL87], [CG97]) from the one-parameter to the multi-parameter case by considering the equivariant algebraic  $K$ -theory of an exotic version of the Steinberg variety. Kato's construction establishes a Deligne-Langlands type classification of irreducible modules of affine Hecke algebras of type C with only very mild restrictions on the parameters. Let  $\mathcal{N}(\mathfrak{gl}_{2m}) \subseteq \mathfrak{gl}_{2m}(\mathbb{C})$  be the ordinary nilpotent cone of type A and let  $\mathcal{S} \subseteq \mathfrak{gl}_{2m}(\mathbb{C})$  denote the  $\mathrm{Sp}_{2m}(\mathbb{C})$ -invariant complement of  $\mathfrak{sp}_{2m}(\mathbb{C})$  in  $\mathfrak{gl}_{2m}(\mathbb{C})$ . The exotic nilpotent cone is the singular affine variety  $\mathfrak{N} = \mathbb{C}^{2m} \times (\mathcal{S} \cap \mathcal{N}(\mathfrak{gl}_{2m}))$ . Many of its properties relating to e.g. intersection cohomology of orbit closures (see [AH08] and [SS14]), theory of special pieces (see [AHS11]), and the Lusztig-Vogan bijection (see [Nan13]) have been explored in follow-up work to [Kat09].

In [Kat09], [Kat11], Kato used the exotic nilpotent cone to construct an exotic version of the Springer correspondence for the Weyl group of type C. This correspondence is less intricate than the classical type C Springer correspondence [Spr76], [Spr78], because it gives in fact a bijection between the orbits under the  $\mathrm{Sp}_{2m}(\mathbb{C})$ -action on  $\mathfrak{N}$  and the isomorphism classes of all complex, finite-dimensional, irreducible representations of the type C Weyl group (i.e. it behaves more like the classical type A Springer correspondence for the symmetric group). In particular, since the irreducible representations of the Weyl group can be labeled by bipartitions (see [Mac95, Appendix B]), it follows that the orbits in  $\mathfrak{N}$  can be labeled by bipartitions (see [AH08] for an explicit bijection between exotic nilpotent orbits and bipartitions).


In analogy to classical Springer theory, the irreducible Weyl group representations can be constructed on the top non-vanishing cohomology of so-called exotic Springer fibers. These algebraic varieties appear as the fibers under a resolution of singularities of the exotic nilpotent cone. More explicitly, given  $(v, x) \in \mathfrak{N}$ , the exotic Springer fiber  $\mathcal{F}l_{(v,x)}$  is the projective variety consisting of all flags  $\{0\} = F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_m \subsetneq \mathbb{C}^{2m}$  such that  $F_m$  is a Lagrangian subspace in  $\mathbb{C}^{2m}$  (with respect to a fixed symplectic form),  $v \in F_m$  and  $x F_i \subseteq F_{i-1}$ ,  $1 \leq i \leq m$ . The varieties  $\mathcal{F}l_{(v,x)}$  depend (up to isomorphism) only on the  $\mathrm{Sp}_{2m}(\mathbb{C})$ -orbit of  $(v, x)$  in  $\mathfrak{N}$  which justifies the notation  $\mathcal{F}l^{(\lambda,\mu)}$ , where  $(\lambda, \mu)$  is a bipartition of  $m$ . The geometric structure of the exotic Springer fibers is only poorly understood. In general, they are not smooth and decompose into many irreducible components. Note that even for the classical Springer fibers of type A the components are not smooth in general (see [FM10] for a parametrization of all smooth irreducible components in type A in terms of standard tableaux).

In this article we study exotic Springer fibers for orbits corresponding (via the bijection from [AH08]) to one-row bipartitions of  $m$ , i.e. pairs of partitions of the form  $((k), (m-k))$ ,  $0 \leq k \leq m$ , where each partition consists of a single number only. In this case we show that the irreducible components exhibit structural properties very similar to the ones discovered by Fung, [Fun03], for two-row Springer fibers of type A (see also [ES16] for type D). The first main result proved in Section 3 (see also Theorem 14(b)) is the following:

**Theorem A.** *Every irreducible component of  $\mathcal{F}l^{((k), (m-k))}$  is an  $m-k$ -fold iterated fiber bundle over  $\mathbb{P}^1$ . In particular, all irreducible components of  $\mathcal{F}l^{((k), (m-k))}$  are smooth.*

A combinatorial parametrization of the irreducible components of exotic Springer fibers in the spirit of [Spa76], [Var79], for type A (see also [Spa82], [vL89], for the remaining classical types) was only recently obtained in [NRS18] using standard bitableaux. Our results are based on a new parametrization in terms of so-called two-boundary cup diagrams. These diagrams enable us to write down explicit relations between the vector spaces of the flags contained in a given irreducible component (see Theorem 14(a) for details) similar to the case of type A two-row Springer fibers treated in [SW12, Section 1.2] based on the earlier work by Fung [Fun03].

Consider a rectangle in the plane with  $m$  vertices evenly spread along its upper horizontal edge. A two-boundary cup diagram is a diagram inside the rectangle obtained by either connecting two vertices by a lower semicircle called a cup, or by connecting a vertex with a point on the left (resp. right) vertical edge of the rectangle by a left-half-cup (resp. right-half-cup). In doing so, we require that the diagram is crossingless and every vertex is connected to exactly one endpoint of either a cup, a left-half-cup or a right-half-cup. If the cups, left-half-cups and right-half-cups of two given cup diagrams are incident with exactly the same vertices in both diagrams (regardless of the actual drawing of the cups, left-half-cups and right-half-cups in the plane) we consider the diagrams as equal. Here is an example:



In Section 5 we show that the set of all two-boundary cup diagrams on  $m$  vertices with a total number of at most  $m-k$  cups and right-half-cups combinatorially describes a Białynicki-Birula paving of  $\mathcal{F}l^{((k), (m-k))}$ . More precisely, each such diagram corresponds to an affine cell and the respective diagram can be used to explicitly determine all flags contained in a given cell (see Theorem 34). The closures of the cells corresponding to the diagrams with the number of cups and right-half-cups being maximal are exactly the irreducible components of  $\mathcal{F}l^{((k), (m-k))}$ . In particular, we get a bijection between the set  $\mathbb{B}^{((k), (m-k))}$  of all two-boundary cup diagrams on  $m$  vertices with a total number of  $m-k$  cups and right-half-cups and the irreducible components of  $\mathcal{F}l^{((k), (m-k))}$  (see also Theorem 14(c)).

Two-boundary cup diagrams appear naturally in the representation theory of two-boundary Temperley-Lieb algebras, [dGN09]. These algebras admit a neat diagrammatic description and can be realized as quotients of multi-parameter affine Hecke algebras of type C, see [Ern12], [Ern18]. In light of [Kat09], the latter fact explains the appearance of the two-boundary cup diagrams when studying exotic Springer fibers. In contrast to the classical Temperley-Lieb algebras for finite types, the two-boundary Temperley-Lieb algebra is infinite-dimensional, which makes its representation theory particularly interesting. The two-boundary cup diagrams naturally label a basis of the so-called spin chain representations which are finite-dimensional irreducible representations of maximal dimension. They play a prominent role in statistical mechanics when studying exactly solvable lattice models with boundary conditions, [dGP04], [dGNPR05], [Nic06a], [Nic06b].

In Section 4 we use the two-boundary cup diagrams to construct a topological model of the exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$  similar to the topological models of two-row Springer fibers constructed in [Kho04], [Weh09], [RT11], [Rus11], for type A and in [ES16], [Wil18], for the remaining classical two-row types. Let  $\mathbb{S}^2 \subseteq \mathbb{R}^3$  be the two-dimensional standard unit sphere with north pole  $p = (0, 0, 1)$ . Given a cup diagram  $\mathbf{a} \in \mathbb{B}^{((k), (m-k))}$ , define  $S_{\mathbf{a}} \subseteq (\mathbb{S}^2)^m$  as the submanifold consisting of all  $(x_1, \dots, x_m) \in (\mathbb{S}^2)^m$  satisfying the relations  $x_j = -x_i$

if vertices  $i$  and  $j$  are connected by a cup in  $\mathbf{a}$ , and  $x_i = p$  if vertex  $i$  is connected to a left-half-cup in  $\mathbf{a}$ . There are no relations involving coordinates indexed by a vertex connected to a right-half-cup. This yields a topological model for the exotic Springer fiber, i.e. we obtain the following homeomorphism (see also Theorem 23):

**Theorem B.** *There exists a homeomorphism*

$$\bigcup_{\mathbf{a} \in \mathbb{B}^{((k), (m-k))}} S_{\mathbf{a}} \cong \mathcal{F}l^{((k), (m-k))}$$

such that the images of the  $S_{\mathbf{a}}$  are the irreducible components of  $\mathcal{F}l^{((k), (m-k))}$ .

We equip  $\mathbb{S}^2$  with the structure of a CW-complex consisting of a 0-cell and a 2-cell and equip  $(\mathbb{S}^2)^m$  with the product CW-structure. Let  $\text{Sk}_{2(m-k)}^m$  denote the  $2(m-k)$ -skeleton of  $(\mathbb{S}^2)^m$ . The next theorem explicitly identifies the homotopy type of the exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$ , i.e. we show that  $\text{Sk}_{2(m-k)}^m$  is isomorphic (but in general not homeomorphic, see Example 55) to  $\mathcal{F}l^{((k), (m-k))}$  in the homotopy category of topological spaces.

**Theorem C.** *There exists a homotopy equivalence  $\mathcal{F}l^{((k), (m-k))} \simeq \text{Sk}_{2(m-k)}^m$ . In particular, the cohomology ring  $H^*(\mathcal{F}l^{((k), (m-k))}, \mathbb{C})$  is isomorphic (as a graded algebra) to  $H^*(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  which has an explicit presentation given by*

$$\mathbb{C}[X_1, \dots, X_m] \left/ \left\langle X_i^2, X_I \mid \begin{array}{l} 1 \leq i \leq m, \\ I \subseteq \{1, \dots, m\}, |I| = m - k + 1 \end{array} \right. \right\rangle,$$

where  $X_I = \prod_{i \in I} X_i$  and  $\deg(X_i) = 2$ .

Even in the case of one-row bipartitions, the type C Weyl group does not act on the exotic Springer fiber (and neither on its topological model from Theorem B). However, it acts on the homotopy equivalent space  $\text{Sk}_{2(m-k)}^m$  in a natural way and induces an action on cohomology which can be described explicitly (see Proposition 57). We prove that the resulting representation  $H^{2l}(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  is isomorphic to the irreducible Specht module of the type C Weyl group labeled by  $((l), (m-l))$ ,  $0 \leq l \leq m-k$ . In particular, the top degree representation is as expected from the exotic Springer correspondence [Kat11]. In fact, by using results from [Kat17], it follows that our representations coincide with the ones constructed by Kato in all cohomological degrees. Furthermore, our explicit presentation of  $H^*(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  comes with a distinguished basis of  $H^{2l}(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  given by monomials with  $l$  pairwise different factors. We show that this monomial basis corresponds to the well known bipolytabloid basis indexed by standard bitableaux of the Specht module, see [Can96]. It would be interesting to extend the construction of homotopy types which admit an action of the Weyl group to more general Springer fibers.

Finally, we describe the structure of the pairwise intersections of the irreducible components of the exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$  using circle diagrams (see Theorem 28). A circle diagram is obtained by putting the cup diagram assigned to a component upside down on top of the diagram associated to another component. We can then consider the vector space  $\mathbb{K}_m$  whose basis is given by all such oriented circle diagrams and obtain the following result.

**Theorem D.** *There is an isomorphism of vector spaces*

$$\mathbb{K}_m \cong \bigoplus_{(\mathbf{a}, \mathbf{b}) \in (\mathbb{B}^{((k), (m-k))})^2} H^*(K_{\mathbf{a}} \cap K_{\mathbf{b}}, \mathbb{C}). \quad (1)$$

The right-hand side of (1) can be equipped with a convolution product by mimicking the construction in [SW12]. This product should have a diagrammatic description in terms of the two-boundary cup diagram combinatorics from the left-hand side of (1). For two-row Springer fibers of type A such a diagrammatic description was obtained in [SW12]. This gives a geometric construction of the (generalized) arc algebras of type A (including their quasi-hereditary covers) which were combinatorially defined in [Str09],[BS11a] (see also [CK14]) based on [Kho02]. These algebras do not only appear in low-dimensional topology in the

context of defining tangle homology theories, but they are known to be related to many interesting representation theoretic categories such as parabolic category  $\mathcal{O}$  [BS11b], perverse sheaves on Grassmannians [Str09], non-semisimple representation theory of the Brauer algebra, [BS12a], and the general linear Lie superalgebra, [BS12b]. Our article can be seen as providing the geometric framework for constructing similar convolution algebras for exotic Springer fibers using two-boundary cup diagrams. The resulting exotic arc algebras should play a role in categorifying Schur-Weyl dualities involving multiparameter affine Hecke algebras of type C studied recently (on the decategorified level) in [DR18] and [FLL<sup>+</sup>18]. Establishing relationships between interesting representation theoretic categories comparable to the aforementioned results in type A is expected to be nontrivial in the exotic case. In particular, since the resolution of the exotic nilpotent cone is not symplectic but only Calabi-Yau, the involved categories would necessarily need to have very different properties from the ones arising from the type A convolution algebras.

**Acknowledgments.** The authors would like to thank Nora Ganter, Anthony Henderson, Vinoth Nandakumar, Arun Ram and Catharina Stroppel for useful discussions and comments. We are particularly grateful to Maud De Visscher for inviting the second author to visit City, University of London, where this project began.

## 2. BASIC DEFINITIONS

**2.1. Exotic nilpotent cone and exotic Springer fibers.** Let  $V$  be a  $2m$ -dimensional complex vector space with basis  $e_1, \dots, e_m, f_1, \dots, f_m$  and symplectic form  $\omega$  given by  $\omega(e_i, f_j) = \delta_{i+j, m+1} = -\omega(f_j, e_i)$  and zero otherwise. Let  $\mathrm{Sp}(V, \omega)$  be the symplectic group of all linear automorphisms of  $V$  preserving  $\omega$  and let  $\mathfrak{sp}(V, \omega)$  be its Lie algebra, i.e. the Lie subalgebra of the general linear Lie algebra  $\mathfrak{gl}(V)$  consisting of all linear endomorphisms  $x$  of  $V$  such that  $\omega(xv, w) = -\omega(v, xw)$  for all  $v, w \in V$ . The conjugation action of  $\mathrm{Sp}(V, \omega)$  on  $\mathfrak{gl}(V)$  yields a direct sum decomposition  $\mathfrak{gl}(V) = \mathfrak{sp}(V, \omega) \oplus \mathcal{S}(V, \omega)$  of  $\mathrm{Sp}(V, \omega)$ -modules, where

$$\mathcal{S}(V, \omega) = \{x \in \mathfrak{gl}(V) \mid \omega(xv, w) - \omega(v, xw) = 0, \forall v, w \in V\}.$$

**Definition 1.** The *exotic nilpotent cone* is the affine variety defined as

$$\mathfrak{N} = V \times (\mathcal{S}(V, \omega) \cap \mathcal{N}(\mathfrak{gl}(V))),$$

where  $\mathcal{N}(\mathfrak{gl}(V)) \subseteq \mathfrak{gl}(V)$  is the ordinary nilpotent cone consisting of all nilpotent (in the usual sense of linear algebra) linear endomorphisms of  $V$ .

Given a partition  $\lambda$ , we write  $|\lambda|$  to denote the sum  $\sum_{i \geq 1} \lambda_i$  of its parts and  $\ell(\lambda)$  denotes the number of parts. Moreover,  $\lambda + \mu$  is the partition defined by  $(\lambda + \mu)_i = \lambda_i + \mu_i$  and  $\lambda \cup \mu$  is the partition obtained by taking both the parts of  $\lambda$  and  $\mu$  and order them such that  $\lambda \cup \mu$  is a partition. A *bipartition* of  $m$  is a pair  $(\lambda, \mu)$  of partitions such that  $|\lambda| + |\mu| = m$ .

**Proposition 2** ([AH08, Theorem 6.1]). *The orbits under the  $\mathrm{Sp}(V, \omega)$ -action on  $\mathfrak{N}$  are in bijective correspondence with bipartitions of  $m$ .*

*Remark 3.* It follows from Section 2 and Section 6 of [AH08] (see also [NRS17, Theorem 2.10]) that a point  $(v, x) \in \mathfrak{N}$  is contained in the orbit labeled by the bipartition  $(\lambda, \mu)$  of  $m$  if and only if there is a basis of  $V$  given by

$$\{e_{ij}, f_{ij} \mid 1 \leq i \leq \ell(\lambda + \mu), 1 \leq j \leq \lambda_i + \mu_i\},$$

where  $\omega(e_{ij}, f_{i'j'}) = \delta_{i,i'} \delta_{j+j', \lambda_i + \mu_i + 1}$ ,  $v = \sum_{i=1}^{\ell(\lambda)} e_{i, \lambda_i}$  and such that the action of  $x$  on this basis is as follows:

$$xe_{ij} = \begin{cases} e_{i, j-1} & \text{if } j \geq 2 \\ 0 & \text{if } j = 1 \end{cases} \quad xf_{ij} = \begin{cases} f_{i, j-1} & \text{if } j \geq 2 \\ 0 & \text{if } j = 1. \end{cases}$$

Note that  $x$  has Jordan type  $(\lambda + \mu) \cup (\lambda + \mu)$ .

By Remark 3 we directly obtain the following distinguished representatives of orbits corresponding to one-row bipartitions.

**Lemma 4.** *Let  $x \in \mathcal{N}(\mathfrak{gl}(V))$  be the nilpotent endomorphism of  $V$  whose action on  $V$  is given as follows:*

$$e_1 \leftarrow e_2 \leftarrow \dots \leftarrow e_m \quad f_1 \leftarrow f_2 \leftarrow \dots \leftarrow f_m. \quad (2)$$

The vectors  $e_1$  and  $f_1$  are sent to 0. Then the point  $(e_k, x) \in \mathfrak{N}$ , where  $e_k \in V$  is the  $k$ th Jordan basis vector of the first block in (2), is contained in the orbit labeled by the bipartition  $((k), (m-k))$ .

Let  $\mathcal{F}l_C$  be the flag variety of type  $C$  consisting of all flags  $\{0\} \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_m \subsetneq V$  such that  $\dim_{\mathbb{C}}(F_i) = i$  and  $F_i \subseteq V$  is isotropic with respect to  $\omega$ ,  $1 \leq i \leq m$ . The nilpotent cone  $\mathfrak{N}$  has a resolution of singularities  $\pi: \tilde{\mathfrak{N}} \rightarrow \mathfrak{N}$ , where

$$\tilde{\mathfrak{N}} = \{((F_1, \dots, F_m), (v, x)) \in \mathcal{F}l_C \times \mathfrak{N} \mid zF_i \subseteq F_{i-1}, v \in F_m\}$$

and  $\pi$  is the projection onto the first component, [Kat11].

**Definition 5.** Given  $(v, x) \in \mathfrak{N}$ , the *exotic Springer fiber*  $\mathcal{F}l_{(v,x)}$  is the variety consisting of all flags  $\mathcal{F}l_C$  satisfying  $v \in F_m$  and  $xF_i \subseteq F_{i-1}$ .

Analogous to the ordinary Springer fibers, the exotic Springer fibers naturally appear as the fibers of a resolution of the exotic nilpotent cone. Up to isomorphism the exotic Springer fiber only depends on the  $\mathrm{Sp}(V, \omega)$ -orbit of  $(v, x)$  in  $\mathfrak{N}$  and not on the chosen point itself. This allows us to talk about the  $(\lambda, \mu)$ -Springer fiber denoted by  $\mathcal{F}l^{(\lambda, \mu)}$  without ambiguity.

**2.2. Classification of irreducible components.** Let  $(\lambda, \mu)$  be a bipartition of  $m$ . A bitableau of shape  $(\lambda, \mu)$  is a filling of the  $m$  boxes of the underlying pair of Young diagrams with the numbers  $1, 2, \dots, m$  such that each number appears exactly once. A bitableau is called standard if the entries increase (to the right and downwards) along rows and columns. Note that this convention differs from the one used in [NRS18, NRS17].

**Proposition 6** ([NRS18, Theorem 2.12]). *There is a bijection between the irreducible components of the  $(\lambda, \mu)$  exotic Springer fiber and the set of all standard bitableaux of shape  $(\lambda, \mu)$ . Moreover, the dimension of the irreducible components equals  $2M(\lambda + \mu) + |\mu|$ , where  $M(\lambda + \mu) = \sum_{i \geq 1} (i-1)(\lambda_i + \mu_i)$  is the dimension of the Springer fiber of type  $A$  corresponding to  $\lambda + \mu$ .*


**2.3. Two-boundary cup diagrams.** Throughout this article we fix a one-row bipartition  $((k), (m-k))$  of the positive integer  $m$ ,  $0 \leq k \leq m$ .

Consider a rectangle in the plane with  $m$  vertices evenly spread along the upper horizontal edge. The vertices are labeled by the consecutive integers  $1, \dots, m$  in increasing order from left to right.

**Definition 7.** A *two-boundary cup diagram* (or shorter *cup diagram*) is obtained by either connecting two vertices by a lower semicircle called a *cup*, or by connecting a vertex with a point on the left (resp. right) vertical edge of the rectangle by a *left-half-cup* (resp. *right-half-cup*). In doing so we require that the diagram is crossingless and every vertex is connected with exactly one endpoint of a cup, left-half-cup or right-half-cup. If the cups, left-half-cups and right-half-cups of two given cup diagrams are incident with exactly the same vertices in both diagrams (regardless of the precise drawing of the cups, left-half-cups and right-half-cups) we consider the diagrams as equal. The notation  $i-j$  means that  $i < j$  are connected by a cup and  $|-i$  means that  $i$  is connected to a left-half-cup.

We write  $\mathbb{B}^{((k), (m-k))}$  to denote the set of all cup diagrams on  $m$  vertices such that the number of cups plus the number of right-half-cups equals  $m-k$ .

*Example 8.* The set  $\mathbb{B}^{((3),(1))}$  consists of the cup diagrams




In the following we will often omit to indicate the vertex labeling on the top horizontal edge of the rectangle.

**Lemma 9.** *There exists a bijection between the set of standard 1-row bitableaux of shape  $((k), (m-k))$  and the set of cup diagrams  $\mathbb{B}^{((k), (m-k))}$ . The map is given by sending a bitableau to the cup diagram whose left endpoints of cups or endpoints of right-half-cups are connected to the vertices whose labels are contained in the second Young tableau.*

*Proof.* Such a cup diagram exists and is unique (thus we have a well-defined map). Obvious inverse is then the following: Given a cup diagram, put the left endpoints of cups and arc segments in the right Young tableaux of the bipartition  $((k), (m-k))$  (in the unique order such that the bitableaux will be standard).  $\square$

*Example 10.* We illustrate this bijection of cup diagrams with bitableaux for the bipartition  $((2), (2))$ . Explicitly the bijection is:



*Remark 11.* By Lemma 9 and Proposition 6 the cup diagrams in  $\mathbb{B}^{((k), (m-k))}$  parametrize the irreducible components of the exotic Springer fiber corresponding to a pair of one-row partitions  $((k), (m-k))$ . By Proposition 6 this exotic Springer fiber has dimension  $m-k$  because  $M((k) + (m-k))$  is zero as the type  $A$  Springer fiber corresponding to the one-row partition  $(k) + (m-k) = (m)$  is just a point. In particular, the dimension of this exotic Springer fiber can be read off the cup diagrams by counting the number of cups plus right-half-cups.

### 3. STRUCTURE OF IRREDUCIBLE COMPONENTS

**3.1. Embedding the exotic Springer fiber into a smooth variety.** Let  $N > 0$  be a large integer (see Remark 12 for a more precise explanation of what is meant by “large”) and let  $z: \mathbb{C}^{2N} \rightarrow \mathbb{C}^{2N}$  be a nilpotent linear endomorphism with two equally sized Jordan blocks, i.e. there is a Jordan basis

$$e_1 \leftarrow e_2 \leftarrow \dots \leftarrow e_N \quad f_1 \leftarrow f_2 \leftarrow \dots \leftarrow f_N \quad (3)$$

on which  $z$  acts as indicated (the vectors  $e_1$  and  $f_1$  are sent to zero). We equip  $\mathbb{C}^{2N}$  with a Hermitian form by defining  $e_1, \dots, e_N, f_1, \dots, f_N$  to be an orthonormal basis.

In the following we will consider the smooth, projective variety

$$Y_m := \{(F_1, \dots, F_m) \mid F_i \subseteq \mathbb{C}^{2N}, \dim F_i = i, F_1 \subseteq \dots \subseteq F_m, zF_i \subseteq F_{i-1}\} \quad (4)$$

defined in [CK08, §2].

*Remark 12.* Note that the conditions  $zF_i \subseteq F_{i-1}$  yield

$$F_m \subseteq z^{-1}F_{m-1} \subseteq \dots \subseteq z^{-m}(0) = \text{span}(e_1, \dots, e_m, f_1, \dots, f_m).$$

Thus, the variety  $Y_m$  is independent of the choice of  $N$  as long as  $N \geq m$ . In particular, we can always assume (by increasing  $N$  if necessary) that all the subspaces of a flag in  $Y_m$  are contained in the image of  $z$ .

Let  $E_m \subseteq \mathbb{C}^{2N}$  be the subspace spanned by  $e_1, \dots, e_m, f_1, \dots, f_m$ . We equip  $E_m$  with a symplectic form  $\omega_m$  defined as follows: for all  $j, j' \in \{1, \dots, m\}$  we set

$$\omega_m(e_j, f_{j'}) = \delta_{j+j', m+1} = -\omega_m(f_{j'}, e_j).$$

Otherwise the pairing is defined to be zero. A direct computation shows that  $\omega_m(z(v), w) = \omega_m(v, z(w))$  for all  $v, w \in E_m$ , i.e. the restriction  $z_m$  of  $z$  to the subspace  $E_m$  is a nilpotent endomorphism which is self-adjoint with respect to  $\omega_m$ .

Thus, we can evidently view the exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$  as a subvariety of  $Y_m$  via the following identification

$$\mathcal{F}l^{((k), (m-k))} \cong \left\{ (F_1, \dots, F_m) \in Y_m \mid \begin{array}{l} e_k \in F_m \subseteq E_m \text{ and } F_m \text{ is} \\ \text{isotropic with respect to } \omega_m \end{array} \right\}. \quad (5)$$

The next proposition shows that the description of the embedding (5) can be further simplified and in our case we can in fact ignore the symplectic structure.

**Proposition 13.** *The exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$  can be viewed as a subvariety of  $Y_m$  via the following identification*

$$\mathcal{F}l^{((k), (m-k))} \cong \{(F_1, \dots, F_m) \in Y_m \mid e_k \in F_m\}. \quad (6)$$

*Proof.* Note that the inclusion  $F_m \subseteq E_m$  is automatically satisfied by Remark 12. In order to reduce the description (5) to (6), it thus suffices to show that the isotropy condition on  $F_m$  is automatically satisfied for all  $(F_1, \dots, F_m) \in Y_m$  and hence can be dropped, too.

Let  $(F_1, \dots, F_m) \in Y_m$  be a flag. By induction we assume that we have already shown that the vector spaces  $F_1, \dots, F_i$  are isotropic with respect to  $\omega_m$ . Furthermore, we assume that there exist linearly independent vectors

$$\begin{array}{cccccc} e_1^{(i)}, & e_2^{(i)}, & \dots, & e_{m-i-1}^{(i)}, & e_{m-i}^{(i)}, \\ f_1^{(i)}, & f_2^{(i)}, & \dots, & f_{m-i-1}^{(i)}, & f_{m-i}^{(i)}, \end{array} \quad (7)$$

in  $\mathbb{C}^{2N}$  such that the following three properties hold:

- (P1) The endomorphism  $z$  maps each vector in (7) to its left neighbor and the two leftmost vectors in each row to  $F_i$ .
- (P2) We have a decomposition  $F_i \oplus G_i = F_i^\perp$ , where  $G_i \subseteq \mathbb{C}^{2N}$  is the subspace spanned by the vectors in (7) and the orthogonal complement is taken with respect to  $\omega_m$ .
- (P3) We have  $\omega_m(e_j^{(i)}, f_{j'}^{(i)}) = \delta_{j+j', m-i+1}$ .

We remark that the induction starts with the zero space  $\{0\}$  and the basis  $e_1, \dots, e_m, f_1, \dots, f_m$  of  $E_m$  which obviously satisfies (P1)-(P3).

Since  $zF_{i+1} \subseteq F_i$ , we can write

$$F_{i+1} = F_i \oplus \text{span}(\alpha e_1^{(i)} + \beta f_1^{(i)})$$

for some  $\alpha, \beta \in \mathbb{C}$ ,  $\alpha \neq 0$  or  $\beta \neq 0$ . Then properties (P2) and (P3) show that  $F_{i+1}$  is indeed isotropic.

In order to complete the inductive argument we need to show the existence of the vectors (7) for the space  $F_{i+1}$ . We distinguish between two different cases depending on the parameters  $\alpha$  and  $\beta$ .

*Case 1:* If  $\alpha \neq 0$  and  $\beta \neq 0$  we consider the linearly independent vectors

$$\begin{array}{l} \alpha e_1^{(i)} - \beta f_1^{(i)}, \alpha e_2^{(i)} - \beta f_2^{(i)}, \dots, \alpha e_{m-i-1}^{(i)} - \beta f_{m-i-1}^{(i)} \\ \alpha e_2^{(i)} + \beta f_2^{(i)}, \alpha e_3^{(i)} + \beta f_3^{(i)}, \dots, \alpha e_{m-i}^{(i)} + \beta f_{m-i}^{(i)} \end{array} \quad (8)$$

and define

$$e_j^{(i+1)} := \frac{1}{\sqrt{2\alpha\beta}} \left( \alpha e_j^{(i)} - \beta f_j^{(i)} \right) \quad \text{and} \quad f_j^{(i+1)} := \frac{1}{\sqrt{2\alpha\beta}} \left( \alpha e_{j+2}^{(i)} + \beta f_{j+2}^{(i)} \right)$$

for  $j \in \{1, \dots, m-i-1\}$ . It is straightforward to check that this collection of vectors satisfies (P1)-(P3) for  $F_{i+1}$ .

*Case 2:* If  $\beta = 0$  we define

$$e_j^{(i+1)} := e_{j+1}^{(i)} \quad \text{and} \quad f_j^{(i+1)} := f_j^{(i)}$$

for  $j \in \{1, \dots, m-i-1\}$  and if  $\alpha = 0$  we set

$$e_j^{(i+1)} := e_j^{(i)} \quad \text{and} \quad f_j^{(i+1)} := f_{j+1}^{(i)}$$

for  $j \in \{1, \dots, m-i-1\}$ . Again, one easily checks that this collection of vectors satisfies (P1)-(P3) for  $F_{i+1}$ .  $\square$

For the remainder of this article we will write  $\mathcal{F}l^{((k),(m-k))}$  to denote the embedded exotic Springer fiber via identification (6) from Proposition 13.

**Theorem 14.** *The following statements hold:*

- (a) Given  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ , define  $K_{\mathbf{a}} \subseteq Y_m$  as the subvariety consisting of all flags  $(F_1, \dots, F_m) \in Y_m$  which satisfy the following conditions imposed by the cup diagram  $\mathbf{a}$ :

- (i) If  $i-j$ , then

$$F_j = z^{-\frac{1}{2}(j-i+1)} F_{i-1}.$$

- (ii) If  $\vdash i$ , then

$$F_i = F_{i-1} + \text{span} \left( e_{\frac{1}{2}(i+\rho_{\mathbf{a}}(i))} \right),$$

where  $\rho_{\mathbf{a}}(i)$  counts the number of left-half-cups to the left of vertex  $i$  (including the vertex  $i$ ) in  $\mathbf{a}$ .

There are no relations for vector spaces indexed by a vertex connected to a right-half-cup. The variety  $K_{\mathbf{a}} \subseteq Y_m$  is an irreducible component of the exotic Springer fiber  $\mathcal{F}l^{((k),(m-k))} \subseteq Y_m$ .

- (b) The irreducible component  $K_{\mathbf{a}} \subseteq \mathcal{F}l^{((k),(m-k))}$  is an  $(m-k)$ -fold iterated fiber bundle over  $\mathbb{P}^1$ , i.e. there exist spaces  $K_{\mathbf{a}} = X_1, X_2, \dots, X_{m-k}, X_{m-k+1} = \text{pt}$  together with maps  $p_1, p_2, \dots, p_{m-k}$  such that  $p_j: X_j \rightarrow \mathbb{P}^1$  is a fiber bundle with typical fiber  $X_{j+1}$ . In particular,  $K_{\mathbf{a}}$  is a smooth variety.
- (c) The map  $\mathbf{a} \mapsto K_{\mathbf{a}}$  defines a bijection between the two-boundary cup diagrams in  $\mathbb{B}^{((k),(m-k))}$  and the irreducible components of  $\mathcal{F}l^{((k),(m-k))}$ .

*Remark 15.* Given a flag  $(F_1, \dots, F_m) \in Y_m$ , we have  $z^{-1}F_i = z_m^{-1}F_i$ , where  $z_m$  is the restriction of  $z$  to  $E_m$ , because  $z^{-1}F_i$  is contained in  $E_m$  for all  $i \in \{1, \dots, m\}$  (see also Remark 12). In particular, we could replace  $z$  by the restriction  $z_m$  of  $z$  to  $E_m$  in Theorem 14. Hence, the description of the irreducible components of the exotic Springer fiber in Theorem 14(a) also makes sense without the embedding into  $Y_m$ .

In order to prove Theorem 14 we consider the smooth subvariety  $X_m^i \subseteq Y_m$ ,  $i \in \{1, \dots, m-1\}$ , defined by

$$X_m^i := \{(F_1, \dots, F_m) \in Y_m \mid F_{i+1} = z^{-1}F_{i-1}\}, \quad (9)$$

and a surjective morphism of varieties  $q_m^i: X_m^i \rightarrow Y_{m-2}$  given by

$$(F_1, \dots, F_m) \mapsto (F_1, \dots, F_{i-1}, zF_{i+2}, \dots, zF_m), \quad (10)$$

see also [CK08, §2].

**Lemma 16.** *Let  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$  be a cup diagram with a cup connecting vertices  $i$  and  $i+1$  and let  $\mathbf{a}' \in \mathbb{B}^{((k-1),(m-k-1))}$  be the cup diagram obtained by deleting this cup. Then we have  $K_{\mathbf{a}} = (q_m^i)^{-1}(K_{\mathbf{a}'})$ .*

*Proof.* We have to show that a flag  $(F_1, \dots, F_m) \in Y_m$  satisfies the conditions (i), (ii) from Theorem 14(a) with respect to the cup diagram  $\mathbf{a}$  if and only if

$$q_m^i(F_1, \dots, F_m) = (F_1, \dots, F_{i-1}, zF_{i+2}, \dots, zF_m) \in Y_{m-2}$$

satisfies these conditions with respect to  $\mathbf{a}'$ .

In order to deal with condition (ii) suppose there is a left-half-cup connected to vertex  $j$  in  $\mathbf{a}$ . If  $j \leq i-1$  there is nothing to show. If  $j > i+1$  we need to show that

$$F_j = F_{j-1} + \text{span}\left(e_{\frac{1}{2}(j-\alpha)}\right) \quad \Leftrightarrow \quad zF_j = zF_{j-1} + \text{span}\left(e_{\frac{1}{2}(j-2-\alpha)}\right).$$


But this follows by applying  $z$  (resp.  $z^{-1}$ ) to the left (resp. right) side of the equivalence.

For the proof involving condition (i) we refer to the proof of [Weh09, Lemma 3.2] for the precise argument.  $\square$

**Proposition 17.** *We have  $K_{\mathbf{a}} \subseteq \mathcal{F}l^{((k),(m-k))}$  for all  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ .*

*Proof.* We proceed by induction on the number of cups in  $\mathbf{a}$ . By Proposition 13 it suffices to show that  $e_k \in F_m$  for all  $(F_1, \dots, F_m) \in K_{\mathbf{a}}$ .

*Induction Start.* Let  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$  be the cup diagram without any cups contained in  $\mathbb{B}^{((k),(m-k))}$ , i.e. the one given by  $k$  subsequent left-half-cups followed by  $m-k$  right-half-cups



Given a flag  $(F_1, \dots, F_m) \in K_{\mathbf{a}}$  it follows from the relations in Theorem 14(a) that  $F_k = \text{span}(e_1, \dots, e_k)$ . In particular, we have  $e_k \in F_k \subseteq F_m$  which implies  $(F_1, \dots, F_m) \in \mathcal{F}l^{((k),(m-k))}$ .

*Inductive Step.* Assume there exists a cup in  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$  and fix a cup connecting neighboring vertices  $i$  and  $i+1$ . Let  $\mathbf{a}' \in \mathbb{B}^{((k-1),(m-k-1))}$  denote the cup diagram obtained by removing this cup.

Given an arbitrary flag  $(F_1, \dots, F_m) \in K_{\mathbf{a}}$ , we have

$$q_m^i(F_1, \dots, F_m) = (F_1, \dots, F_{i-1}, zF_{i+2}, \dots, zF_m) \in K_{\mathbf{a}'}$$

by Lemma 16. By induction we assume that  $K_{\mathbf{a}'} \subseteq \mathcal{F}l^{((k-1),(m-k-1))}$ . In particular, we have  $e_{k-1} \in zF_m$ , i.e.  $e_{k-1} = z(v)$  for some  $v \in F_m$ . Any such  $v \in F_m$  must be of the form  $v = e_k + \alpha e_1 + \beta f_1$  for  $\alpha, \beta \in \mathbb{C}$ . Since  $\text{span}(e_1, f_1) = \ker(z) = z^{-1}(0) \subseteq z^{-1}F_{i-1} = F_{i+1} \subseteq F_m$  we see that  $v \in F_m$  implies  $e_k \in F_m$ . But  $e_k \in F_m$  means  $(F_1, \dots, F_m) \in \mathcal{F}l^{((k),(m-k))}$ .  $\square$

*Proof (Theorem 14).* We first show that  $K_{\mathbf{a}}$  is an  $m-k$ -fold iterated  $\mathbb{P}^1$ -bundle. This follows as in [Fun03, Proposition 5.1] (see also [Sch12, Section 8]). We briefly recall the argument in our setup. Let  $i_1 < i_2 < \dots < i_{m-k}$  denote the vertices connected to either a left endpoint of a cup or a right-half-cup in  $\mathbf{a}$ . Note that the space  $F_{i_1-1}$  is the same for every flag  $(F_1, \dots, F_m) \in K_{\mathbf{a}}$  because each vertex to the left of  $i_1$  is connected to a left-half-cup and successively applying relation (ii) in Theorem 14 (a) determines this space uniquely. Hence, we can define the fiber bundle

$$p_1: K_{\mathbf{a}} \rightarrow \mathbb{P}(z^{-1}F_{i_1-1}/F_{i_1-1}) \cong \mathbb{P}^1, \quad (F_1, \dots, F_m) \mapsto F_{i_1}/F_{i_1-1}$$

which has typical fiber  $X_2$  consisting of all flags in  $(F_1, \dots, F_m) \in K_{\mathbf{a}}$  with  $F_{i_1}$  (and  $F_j$ , if  $(i_1+1)$  and  $j$  are connected by a cup) fixed. Now we repeat the above construction replacing  $X_1$  by  $X_2$  and the vertex  $i_1$  by  $i_2$  and continue until we have exhausted all of the vertices  $i_1 < \dots < i_{m-k}$ .

In order to prove parts (a) and (b) of Theorem 14 we first note that  $K_{\mathbf{a}}$  is smooth since it is an  $m-k$ -fold iterated fiber bundle over  $\mathbb{P}^1$ . Furthermore,  $K_{\mathbf{a}}$  is connected (in Proposition 22 we show that  $K_{\mathbf{a}}$  is the preimage under a diffeomorphism of a topological space  $T_{\mathbf{a}}$  which will evidently be connected). This shows that  $K_{\mathbf{a}}$  is irreducible. By Proposition 17 the variety  $K_{\mathbf{a}}$  is contained in  $\mathcal{F}l^{((k),(m-k))}$ . Finally, the dimension of  $K_{\mathbf{a}}$  equals  $m-k$  which is the dimension of  $\mathcal{F}l^{((k),(m-k))}$  by [NRS18, Theorem 2.12]. Hence,  $K_{\mathbf{a}}$  is an irreducible component of the (embedded) exotic Springer fiber  $\mathcal{F}l^{((k),(m-k))}$ . In particular, parts (a) and (b) of Theorem 14 are now clear from the above.

By Proposition 6 and Lemma 9 we know that the cup diagrams in  $\mathbb{B}^{((k),(m-k))}$  are in bijective correspondence with the irreducible components of  $\mathcal{F}l^{((k),(m-k))}$ . Since the irreducible components  $K_{\mathbf{a}}$  are evidently different for different  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ , we see that the map  $\mathbf{a} \mapsto K_{\mathbf{a}}$  explicitly realizes this bijection which proves part (c) of the theorem.  $\square$

*Example 18.* For the cup diagrams from Example 8, Theorem 14(a) yields the following sets of flags

- $K_{\mathbf{a}} = \{F_1 \subseteq \text{span}(e_1, f_1) \subseteq \text{span}(e_1, e_2, f_1) \subseteq \text{span}(e_1, e_2, e_3, f_1)\}$ ,
- $K_{\mathbf{b}} = \{\text{span}(e_1) \subseteq F_2 \subseteq \text{span}(e_1, e_2, f_1) \subseteq \text{span}(e_1, e_2, e_3, f_1)\}$ ,
- $K_{\mathbf{c}} = \{\text{span}(e_1) \subseteq \text{span}(e_1, e_2) \subseteq F_3 \subseteq \text{span}(e_1, e_2, e_3, f_1)\}$ ,
- $K_{\mathbf{d}} = \{\text{span}(e_1) \subseteq \text{span}(e_1, e_2) \subseteq \text{span}(e_1, e_2, e_3) \subseteq F_4\}$ ,

each of which is an irreducible component of  $\mathcal{F}l^{((3),(1))}$  isomorphic to  $\mathbb{P}^1$ .

#### 4. TOPOLOGY OF EXOTIC SPRINGER FIBERS

**4.1. A topological model.** Let  $\mathbb{S}^2 \subseteq \mathbb{R}^3$  be the two-dimensional standard unit sphere with north pole  $p = (0, 0, 1)$ . Given a cup diagram  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ , define

$$S_{\mathbf{a}} = \{(x_1, \dots, x_m) \in (\mathbb{S}^2)^m \mid x_j = -x_i \text{ if } i-j, \text{ and } x_i = p \text{ if } \vdash i\}.$$

Note that there are no relations involving coordinates indexed by a vertex connected to a right-half-cup. Note that  $S_{\mathbf{a}}$  is homeomorphic to a product of two-spheres in which each cup and right-half-cup in  $\mathbf{a}$  contributes exactly one sphere. The  $((k), (m-k))$  *topological exotic Springer fiber*  $\mathcal{S}^{((k),(m-k))}$  is defined as the union

$$\mathcal{S}^{((k),(m-k))} := \bigcup_{\mathbf{a} \in \mathbb{B}^{((k),(m-k))}} S_{\mathbf{a}} \subseteq (\mathbb{S}^2)^m.$$

Let  $e, f$  be the standard basis of  $\mathbb{C}^2$  and let  $C: \mathbb{C}^{2N} \rightarrow \mathbb{C}^2$  be the linear map defined by  $C(e_i) = e$  and  $C(f_i) = f$ ,  $i \in \{1, \dots, N\}$ .

Consider the stereographic projection  $\sigma: \mathbb{S}^2 \setminus \{p\} \rightarrow \mathbb{C}$  and its analog for projective space

$$\theta: \mathbb{P}^1 \setminus \{\text{span}(e)\} \rightarrow \mathbb{C}; \text{span}(\alpha e + \beta f) \mapsto \frac{\alpha}{\beta}.$$

We can use  $\sigma$  and  $\theta$  to define a diffeomorphism  $\gamma: \mathbb{P}^1 \rightarrow \mathbb{S}^2$  by

$$\text{span}(\alpha e + \beta f) \mapsto \begin{cases} \sigma^{-1}(\theta(\text{span}(\alpha e + \beta f))) & \text{if } \text{span}(\alpha e + \beta f) \neq \text{span}(e), \\ p = (0, 0, 1) & \text{if } \text{span}(\alpha e + \beta f) = \text{span}(e). \end{cases}$$

This induces a diffeomorphism  $\gamma_m: (\mathbb{P}^1)^m \rightarrow (\mathbb{S}^2)^m$  on the  $m$ -fold products by setting  $\gamma_m(l_1, \dots, l_m) := (\gamma(l_1), \dots, \gamma(l_m))$ .

**Lemma 19.** *There is an equality of sets  $\gamma_m(T_{\mathbf{a}}) = S_{\mathbf{a}}$  for every  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ , where  $T_{\mathbf{a}} \subseteq (\mathbb{P}^1)^m$  is defined by*

$$T_{\mathbf{a}} := \{(l_1, \dots, l_m) \in (\mathbb{P}^1)^m \mid l_i^\perp = l_j \text{ if } i-j \text{ and } l_i = \text{span}(e) \text{ if } \vdash i\},$$

where the orthogonal complement is taken with respect to the standard Hermitian form on  $\mathbb{C}^2$ .

*Proof.* This follows directly from the definition of  $\gamma_m$ .  $\square$

**4.2. An explicit homeomorphism.** Recall the definition of the Hermitian form on  $\mathbb{C}^{2N}$  from the beginning of Section 3 as well as the definition of the variety  $Y_m$  in (4).

**Proposition 20** ([CK08, Theorem 2.1]). *The map  $\phi_m: Y_m \rightarrow (\mathbb{P}^1)^m$  given by the assignment*

$$(F_1, \dots, F_m) \mapsto (C(F_1), C(F_2 \cap F_1^\perp), \dots, C(F_m \cap F_{m-1}^\perp))$$

*is a diffeomorphism, where the orthogonal complement is taken with respect to the Hermitian form of  $\mathbb{C}^{2N}$ .*

Next, recall the definitions of the variety  $X_i^m \subseteq Y_m$  from (9) and the morphism  $q_m^i$  from (10).

**Lemma 21.** *The diffeomorphism  $\phi_m$  maps  $X_m^i$  bijectively to the set*

$$A_m^i := \{(l_1, \dots, l_m) \in (\mathbb{P}^1)^m \mid l_{i+1} = l_i^\perp\}. \quad (12)$$

Moreover, we have a commutative diagram

$$\begin{array}{ccc} X_m^i & \xrightarrow{q_m} & Y_{m-2} \\ \phi_m|_{X_m^i} \downarrow \cong & & \cong \downarrow \phi_{m-2} \\ A_m^i & \xrightarrow{f_m^i} & (\mathbb{P}^1)^{m-2} \end{array} \quad (13)$$

where  $f_m^i: (\mathbb{P}^1)^m \rightarrow (\mathbb{P}^1)^{m-2}$  is the map which forgets coordinates  $i$  and  $i+1$ . The orthogonal complement in (12) is taken with respect to the Hermitian structure of  $\mathbb{C}^2$ .

Furthermore, we have a commutative diagram

$$\begin{array}{ccc} Y_m & \xrightarrow{\psi_m} & Y_{m-1} \\ \phi_m \downarrow \cong & & \cong \downarrow \phi_{m-1} \\ (\mathbb{P}^1)^m & \xrightarrow{g_m} & (\mathbb{P}^1)^{m-1} \end{array} \quad (14)$$

where  $g_m$  (resp.  $\psi_m$ ) is the map which forgets the last coordinate (resp. vector space of the flag).

*Proof.* For the proof of the commutativity of the first diagram (13) we refer to [CK08, Theorem 2.1] or [Weh09, Lemma 2.4]. The commutativity of the second diagram (14) follows directly from the definitions of the respective maps.  $\square$

**Proposition 22.** *We have  $K_{\mathbf{a}} = \phi_m^{-1}(T_{\mathbf{a}})$ .*

*Proof.* We prove the proposition by induction on the number of cups in  $\mathbf{a}$ .

*Base of Induction.* We first prove Proposition 22 for all cup diagrams  $\mathbf{a} \in \mathbb{B}^{((k), (m-k))}$  as in (11) consisting of left-half-cups and right-half-cups only.

We first consider the cup diagram  $\mathbf{a} \in \mathbb{B}^{((m), (0))}$  consisting of  $m$  successive left-half-cups. In this case  $T_{\mathbf{a}}$  consists of a single element  $(\text{span}(e), \dots, \text{span}(e)) \in (\mathbb{P}^1)^m$  whose preimage  $(F_1, \dots, F_m)$  under the diffeomorphism  $\phi_m$  is given by the flag

$$(\text{span}(e_1), \text{span}(e_1, e_2), \dots, \text{span}(e_1, \dots, e_m))$$

since

$$\begin{aligned} \text{span}(e) &= C(\text{span}(e_i)) = C(\text{span}(e_1, \dots, e_i) \cap \text{span}(e_i, \dots, e_m, f_1, \dots, f_m)) \\ &= C(F_i \cap F_{i-1}^\perp). \end{aligned}$$

The relations in part (ii) of Theorem 14 show that this flag is indeed the unique element in  $K_{\mathbf{a}}$ .

Assume that there exists a right-half-cup in  $\mathbf{a} \in \mathbb{B}^{((k), (m-k))}$  (otherwise we are done by the above). Let  $\mathbf{a}' \in \mathbb{B}^{((k), (m-k-1))}$  be the cup diagram obtained by deleting the right-half-cup connected to the rightmost vertex  $m$ . Thus, we have

$$K_{\mathbf{a}} = \psi_m^{-1}(K_{\mathbf{a}'}) = \psi_m^{-1}(\phi_{m-2}^{-1}(T_{\mathbf{a}'})) = \phi_m^{-1}(g_m^{-1}(T_{\mathbf{a}'})) = \phi_m^{-1}(T_{\mathbf{a}}), \quad (15)$$

where the first equality is evident, the second one follows from the inductive hypothesis  $\phi_{m-2}^{-1}(T_{\mathbf{a}'}) = K_{\mathbf{a}'}$ , the third one follows from the commutativity of the diagram (14) and the last one is the obvious fact that  $g_m^{-1}(T_{\mathbf{a}'}) = T_{\mathbf{a}}$ .

*Inductive Step.* Now we assume that there exists at least one cup in  $\mathbf{a}$ . Thus, there exists a cup connecting neighboring vertices, say  $i$  and  $i+1$  (because the diagram is crossingless). Let  $\mathbf{a}'' \in \mathbb{B}^{((k-1), (m-k-1))}$  be the cup diagram obtained by removing this cup.

By an analogous argument as in the base case of the induction we obtain

$$K_{\mathbf{a}} = (q_m^i)^{-1}(K_{\mathbf{a}''}) = (q_m^i)^{-1}(\phi_{m-2}^{-1}(T_{\mathbf{a}''})) = \phi_m^{-1}((f_m^i)^{-1}(T_{\mathbf{a}''})) = \phi_m^{-1}(T_{\mathbf{a}}), \quad (16)$$

where the first equality follows from Lemma 16, the second one from the inductive hypothesis, the third one from the commutativity of the diagram (13) and the last one follows from  $(f_m^i)^{-1}(T_{\mathbf{a}''}) = T_{\mathbf{a}}$  which is evident.  $\square$

**Theorem 23.** *The diffeomorphism  $(\mathbb{S}^2)^m \xrightarrow{\gamma_m^{-1}} (\mathbb{P}^1)^m \xrightarrow{\phi_m^{-1}} Y_m$  restricts to a homeomorphism*

$$\mathcal{S}^{((k),(m-k))} \xrightarrow{\cong} \mathcal{F}l^{((k),(m-k))}$$

such that the images of the  $S_{\mathbf{a}}$  under this homeomorphism are precisely the irreducible components  $K_{\mathbf{a}}$  of  $\mathcal{F}l^{((k),(m-k))}$  for all  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ .

*Proof.* The image of  $\mathcal{S}^{((k),(m-k))} \subseteq (\mathbb{S}^2)^m$  under the diffeomorphism  $\phi_m^{-1} \circ \gamma_m^{-1}$  is given by


$$\begin{aligned} \phi_m^{-1} \left( \gamma_m^{-1} \left( \mathcal{S}^{((k),(m-k))} \right) \right) &= \bigcup_{\mathbf{a} \in \mathbb{B}^{((k),(m-k))}} \phi_m^{-1} \left( \gamma_m^{-1} (S_{\mathbf{a}}) \right) \\ &= \bigcup_{\mathbf{a} \in \mathbb{B}^{((k),(m-k))}} \phi_m^{-1} (T_{\mathbf{a}}) \quad (\text{by Lemma 19}) \\ &= \bigcup_{\mathbf{a} \in \mathbb{B}^{((k),(m-k))}} K_{\mathbf{a}}. \quad (\text{by Proposition 22}) \end{aligned}$$

The last line of the above chain of equalities equals  $\mathcal{F}l^{((k),(m-k))}$  by Theorem 14.  $\square$

*Example 24.* The submanifolds of  $(\mathbb{S}^2)^4$  associated with the cup diagrams in  $\mathbb{B}^{((3),(1))}$  (see Example 8) are the following:

$$\begin{aligned} \bullet S_{\mathbf{a}} &= \{(x, -x, p, p) \mid x \in \mathbb{S}^2\}, & \bullet S_{\mathbf{c}} &= \{(p, p, x, -x) \mid x \in \mathbb{S}^2\}, \\ \bullet S_{\mathbf{b}} &= \{(p, x, -x, p) \mid x \in \mathbb{S}^2\}, & \bullet S_{\mathbf{d}} &= \{(p, p, p, x) \mid x \in \mathbb{S}^2\}. \end{aligned}$$

Each of these manifolds is homeomorphic to a two-sphere. Their pairwise intersection is either a point or empty, e.g. we have  $S_{\mathbf{a}} \cap S_{\mathbf{b}} = \{(p, -p, p, p)\}$  and  $S_{\mathbf{a}} \cap S_{\mathbf{c}} = \emptyset$  (see also Theorem 28 for a more general statement). The  $((3),(1))$  topological exotic Springer fiber is a chain of four two-spheres:



More generally, it is straightforward to check that the  $((m-1),(1))$  topological exotic Springer fiber is chain of  $m$  two-spheres.


### 4.3. Intersections of irreducible components.

**Definition 25.** Let  $\mathbf{a}, \mathbf{b} \in \mathbb{B}^{((k),(m-k))}$ . The *circle diagram*  $\bar{\mathbf{a}}\mathbf{b}$  is defined as the diagram obtained by reflecting the diagram  $\mathbf{a}$  in its horizontal middle line and then sticking the resulting diagram, denoted by  $\bar{\mathbf{a}}$ , on top of the cup diagram  $\mathbf{b}$ , i.e. we glue the two diagrams along the horizontal line containing the vertices (thereby identifying the vertices of  $\bar{\mathbf{a}}$  and  $\mathbf{b}$  pairwise from left to right).

*Remark 26.* In general the diagram  $\bar{\mathbf{a}}\mathbf{b}$  consists of several connected components each of which is either a circle (i.e. a closed connected component) or a line segment. Note that there are three types of line segments: lines with both endpoints connected to the right vertical edge of the rectangle (we refer to this edge as the right wall), lines with both endpoints connected to the left vertical edge (also referred to as the left wall), and lines with one endpoint connected to the left wall and the other one connected to the right wall. A line which contains a half-cup (either left or right) of  $\mathbf{a}$  and a half-cup (either left or right) of  $\mathbf{b}$  is called a *propagating line*.

*Example 27.* Here is an example illustrating the gluing of two cup diagrams in order to obtain a circle diagram:





Hence,  $\bar{\mathbf{a}}\mathbf{b}$  consists of one circle and three line segments. The line with both endpoints connected to the left (resp. right) wall is propagating and the line which connects the left and the right wall is an example of a non-propagating line.

**Theorem 28.** *Let  $\mathbf{a}, \mathbf{b} \in \mathbb{B}^{((k), (m-k))}$ . We have  $S_{\mathbf{a}} \cap S_{\mathbf{b}} \neq \emptyset$  if and only if all lines containing two left-half-cups in  $\bar{\mathbf{a}}\mathbf{b}$  are propagating. Moreover, if  $S_{\mathbf{a}} \cap S_{\mathbf{b}} \neq \emptyset$ , we have a homeomorphism  $S_{\mathbf{a}} \cap S_{\mathbf{b}} \cong (\mathbb{S}^2)^k$ , where  $k$  counts the number of circles plus the number of line segments in  $\bar{\mathbf{a}}\mathbf{b}$  with both endpoints connected to the right wall.*

*Proof.* Assume that all lines in  $\bar{\mathbf{a}}\mathbf{b}$  containing two left-half-cups are propagating. Given a circle or a line with two endpoints connected to the right wall in  $\bar{\mathbf{a}}\mathbf{b}$ , let  $i$  denote the leftmost vertex (i.e.  $i$  is the smallest element) on this component. Choose an element  $x_i \in (\mathbb{S}^2)^m$ . This determines all other  $x_j$  for  $j$  in the component by applying the relation  $x_k = -x_l$  for the cups on the circle. Note that a circle always consists of an even number of cups. Hence, going around the circle once therefore yields an even number of sign changes and the relations are consistent. For line segments with both endpoints at the right wall there is nothing to show. Given a line, we pick a vertex  $i$  connected to a left-half-cup and set  $x_i = p$ . The remaining coordinates in the component are then determined. Note that for a given line the coordinate relations are consistent because the line is propagating and thus there is an even number of cups between the two left-half-cups. For a line segment with one endpoint at the right wall the relations are always consistent. Projecting onto the leftmost endpoints of circles and line segments with both endpoints at the right wall yields the homeomorphism  $S_{\mathbf{a}} \cap S_{\mathbf{b}} \cong (\mathbb{S}^2)^k$ .

On the other hand, if there is a non-propagating line containing two left-half-cups in  $\bar{\mathbf{a}}\mathbf{b}$ , there is an odd number of cups connecting the two left-half-cups of the line. Hence the relations coming from the cups are not consistent with the ones coming from the left-half-cups. Hence, the intersection  $S_{\mathbf{a}} \cap S_{\mathbf{b}}$  must be empty.  $\square$

## 5. $\mathbb{C}^*$ -ACTION AND BIALYNICKI-BIRULA PAVING

There is a  $\mathbb{C}^*$ -action on  $\mathbb{C}^{2N}$  given by  $t.e_i = te_i$  and  $t.f_i = t^{-1}f_i$  which yields a  $\mathbb{C}^*$ -action on  $Y_m$ . This action restricts to  $\mathcal{F}l^{((k), (m-k))} \subseteq Y_m$ . In order to study the isolated fixed points of this  $\mathbb{C}^*$ -action on the exotic Springer fiber we introduce the notion of a combinatorial weight.

**Definition 29.** A *combinatorial weight of type  $((k), (m-k))$*  is a  $m$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_m)$  of symbols  $\alpha_i \in \{\wedge, \vee\}$  with at most  $m-k$   $\vee$ 's. Let  $W^{((k), (m-k))}$  denote the set of all combinatorial weights of type  $((k), (m-k))$ .

*Example 30.* Here is a list of all combinatorial weights of type  $((3), (1))$ :

$$\alpha = \wedge \wedge \wedge \wedge, \quad \beta = \vee \wedge \wedge \wedge, \quad \gamma = \wedge \vee \wedge \wedge, \quad \delta = \wedge \wedge \vee \wedge, \quad \epsilon = \wedge \wedge \wedge \vee.$$

Given  $\alpha \in W^{((k), (m-k))}$ , we define  $F_l^\alpha = \text{span}(e_1, \dots, e_i, f_1, \dots, f_j)$ , where  $i$  is the number of  $\wedge$ 's amongst the first  $l$  entries  $\alpha_1, \dots, \alpha_l$  of  $\alpha$  and  $j$  is the number of  $\vee$ 's amongst them. This yields a flag  $(F_1^\alpha, \dots, F_m^\alpha) \in Y_m$  associated to  $\alpha \in W^{((k), (m-k))}$ . Note that we have  $e_k \in F_m^\alpha$  because by assumption there are at least  $k$   $\wedge$ 's in  $\alpha$ . In particular,  $(F_1^\alpha, \dots, F_m^\alpha) \in \mathcal{F}l^{((k), (m-k))}$ .

**Lemma 31.** *The map  $\alpha \mapsto (F_1^\alpha, \dots, F_m^\alpha)$  defines a bijection between  $W^{((k), (m-k))}$  and the fixed points under the  $\mathbb{C}^*$ -action on  $\mathcal{F}l^{((k), (m-k))}$ .*

*Proof.* The flag  $(F_1^\alpha, \dots, F_m^\alpha) \in Y_m$  is clearly a  $\mathbb{C}^*$ -fixed point for every  $\alpha \in W^{((k), (m-k))}$  because each of its vector spaces is spanned by weight vectors with respect to the  $\mathbb{C}^*$ -action. Thus, we have a well-defined map which is injective by construction.

In order to see the surjectivity let  $(F_1, \dots, F_m) \in Y_m$  be a flag fixed by the  $\mathbb{C}^*$ -action. By induction on  $l$  one can see that  $F_l$  must be spanned by  $e_1, \dots, e_r, f_1, \dots, f_s$  for some  $r, s$

in order to be fixed by the  $\mathbb{C}^*$ -action. In particular, in doing so, one can simultaneously construct the preimage  $\alpha$  of  $(F_1, \dots, F_m)$ . Since  $e_k \in F_m$ , there must be at least  $k$   $\wedge$ 's in  $\alpha$  or equivalently at most  $m - k$   $\vee$ 's and it follows that  $\alpha \in W^{((k), (m-k))}$ .  $\square$

Given a combinatorial weight  $\alpha \in W^{((k), (m-k))}$ , we associate with it a cup diagram  $C(\alpha)$  by successively connecting pairs of symbols  $\vee \wedge$  by a cup (ignoring the symbols which are already paired). If there is no more  $\vee$  to the left of a  $\wedge$  we attach a left-half-cup to all remaining  $\wedge$ 's and a right-half-cup to all remaining  $\vee$ 's.

*Example 32.* The cup diagrams associated with the combinatorial weights from Example 30 are given by

$$C(\alpha) = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} \quad , \quad C(\beta) = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} \quad , \quad C(\gamma) = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array}$$

$$C(\delta) = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} \quad , \quad C(\epsilon) = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array}$$

Let  $P = \text{span}(e_1, \dots, e_N)$  and  $Q = \text{span}(f_1, \dots, f_N)$ . Given a flag  $(F_1, \dots, F_m) \in \mathcal{F}l^{((k), (m-k))}$ , we associate with it a new flag  $(F_1^{\text{ass}}, \dots, F_m^{\text{ass}}) \in \mathcal{F}l^{((k), (m-k))}$  by setting  $F_i^{\text{ass}} = P_i + Q_i \subseteq P \oplus Q = V$ , where  $P_i = F_i \cap P$  and  $Q_i = (F_i + P)/P$  (we identify  $V/P \cong Q$ ). Note that this flag is fixed under the  $\mathbb{C}^*$ -action, i.e.  $(F_1^{\text{ass}}, \dots, F_m^{\text{ass}}) = (F_1^\alpha, \dots, F_m^\alpha)$  for some combinatorial weight  $\alpha$ .

**Lemma 33.** *A flag  $(F_1, \dots, F_m)$  in  $\mathcal{F}l^{((k), (m-k))}$  is contained in the attracting cell*

$$K_\alpha := \{(F_1, \dots, F_m) \in \mathcal{F}l^{((k), (m-k))} \mid \lim_{t \rightarrow \infty} t \cdot (F_1, \dots, F_m) = (F_1^\alpha, \dots, F_m^\alpha)\}$$

*corresponding to  $\alpha \in W^{((k), (m-k))}$  if and only if  $(F_1^{\text{ass}}, \dots, F_m^{\text{ass}}) = (F_1^\alpha, \dots, F_m^\alpha)$ .*

*Proof.* This follows as in the proof of [SW12, Proposition 14].  $\square$

**Theorem 34.** *The attracting cell*

$$K_\alpha = \{(F_1, \dots, F_m) \in \mathcal{F}l^{((k), (m-k))} \mid \lim_{t \rightarrow \infty} t \cdot (F_1, \dots, F_m) = (F_1^\alpha, \dots, F_m^\alpha)\}$$

*associated with the  $\mathbb{C}^*$ -fixed point  $(F_1^\alpha, \dots, F_m^\alpha) \in Y_m$  consists of all flags  $(F_1, \dots, F_m) \in \mathcal{F}l^{((k), (m-k))}$  satisfying the following conditions imposed by the cup diagram  $C(\alpha)$ :*

$$(i) F_j = x^{-\frac{1}{2}(j-i+1)} F_{i-1} \quad \text{if } i-j \quad (ii) F_i = F_{i-1} + \text{span}\left(e_{\frac{1}{2}(i+\rho_{\mathbf{a}}(i))}\right) \quad \text{if } \vdash i$$

*where  $\rho_{\mathbf{a}}(i)$  is as in Theorem 23. Additionally, we have the condition  $F_i \cap P = F_{i-1} \cap P$  whenever  $i$  is connected to a left endpoint of a cup or a right-half-cup.*

*Via the homeomorphism from Theorem 23, the attracting cell  $S_\alpha$  corresponding to  $\alpha$  consists of all  $(x_1, \dots, x_m) \in (\mathbb{S}^2)^m$  which satisfy the conditions  $x_i = -x_j$  and  $x_i \neq p$  if  $i < j$  are connected by a cup in  $C(\alpha)$ ,  $x_i \neq p$  if  $i$  is connected to a right-half-cup, and  $x_i = p$  if  $i$  is connected to a left-half-cup in  $C(\alpha)$ .*

*Proof.* We prove the claim by induction on the number of cups in  $C(\alpha)$ . First assume that  $\alpha$  is of the form  $\wedge \wedge \dots \wedge$  such that  $C(\alpha)$  consists of left-half-cups only. Then  $(F_1^\alpha, \dots, F_m^\alpha)$  is given by  $F_i(\alpha) = \text{span}(e_1, \dots, e_i)$ . For any other flag  $(F_1, \dots, F_m)$  attracted to  $(F_1^\alpha, \dots, F_m^\alpha)$  we have  $P_i = F_i \cap P = \text{span}(e_1, \dots, e_i) = F_i(\alpha)$  and thus  $Q_i \cong 0$ . In particular, it follows that any such  $(F_1, \dots, F_m)$  must already be equal to  $(F_1^\alpha, \dots, F_m^\alpha)$ .

Next assume that  $\alpha$  is of the form  $\wedge \wedge \dots \wedge \vee \dots \vee$  such that  $C(\alpha)$  consists of a sequence of left-half-cups followed by a sequence of right-half-cups (assume there exists at least one  $\vee$ ). Let  $\alpha'$  be the sequence obtained by deleting the rightmost  $\vee$  in  $\alpha$ . By induction we have that  $K_{\alpha'} = K_{\alpha'}^0$ , where  $K_{\alpha'}^0$  is defined as the set of flags satisfying the conditions (i) and (ii) of the theorem. Since  $K_\alpha \subseteq \psi_m^{-1}(K_{\alpha'})$  we obtain

$$K_\alpha \subseteq \psi_m^{-1}(K_{\alpha'}) = \psi_m^{-1}(K_{\alpha'}^0),$$

where  $\psi_m^{-1}(K_{\alpha'}^0)$  consists of all flags such that  $F_1, \dots, F_{m-1}$  satisfy the relations (i) and (ii) of  $C(\alpha)$ , and  $F_m$  can be chosen freely. We even have

$$K_\alpha \subseteq \{(F_1, \dots, F_m) \in \psi_m^{-1}(K_{\alpha'}^0) \mid F_i \cap P = F_{i-1} \cap P\}$$

because if  $\alpha_i = \vee$ , then we have  $F_i \cap P = F_{i-1} \cap P$ . On the other hand, if  $P_m = F_m \cap P = F_{m-1} \cap P$  we see that  $F_m^{\text{ass}}$  is obtained from  $F_{m-1}^{\text{ass}}$  by adding a vector from  $Q$ . Thus, the inclusion is in fact an equality.

Assume that there exists a cup in  $C(\alpha)$  and assume that  $i$  and  $i+1$  are connected by a cup. In particular,  $\alpha$  has symbols  $\vee \wedge$  at positions  $i$  and  $i+1$ . Let  $\alpha'$  be the combinatorial weight obtained by removing these two symbols. By induction we have  $K_{\alpha'}$  satisfies the conditions of the theorem and since  $K_{\alpha} \subseteq (\psi_m^i)^{-1}(K_{\alpha'})$  (by the same argument as in [SW12]) we obtain  $K_{\alpha} \subseteq (\psi_m^i)^{-1}(K_{\alpha'}) = (\psi_m^i)^{-1}(K_{\alpha'}^0)$ . By arguing as in the case of a right-half-cup, we see that  $K_{\alpha} = \{(F_1, \dots, F_m) \in (\psi_m^i)^{-1}(K_{\alpha'}^0) \mid F_i \cap P = F_{i-1} \cap P\}$ .  $\square$

*Example 35.* The attracting cells corresponding to the diagrams in Example 32 are given by  $K_{\alpha} = \{(\text{span}(e_1), \text{span}(e_1, e_2), \text{span}(e_1, e_2, e_3), \text{span}(e_1, e_2, e_3, e_4))\}$  and

$$\begin{aligned} K_{\beta} &= \{(F_1, \text{span}(e_1, f_1), \text{span}(e_1, e_2, f_1), \text{span}(e_1, e_2, e_3, f_1)) \mid F_1 \neq \text{span}(e_1)\}, \\ K_{\gamma} &= \{(\text{span}(e_1), F_2, \text{span}(e_1, e_2, f_1), \text{span}(e_1, e_2, e_3, f_1)) \mid F_2 \neq \text{span}(e_1, e_2)\}, \\ K_{\delta} &= \{(\text{span}(e_1), \text{span}(e_1, e_2), F_3, \text{span}(e_1, e_2, e_3, f_1)) \mid F_3 \neq \text{span}(e_1, e_2, e_3)\}, \\ K_{\epsilon} &= \{(\text{span}(e_1), \text{span}(e_1, e_2), \text{span}(e_1, e_2, e_3), F_4) \mid F_4 \neq \text{span}(e_1, e_2, e_3, e_4)\}. \end{aligned}$$

Topologically, we have  $S_{\alpha} = \{(p, p, p, p)\}$  and

$$\begin{aligned} S_{\beta} &= \{(x, -x, p, p) \mid x \in \mathbb{S}^2 \setminus \{p\}\}, & S_{\gamma} &= \{(p, x, -x, p) \mid x \in \mathbb{S}^2 \setminus \{p\}\}, \\ S_{\delta} &= \{(p, p, x, -x) \mid x \in \mathbb{S}^2 \setminus \{p\}\}, & S_{\epsilon} &= \{(p, p, p, x) \mid x \in \mathbb{S}^2 \setminus \{p\}\}. \end{aligned}$$

**Corollary 36.** *The  $2i$ -th Betti number of the exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$  is  $b_{2i} = \binom{m}{i}$ ,  $0 \leq i \leq m-k$ .*

*Proof.* This is now a direct consequence of the paving constructed in the above theorem.  $\square$

The next goal will be to give a combinatorial description of the topology of the pairwise intersections of the generalized irreducible components using oriented circle diagrams (see Theorem 28 below).

**Definition 37.** Let  $\alpha, \beta \in W^{((k), (m-k))}$  and let  $\alpha C(\beta)$  be the diagram obtained by sticking the weight  $\alpha$  on top of the cup diagram  $C(\beta)$ . We say that  $\alpha C(\beta)$  is an *oriented cup diagram* if  $\alpha_i \neq \alpha_j$  for  $i-j$  and  $\alpha_i = \wedge$  if  $\mid -i$ . Furthermore, we allow any symbol  $\alpha_i \in \{\wedge, \vee\}$  if  $i$  is connected to a right-half-cup.

**Definition 38.** Given  $\alpha, \beta \in W^{((k), (m-k))}$ , an *oriented circle diagram* is obtained by sticking a weight  $\gamma \in W^{((k), (m-k))}$  between the two diagrams  $\overline{C(\alpha)}$  and  $C(\beta)$  such that both  $\gamma C(\alpha)$  and  $\gamma C(\beta)$  are oriented cup diagrams. This induces orientations of the connected components, i.e. the circle and lines, of the circle diagram. We say that a component is oriented clockwise (resp. counterclockwise) if the leftmost vertex on such a component is a  $\wedge$  (resp.  $\vee$ ).

The following theorem describes the intersections of the generalized (and thus in particular also the ordinary) irreducible components.

*Remark 39.* It is straightforward to check that there exists a  $\gamma \in W^{((k), (m-k))}$  such that  $\overline{C(\alpha)}\gamma C(\beta)$  is an oriented circle diagram if and only if all lines in  $\overline{C(\alpha)}C(\beta)$  containing two left-half-cups are propagating.

Let  $\alpha, \beta \in W^{((k), (m-k))}$  be such that  $S_{C(\alpha)} \cap S_{C(\beta)} \neq \emptyset$ . Then  $S_{C(\alpha)} \cap S_{C(\beta)} \cong (\mathbb{S}^2)^k$  by Theorem 28, where  $k$  counts the number of circles and line segments connected to the right wall. In particular, we can find  $2^k$  different  $\gamma \in W^{((k), (m-k))}$  such that  $\overline{C(\alpha)}\gamma C(\beta)$  is an oriented circle diagram because any circle or line segment with both endpoints connected to the right wall can be oriented either clockwise or counterclockwise. Since

$$H^*(S_{C(\alpha)} \cap S_{C(\beta)}, \mathbb{C}) \cong H^*((\mathbb{S}^2)^k, \mathbb{C}) \cong \mathbb{C}[X]/(X^2)^{\otimes k},$$

we obtain an isomorphism of vector spaces

$$H^*(S_{C(\alpha)} \cap S_{C(\beta)}, \mathbb{C}) \cong \mathbb{C}[\overline{C(\alpha)}\gamma C(\beta) \mid \overline{C(\alpha)}\gamma C(\beta) \text{ is oriented}]$$

sending an elementary tensor  $a_1 \otimes \dots \otimes a_k$ ,  $a_i \in \{1, X\} \subseteq \mathbb{C}[X]/(X^2)$  to the oriented circle diagram in which the components corresponding to spheres with  $a_i = X$  are oriented clockwise.

## 6. COHOMOLOGY RING OF EXOTIC SPRINGER FIBERS FOR ONE-ROW BIPARTITIONS

In the following we will write  $H_*(X)$  (resp.  $H^*(X)$ ) as a shorthand notation for  $H_*(X, \mathbb{Z})$  (resp.  $H^*(X, \mathbb{Z})$ ).

**Definition 40.** A *line diagram* is constructed by attaching  $m$  vertical lines, each possibly decorated with a single dot, to  $m$  vertices (numbered from left to right) on a horizontal line. Given  $U \subseteq \{1, \dots, m\}$ , let  $l_U$  denote the unique line diagram with dots precisely on the lines whose endpoints are not contained in  $U$ . We denote the set of all line diagrams on  $m$  vertices by  $\mathfrak{L}_m$ .

*Example 41.* The line diagrams in  $\mathfrak{L}_2$  are given as follows:

$$l_\emptyset = \begin{array}{|c|} \hline \circ \\ \hline \end{array} \begin{array}{|c|} \hline \circ \\ \hline \end{array}, \quad l_{\{1\}} = \begin{array}{|c|} \hline \\ \hline \end{array} \begin{array}{|c|} \hline \circ \\ \hline \end{array}, \quad l_{\{2\}} = \begin{array}{|c|} \hline \circ \\ \hline \end{array} \begin{array}{|c|} \hline \\ \hline \end{array}, \quad l_{\{1,2\}} = \begin{array}{|c|} \hline \\ \hline \end{array} \begin{array}{|c|} \hline \\ \hline \end{array}.$$

The two-sphere has a cell decomposition  $\mathbb{S}^2 = \{p\} \cup (\mathbb{S}^2 \setminus \{p\})$  consisting of a 0-cell and a 2-cell. Henceforth, we fix this CW-structure and equip  $(\mathbb{S}^2)^m$  with the Cartesian product CW-structure. Note that we obtain a bijection between  $\mathfrak{L}_m$  and the cells of the CW-complex  $(\mathbb{S}^2)^m$  by mapping a line diagram  $l_U$  to the cell  $C_{l_U}$ , where  $C_{l_U}$  is defined by choosing the 0-cell (resp. 2-cell) for the  $i$ th sphere in  $(\mathbb{S}^2)^m$  if the  $i$ th line of  $l_U$  is dotted (resp. undotted). Since the homology classes  $[C_{l_U}]$  of the cells form a basis of  $H_*((\mathbb{S}^2)^m)$  as a  $\mathbb{Z}$ -module, we obtain an isomorphism of  $\mathbb{Z}$ -modules

$$\mathbb{Z}[\mathfrak{L}_m] \xrightarrow{\cong} H_*((\mathbb{S}^2)^m); \quad l_U \mapsto [C_{l_U}], \quad (17)$$

where  $\mathbb{Z}[\mathfrak{L}_m]$  is the free  $\mathbb{Z}$ -module with basis  $\mathfrak{L}_m$ . The homological degree of a line diagram is given by twice the number of undotted lines.

*Example 42.* The cells of  $\mathbb{S}^2 \times \mathbb{S}^2$  corresponding to the line diagrams in  $\mathfrak{L}_2$  (see Example 41) are given by

$$\begin{aligned} \bullet C_{l_\emptyset} &= \{(p, p)\}, & \bullet C_{l_{\{2\}}} &= \{(p, x) \mid x \in \mathbb{S}^2 \setminus \{p\}\}, \\ \bullet C_{l_{\{1\}}} &= \{(x, p) \mid x \in \mathbb{S}^2 \setminus \{p\}\}, & \bullet C_{l_{\{1,2\}}} &= \{(x, y) \mid x, y \in \mathbb{S}^2 \setminus \{p\}\}. \end{aligned}$$

The homology class  $[C_{l_\emptyset}]$  (resp.  $[C_{l_{\{1,2\}}}]$ ) is a basis of  $H_0(\mathbb{S}^2 \times \mathbb{S}^2)$  (resp.  $H_4(\mathbb{S}^2 \times \mathbb{S}^2)$ ) and  $[C_{l_{\{1\}}}]$ ,  $[C_{l_{\{2\}}}]$  form a basis of  $H_2(\mathbb{S}^2 \times \mathbb{S}^2)$ .

**Definition 43.** An *enriched cup diagram* is a cup diagram (as in Definition 7) in which the cups and right-half-cups are allowed to be decorated with a single dot each. Additionally, every left-half-cup is decorated with a single dot. Let  $\widetilde{\mathbb{B}}^{((k),(m-k))}$  be the set of all enriched cup diagrams on  $m$  vertices such that the total number of cups plus right-half-cups is  $m - k$ .

*Example 44.* Here are two examples of enriched cup diagrams in  $\widetilde{\mathbb{B}}^{((4),(5))}$ :



Given  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ , let  $i_1 < i_2 < \dots < i_{m-k}$  denote the left endpoints of the cups and right-half-cups in  $\mathbf{a}$ . Note that we have a homeomorphism

$$\xi_{\mathbf{a}}: S_{\mathbf{a}} \xrightarrow{\cong} (\mathbb{S}^2)^{m-k}; \quad (x_1, \dots, x_m) \mapsto (y_1, \dots, y_{m-k}), \quad (18)$$

where  $y_j = x_{i_j}$ ,  $1 \leq j \leq m - k$ , and we equip  $S_{\mathbf{a}}$  with the structure of the CW-complex whose cells are defined as the preimages of the cells of  $(\mathbb{S}^2)^{m-k}$  under the homeomorphism (18). We obtain a bijection between  $\widetilde{\mathbb{B}}^{((k),(m-k))}$  and the cells of  $S_{\mathbf{a}}$  by mapping an enriched cup diagram  $M$  to the cell  $C_M$ , where a dotted cup (resp. undotted cup) means that we have chosen the 0-cell (resp. 2-cell) for the corresponding sphere via (18). Since the homology

classes of the the cells of  $S_{\mathbf{a}}$  form a basis of the homology  $H_*(S_{\mathbf{a}})$ , we obtain an isomorphism of  $\mathbb{Z}$ -modules

$$\mathbb{Z}[\widetilde{\mathbb{B}}^{((k),(m-k))}] \xrightarrow{\cong} H_*(S_{\mathbf{a}}); M \mapsto [C_M]. \quad (19)$$


An enriched cup diagram for which the number of undotted cups plus undotted right-half-cups is equal to  $l$  has homological degree  $2l$ .

**Definition 45.** Given an enriched cup diagram  $M \in \widetilde{\mathbb{B}}^{((k),(m-k))}$ , we define the associated *line diagram sum* as

$$L_M = \sum_{U \in \mathcal{U}_M} (-1)^{\Lambda_M(U)} l_U \in \mathbb{Z}[\mathfrak{L}_m],$$

where  $\mathcal{U}_M$  is the set of all subsets  $U \subseteq \{1, \dots, m\}$  containing precisely one endpoint of every undotted cup as well as the endpoints of every undotted right-half-cup in  $M$ . Moreover,  $\Lambda_M(U)$  counts the total number of right endpoints of cups in  $U$ .

*Example 46.* Consider the enriched cup diagram



We have

$$\mathcal{U}_M = \{\{4, 6, 7\}, \{4, 6, 8\}, \{5, 6, 7\}, \{5, 6, 8\}\}$$

with

$$\Lambda_M(\{4, 6, 7\}) = 0, \Lambda_M(\{4, 6, 8\}) = \Lambda_M(\{5, 6, 7\}) = 1 \text{ and } \Lambda_M(\{5, 6, 8\}) = 2.$$

Therefore, the associated line diagram sum is

$$L_M = l_{\{4,6,7\}} - l_{\{4,6,8\}} - l_{\{5,6,7\}} + l_{\{5,6,8\}}.$$

**Lemma 47.** *The map  $(\psi_{\mathbf{a}})_*: H_*(S_{\mathbf{a}}) \rightarrow H_*((\mathbb{S}^2)^m)$  induced by the natural inclusion*

$$\psi_{\mathbf{a}}: S_{\mathbf{a}} \hookrightarrow (\mathbb{S}^2)^m$$

*is explicitly given by the assignment  $M \mapsto L_M$  (via identifications (17) and (19)).*

*Proof.* We first consider a cup diagram  $\mathbf{a}_0 \in \mathbb{B}^{((k),(m-k))}$  in which all  $l$  right-half-cups,  $0 \leq l \leq m - k$ , are connected to the rightmost vertices  $\{m - l + 1, \dots, m\}$ . Let  $\mathbf{a}'_0$  be the cup diagram on  $m - l$  vertices obtained by deleting the  $l$  rightmost vertices in  $\mathbf{a}_0$  and let  $\mathbf{a}''_0$  be the cup diagram on  $l$  vertices obtained by deleting the leftmost  $m - l$  vertices in  $\mathbf{a}_0$ . Then  $\mathbf{a}'_0$  consists of cups and left-half-cups only and  $\mathbf{a}''_0$  consists of right-half-cups only. Let  $M_0$  be an enriched cup diagram obtained by placing dots on  $\mathbf{a}_0$  and let  $M'_0$  (resp.  $M''_0$ ) be the corresponding diagram for  $\mathbf{a}'_0$  (resp.  $\mathbf{a}''_0$ ).

By [Rus11, §2], the manifold  $S_{\mathbf{a}'_0}$  also appears as a component of a topological Springer fiber of type A (by thinking of the left-half-cups as “rays”) and it follows from [RT11, Lemma 3.12] and [Rus11, §5] (see also [SW16, Proposition 12]) that the map  $H_*(S_{\mathbf{a}'_0}) \rightarrow H_*((\mathbb{S}^2)^{m-l})$  induced by the natural inclusion  $\psi_{\mathbf{a}'_0}$  is given by the assignment  $M'_0 \mapsto L_{M'_0}$  as claimed.

Next, note that we can identify  $S_{\mathbf{a}''_0}$  with  $(\mathbb{S}^2)^l$  and the inclusion  $\psi_{\mathbf{a}''_0}$  with the identity  $\text{id}_{(\mathbb{S}^2)^l}$ . In particular, the map  $(\psi_{\mathbf{a}''_0})_*$  is evidently given by  $M''_0 \mapsto l_V$ , where  $V \subseteq \{1, \dots, l\}$  contains all vertices connected to the undotted right-half-cups in  $M''_0$ .

Thus, by the preceding two paragraphs, we can compute the image of  $M'_0 \otimes M''_0$  under the map

$$(\psi_{\mathbf{a}'_0})_* \otimes (\psi_{\mathbf{a}''_0})_*: H_*(S_{\mathbf{a}'_0}) \otimes H_*(S_{\mathbf{a}''_0}) \rightarrow H_*((\mathbb{S}^2)^{m-l}) \otimes H_*((\mathbb{S}^2)^l) \quad (20)$$

as follows:

$$\begin{aligned} (\psi_{\mathbf{a}'_0})_* \otimes (\psi_{\mathbf{a}''_0})_*(M'_0 \otimes M''_0) &= L_{M'_0} \otimes l_V = \left( \sum_{U' \in \mathcal{U}_{M'_0}} (-1)^{\Lambda_{M'_0}(U')} l_{U'} \right) \otimes l_V \\ &= \sum_{U' \in \mathcal{U}_{M'_0}} (-1)^{\Lambda_{M'_0}(U')} (l_{U'} \otimes l_V). \end{aligned}$$

Note that the Künneth isomorphism

$$H_*(S_{\mathbf{a}'_0}) \otimes H_*(S_{\mathbf{a}''_0}) \cong H_*(S_{\mathbf{a}'_0} \times S_{\mathbf{a}''_0}) = H_*(S_{\mathbf{a}_0}) \quad (21)$$

sends  $M'_0 \otimes M''_0$  to  $M_0$  and

$$H_*((\mathbb{S}^2)^{m-l}) \otimes H_*((\mathbb{S}^2)^l) \cong H_*((\mathbb{S}^2)^{m-l} \times (\mathbb{S}^2)^l) = H_*((\mathbb{S}^2)^m) \quad (22)$$

sends  $l_{U'} \otimes l_V$  to  $l_U$ , where  $U$  contains all numbers in  $U'$  and all numbers obtained by adding  $m-l$  to the numbers in  $V$ . Since the map  $(\psi_{\mathbf{a}_0})_* = (\psi_{\mathbf{a}'_0} \times \psi_{\mathbf{a}''_0})_*$  is given by the composition of the maps in (20), (21) and (22), it is straightforward to check that it maps  $M_0$  to  $L_{M_0}$  as claimed (note that  $\Lambda_{M'_0}(U') = \Lambda_{M_0}(U)$ ).

For the general case let  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$  be a cup diagram and let  $M$  be an enriched cup diagram obtained by decorating  $\mathbf{a}$  with dots. Let  $\mathbf{a}_0 \in \mathbb{B}^{((k),(m-k))}$  be the cup diagram obtained by rearranging the components of  $\mathbf{a}$  in such a way that the right-half-cups are all connected to the rightmost vertices and let  $M_0$  be the enriched cup diagram for  $\mathbf{a}_0$  in which a component has a dot if and only if the corresponding component of  $M$  has a dot. Define  $\tau_{\mathbf{a}}$  to be the permutation of the vertices realizing this rearrangement. We obtain an induced homeomorphism  $\tau_{\mathbf{a}}: (\mathbb{S}^2)^m \rightarrow (\mathbb{S}^2)^m$  which permutes the coordinates and restricts to a homeomorphism  $S_{\mathbf{a}} \rightarrow S_{\mathbf{a}_0}$ . We can then write the inclusion  $\psi_{\mathbf{a}}: S_{\mathbf{a}} \hookrightarrow (\mathbb{S}^2)^m$  as the composition  $\tau_{\mathbf{a}} \circ \psi_{\mathbf{a}_0} \circ (\tau_{\mathbf{a}}^{-1}|_{S_{\mathbf{a}}})$ . Then the claim in Lemma 47 follows from the following calculation:

$$\begin{aligned} (\psi_{\mathbf{a}})_*(M) &= (\tau_{\mathbf{a}})_* \circ (\psi_{\mathbf{a}_0})_* \circ (\tau_{\mathbf{a}}^{-1}|_{S_{\mathbf{a}}})_*(M) = (\tau_{\mathbf{a}})_* \circ (\psi_{\mathbf{a}_0})_*(M_0) \\ &= (\tau_{\mathbf{a}})_* \left( \sum_{U \in \mathcal{U}_{M_0}} (-1)^{\Lambda_{M_0}(U)} l_U \right) = \sum_{U \in \mathcal{U}_M} (-1)^{\Lambda_M(U)} l_U. \quad \square \end{aligned}$$


For technical purposes we introduce a map

$$\beta^{((k),(m-k))}: \bigsqcup_{l=0}^{m-k} \mathbb{B}^{((m-l),(l))} \longrightarrow \widetilde{\mathbb{B}}^{((k),(m-k))}.$$

In order to define this map let  $\mathbf{a} \in \mathbb{B}^{((m-l),(l))}$  be a cup diagram,  $0 \leq l \leq m-k$ . Let  $T_{\mathbf{a}}$  be the standard one-row bitableau of shape  $((m-l), (l))$  associated to  $\mathbf{a}$  via Lemma 9 and let  $\widetilde{T}_{\mathbf{a}}$  be the unique standard one-row bitableau of shape  $((k), (m-k))$  obtained by moving the  $m-l-k$  largest entries in the left tableau of  $T_{\mathbf{a}}$  to the right tableau (and reordering the right tableau to make it standard). The tableau  $\widetilde{T}_{\mathbf{a}}$  gives rise to a cup diagram in  $\mathbb{B}^{((m-k),(k))}$  (again, via Lemma 9) which we additionally decorate by putting a dot on all left-half-cups as well as on each component which is connected to a vertex labeled by one of the  $m-l-k$  numbers which we moved. We define the resulting enriched cup diagram in  $\widetilde{\mathbb{B}}^{((k),(m-k))}$  as the image of  $\mathbf{a}$  under  $\beta^{((k),(m-k))}$ .


*Example 48.* Let  $m = 5$  and  $k = 3$ . We give an example of the map  $\beta^{((3),(2))}$  from

$$\mathbb{B}^{((5),(-))} \sqcup \mathbb{B}^{((4),(1))} \sqcup \mathbb{B}^{((3),(2))} \longrightarrow \widetilde{\mathbb{B}}^{((3),(2))}.$$


(1) Let  $\mathbf{a} =$    $\in \mathbb{B}^{((5),(-))}$ . The bitableau associated to  $\mathbf{a}$  is

$$T_{\mathbf{a}} = (\boxed{1 \ 2 \ 3 \ 4 \ 5}, -).$$

Since  $l = 0$  in this case, we have  $m-l-k = 2$  and so  $\widetilde{T}_{\mathbf{a}} = (\boxed{1 \ 2 \ 3}, \boxed{4 \ 5})$ , which yields the following image:


$$\beta^{((3),(2))}(\mathbf{a}) =$$



$$\in \widetilde{\mathbb{B}}^{((3),(2))}.$$

(2) Now let  $\mathbf{b} =$    $\in \mathbb{B}^{((4),(1))}$ . The bitableau associated to  $\mathbf{b}$  is

$$T_{\mathbf{b}} = (\boxed{1 \ 2 \ 4 \ 5}, \boxed{3}).$$


In this case  $l = 1$  and so  $m - k - l = 1$  giving  $\tilde{T}_{\mathbf{b}} = (\boxed{1 \ 2 \ 4}, \boxed{3 \ 5})$ . Therefore:

$$\beta^{((3),(2))}(\mathbf{b}) =$$
   $\in \tilde{\mathbb{B}}^{((3),(2))}$ .

(3) As a final example, let  $\mathbf{c} =$    $\in \mathbb{B}^{((3),(2))}$ . In this case we have


$$T_{\mathbf{c}} = \tilde{T}_{\mathbf{c}} = (\boxed{3 \ 4 \ 5}, \boxed{1 \ 2})$$

since  $l = 0$ . Therefore we have,

$$\beta^{((3),(2))}(\mathbf{c}) =$$
   $\in \tilde{\mathbb{B}}^{((3),(2))}$ .

**Definition 49.** An enriched cup diagram in  $\tilde{\mathbb{B}}^{((k),(m-k))}$  contained in the image of  $\beta^{((k),(m-k))}$  is called a *standard enriched cup diagram*. We write  $s\tilde{\mathbb{B}}^{((k),(m-k))}$  to denote the set of all standard enriched cup diagrams on  $m$  vertices such that the total number of cups and right-half-cups is  $m - k$ .

*Example 50.* The set  $s\tilde{\mathbb{B}}^{((3),(1))}$  consists of the following five diagrams:



The first 4 diagrams constitute the image of the set  $\mathbb{B}^{((3),(1))}$  and the last diagram is the image of  $\mathbb{B}^{((4),(-))}$  under the map  $\beta^{((3),(1))}$ .

*Remark 51.* The left endpoints of the cups and right-half-cups in  $\mathbf{a} \in \mathbb{B}^{((m-l),(l))}$  are precisely the left endpoints of the undotted cups and undotted right-half-cups in  $\beta^{((k),(m-k))}(\mathbf{a})$ . Thus,  $\beta^{((k),(m-k))}$  maps  $\mathbb{B}^{((m-l),(l))}$  to the enriched cup diagrams of degree  $2l$ . Moreover, since the sets containing all vertices connected to the left endpoints of cups and right-half-cups are different for  $\mathbf{a}, \mathbf{b} \in \mathbb{B}^{((m-l),(l))}$  if  $\mathbf{a} \neq \mathbf{b}$  (because these vertices are precisely the entries of the right tableau in the bitableau associated to a cup diagram via the bijection in Lemma 9), it follows that the standard enriched cup diagrams associated to  $\mathbf{a}$  and  $\mathbf{b}$  via  $\beta^{((k),(m-k))}$  must be different. In other words, a standard enriched cup diagram is uniquely determined by its left endpoints of undotted cups and undotted right-half-cups.

**Lemma 52.** *The line diagram sums  $L_M$ , where  $M$  varies over all standard enriched cup diagrams in  $s\tilde{\mathbb{B}}^{((k),(m-k))}$ , are  $\mathbb{Z}$ -linearly independent in  $\mathbb{Z}[\mathfrak{L}_m]$ .*

*Proof.* Note that it suffices to prove that the elements  $L_M$ , where  $M$  varies over all standard enriched cup diagrams in  $s\tilde{\mathbb{B}}^{((k),(m-k))}$  such that the total number of undotted cups and undotted right-half-cups is precisely  $l$ ,  $0 \leq l \leq m$ , are  $\mathbb{Z}$ -linearly independent. In the following we write  $s\tilde{\mathbb{B}}_l^{((k),(m-k))}$  to denote the set of all such standard enriched cup diagrams.

We begin by defining a total order on the subsets of  $\{1, \dots, m\}$  of cardinality  $l$  by setting

$$\{i_1 < \dots < i_l\} < \{i'_1 < \dots < i'_l\} \Leftrightarrow \exists r : i_r < i'_r \text{ and } i_1 = i'_1, \dots, i_{r-1} = i'_{r-1}. \quad (23)$$

This induces a total order on all line diagrams  $l_U \in \mathfrak{L}_m$ , where  $U \subseteq \{1, \dots, m\}$  has cardinality  $l$ , i.e. we obtain a total order on our basis of the submodule of  $\mathbb{Z}[\mathfrak{L}_m]$  spanned by all line diagrams of degree  $2l$ .

Given a standard enriched cup diagram  $M \in \widetilde{s\mathbb{B}}_l^{((k),(m-k))}$ , we define  $U_M \in \mathcal{U}_M$  as the set containing the left endpoints of all undotted cups as well as the endpoints of all undotted right-half-cups. Given a different standard enriched cup diagrams  $N \in \widetilde{s\mathbb{B}}_l^{((k),(m-k))} \setminus \{M\}$ , we have  $U_M \neq U_N$  because knowing the left endpoints of undotted cups and undotted right-half-cups uniquely determines a standard enriched cup diagram in  $\widetilde{s\mathbb{B}}_l^{((k),(m-k))}$  by Remark 51. Hence, it makes sense to define a total order on  $\widetilde{s\mathbb{B}}_l^{((k),(m-k))}$  by setting  $M < N$  if and only if  $U_M < U_N$  with respect to the order (23).

Let  $M_1 < M_2 < \dots < M_{\binom{m}{l}}$  denote the elements of  $\widetilde{s\mathbb{B}}_l^{((k),(m-k))}$  and assume  $\sum_i \lambda_i L_{M_i} = 0$  for  $\lambda_i \in \mathbb{Z}$ . Then the line diagram  $l_{U_{M_1}}$  appears in  $L_{M_1}$  with non-zero coefficient but it does not appear in any other  $L_{M_i}$ ,  $i \neq 1$ , because we have  $l_{U_{M_1}} < l_{U_{M_i}}$  and  $l_{U_{M_i}} < l_U$  for all  $U \in \mathcal{U}_{M_i} \setminus \{U_{M_i}\}$ . Hence, we deduce that  $\lambda_1 = 0$ . By repeating the same argument with  $M_2 < \dots < M_{\binom{m}{l}}$  we obtain that  $\lambda_2 = 0$  and continuing in this way shows that all coefficients must be zero.  $\square$

**Proposition 53.** *The natural inclusion of  $\mathcal{S}^{((k),(m-k))}$  into  $(\mathbb{S}^2)^m$  induces an injection  $H_*(\mathcal{S}^{((k),(m-k))}) \hookrightarrow H_*((\mathbb{S}^2)^m)$  in homology. We even have an isomorphism*

$$H_{2l}(\mathcal{S}^{((k),(m-k))}) \cong H_{2l}((\mathbb{S}^2)^m)$$

of  $\mathbb{Z}$ -modules for  $0 \leq l \leq m - k$ .

*Proof.* Given  $\mathbf{a} \in \mathbb{B}^{((k),(m-k))}$ , let  $\phi_{\mathbf{a}}: S_{\mathbf{a}} \hookrightarrow \mathcal{S}^{((k),(m-k))}$ ,  $\psi_{\mathbf{a}}: S_{\mathbf{a}} \hookrightarrow (\mathbb{S}^2)^m$  be the natural inclusions. We obtain a commutative diagram

$$\bigoplus_{\mathbf{a} \in \mathbb{B}^{((k),(m-k))}} H_*(S_{\mathbf{a}}) \begin{array}{c} \xrightarrow{\phi_{((k),(m-k))}} \\ \xrightarrow{\psi_{((k),(m-k))}} \end{array} H_*(\mathcal{S}^{((k),(m-k))}) \xrightarrow{\gamma_{((k),(m-k))}} H_*((\mathbb{S}^2)^m), \quad (24)$$

where  $\phi_{((k),(m-k))}$  (resp.  $\psi_{((k),(m-k))}$ ) is the direct sum of the maps induced by  $\phi_{\mathbf{a}}$  (resp.  $\psi_{\mathbf{a}}$ ) and  $\gamma_{((k),(m-k))}$  is induced by the inclusion  $\mathcal{S}^{((k),(m-k))} \hookrightarrow (\mathbb{S}^2)^m$ .

We have a chain of equalities

$$\gamma_{((k),(m-k))}(\phi_{((k),(m-k))}(M)) = \psi_{((k),(m-k))}(M) = L_M,$$

where the first equality follows from the commutative diagram (24) and the second one follows from Lemma 47. Hence, by Lemma 52, we see that the elements

$$\gamma_{((k),(m-k))}(\phi_{((k),(m-k))}(M))$$

are linearly independent if  $M$  varies over all the whole set  $\widetilde{s\mathbb{B}}^{((k),(m-k))}$ . In particular, it follows that  $\phi_{((k),(m-k))}(M)$  must be  $\mathbb{Z}$ -linearly independent for  $M \in \widetilde{s\mathbb{B}}^{((k),(m-k))}$ . Since the cardinality of  $\widetilde{s\mathbb{B}}^{((k),(m-k))}$  equals the dimension of  $H_*(\mathcal{S}^{((k),(m-k))})$  by Corollary 36 and Remark 51, these elements form a basis of  $H_*(\mathcal{S}^{((k),(m-k))})$ . In particular, it directly follows that the map  $\gamma_{((k),(m-k))}$  is injective.  $\square$

**Theorem 54.** *The following statements hold:*

- (1) *There exists a homotopy equivalence*

$$\mathcal{F}l^{((k),(m-k))} \simeq \text{Sk}_{2(m-k)}^m = \bigcup_{\substack{U \subseteq \{1, \dots, m\} \\ |U|=m-k}} C_U \subseteq (\mathbb{S}^2)^m \quad (25)$$

*between the  $((k),(m-k))$ -exotic Springer fiber and the  $2(m-k)$ -skeleton  $\text{Sk}_{2(m-k)}^m$  of the CW-complex  $(\mathbb{S}^2)^m$ .*

- (2) *We have an isomorphism of graded algebras*

$$H^*(\mathcal{F}l^{((k),(m-k))}, \mathbb{C}) \cong H^*(\text{Sk}_{2(m-k)}^m, \mathbb{C}). \quad (26)$$

In particular, using (26), we obtain an explicit presentation of the graded algebra  $H^*(\mathcal{F}l^{((k),(m-k))}, \mathbb{C})$  given by

$$\mathbb{C}[X_1, \dots, X_m] \Big/ \left\langle X_i^2, X_I \mid I \subseteq \{1, \dots, m\}, |I| = m - k + 1, 1 \leq i \leq m \right\rangle, \quad (27)$$

where  $X_I = \prod_{i \in I} X_i$  and  $\deg(X_i) = 2$ . Furthermore, the presentation (27) yields a distinguished basis of  $H^{2l}(\mathcal{F}l^{((k),(m-k))}, \mathbb{C})$  given by all monomials  $X_I$  such that  $|I| = l$ ,  $I \subseteq \{1, \dots, m\}$ ,  $0 \leq l \leq m - k$ .

*Proof.* Using the homeomorphism from Theorem 23 we can work with  $\mathcal{S}^{((k),(m-k))}$  instead of  $\mathcal{F}l^{((k),(m-k))}$  when arguing topologically. By the cellular approximation theorem we can replace the natural inclusion of  $\mathcal{S}^{((k),(m-k))}$  into  $(\mathbb{S}^2)^m$  (which in general is not a cellular map) by a homotopic map  $j$  which factors through  $\text{Sk}_{2(m-k)}^m$  (note that by Remark 11 the complex dimension of  $\mathcal{F}l^{((k),(m-k))}$  is  $m - k$  so there are no cells of real dimension greater than  $2(m - k)$ ) and we obtain a commutative diagram

$$\begin{array}{ccc} \mathcal{S}^{((k),(m-k))} & \xrightarrow{j} & (\mathbb{S}^2)^m \\ & \searrow j & \nearrow \\ & \text{Sk}_{2(m-k)}^m & \end{array} \quad (28)$$

The horizontal map  $j$  in (28) induces isomorphisms  $H_{2l}(\mathcal{S}^{((k),(m-k))}) \cong H_{2l}((\mathbb{S}^2)^m)$  for all  $0 \leq l \leq m - k$  because the natural inclusion induces such isomorphisms by Proposition 53. Since the natural inclusion of  $\text{Sk}_{2(m-k)}^m$  into  $(\mathbb{S}^2)^m$  is well known to induce isomorphisms  $H_{2l}(\text{Sk}_{2(m-k)}^m) \cong H_{2l}((\mathbb{S}^2)^m)$  for all  $0 \leq l \leq m - k$ , we deduce that  $j$  induces isomorphisms  $H_{2l}(\mathcal{S}^{((k),(m-k))}) \cong H_{2l}(\text{Sk}_{2(m-k)}^m)$  for all  $0 \leq l \leq m - k$ . Since  $\mathcal{S}^{((k),(m-k))}$  and  $\text{Sk}_{2(m-k)}^m$  are simply-connected CW-complexes we can apply the homological version of the Whitehead theorem to the map  $j$  to prove that it yields the desired homotopy equivalence  $\mathcal{F}l^{((k),(m-k))} \cong \mathcal{S}^{((k),(m-k))} \simeq \text{Sk}_{2(m-k)}^m$ .

Since  $\mathcal{S}^{((k),(m-k))} \simeq \text{Sk}_{2(m-k)}^m$  are homotopy equivalent it suffices to compute the cohomology ring of  $\text{Sk}_{2(m-k)}^m$ . Note that the map  $H^*((\mathbb{S}^2)^m) \rightarrow H^*(\text{Sk}_{2(m-k)}^m)$  induced by the inclusion  $\text{Sk}_{2(m-k)}^m \hookrightarrow (\mathbb{S}^2)^m$  is surjective. This follows from the standard isomorphisms  $H^{2l}(\text{Sk}_{2(m-k)}^m) \cong H^{2l}((\mathbb{S}^2)^m)$  for  $0 \leq l \leq m - k$  and it is obviously true for  $l > m - k$  because  $H^{2l}(\text{Sk}_{2(m-k)}^m) \cong H_{2l}(\text{Sk}_{2(m-k)}^m) \cong \{0\}$ . In particular, the kernel of the surjection is given by the ideal  $\bigoplus_{i>2(m-k)}^{\infty} H^i((\mathbb{S}^2)^m)$  and we obtain an isomorphism of algebras

$$\bigoplus_{i=0}^{\infty} H^i((\mathbb{S}^2)^m) \Big/ \bigoplus_{i>2(m-k)}^{\infty} H^i((\mathbb{S}^2)^m) \xrightarrow{\cong} H^*(\text{Sk}_{2(m-k)}^m). \quad (29)$$

Using the isomorphism

$$H^*((\mathbb{S}^2)^m) \cong \mathbb{C}[X_1, \dots, X_m] \Big/ \langle X_i^2 \mid 1 \leq i \leq m \rangle$$

we see that the quotient in (29) can be identified with the quotient in Theorem 54.

The statement about the basis of the algebra in (27) is evident.  $\square$

*Example 55.* Note that  $\text{Sk}_2^m$  is a bouquet of  $m$  two-spheres, i.e.  $m$  two-spheres glued together in a single point. It is a standard exercise to show that a bouquet of  $m$  two-spheres is homotopy equivalent to a chain of  $m$  two-spheres. In particular, for the special case  $\mathcal{F}l^{((m-1),(1))} \cong \mathcal{S}^{((m-1),(1))} \simeq \text{Sk}_2^m$  we already obtain the homotopy equivalence (25) from the discussion in Example 24. Note, that  $\text{Sk}_{2(m-k)}^m$  and  $\mathcal{F}l^{((k),(m-k))}$  are in general not homeomorphic, e.g. removing the gluing point of the bouquet  $\text{Sk}_2^m$  yields  $m$  distinct connected components, whereas removing any point in  $\mathcal{F}l^{((m-1),(1))} \cong \mathcal{S}^{((m-1),(1))}$  yields at most two connected components.

Let  $\mathcal{W}_{C_m}$  be the Weyl group of type  $C_m$ . This is a Coxeter group with generators  $s_0, s_1, \dots, s_{m-1}$  subject to the relations  $s_i^2 = e$  for all  $i$ , and for  $i, j \neq 0$   $s_i s_j = s_j s_i$  if

$|i - j| > 1$  and  $s_i s_j s_i = s_j s_i s_j$  if  $|i - j| = 1$ , and additionally  $s_0 s_1 s_0 s_1 = s_1 s_0 s_1 s_0$  and  $s_0 s_j = s_j s_0$  for  $j \neq 1$ .

The following proposition is a classical result, see e.g. [Ser77, §8.2], [CM93, Theorem 10.1.2], [Mac95, Appendix B] and also [MS16, Proposition 3] for a categorical approach.

**Proposition 56.** *There exists a bijection between complex, finite-dimensional, irreducible  $\mathcal{W}_{C_m}$ -modules (up to isomorphism) and bipartitions of  $m$ . Given a bipartition  $(\lambda, \mu)$  of  $m$ , we write  $V_{(\lambda, \mu)}$  to denote the corresponding irreducible  $\mathcal{W}_{C_m}$ -module.*

Let  $\sigma_0: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the homeomorphism given by  $(x, y, z) \mapsto (-x, y, z)$ . It induces a homeomorphism  $\mathbb{S}^2 \xrightarrow{\cong} \mathbb{S}^2$  which (by abuse of notation) we also denote by  $\sigma_0$ .

**Proposition 57.** *Let  $0 \leq l \leq m - k$ . The following statements hold:*

(a) *The Weyl group  $\mathcal{W}_{C_m}$  acts on  $\text{Sk}_{2(m-k)}^m$  by*

$$s_0 \cdot (x_1, x_2, \dots, x_m) = (\sigma_0(x_1), x_2, \dots, x_m),$$

$$s_i \cdot (x_1, \dots, x_i, x_{i+1}, \dots, x_m) = (x_1, \dots, x_{i+1}, x_i, \dots, x_m), \quad i \neq 0.$$

(b) *The induced  $\mathcal{W}_{C_m}$ -action on  $H^*(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  can be described explicitly on the monomial basis of  $H^{2l}(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  from Theorem 54 as follows: a generator  $s_i$ ,  $i \neq 0$ , acts by exchanging  $X_i$  and  $X_{i+1}$  and leaves  $X_j$  invariant for  $j \neq i, i + 1$ , and  $s_0$  acts by sending  $X_1$  to  $-X_1$  and leaves all other  $X_i$ ,  $i \neq 1$ , invariant.*

(c) *The action from (b) yields an isomorphism  $H^{2l}(\text{Sk}_{2(m-k)}^m, \mathbb{C}) \cong V_{((m-l), (l))}$  of  $\mathcal{W}_{C_m}$ -modules.*

*Proof.* Since  $\sigma_0$  is an involution of  $(\mathbb{S}^2)^m$  fixing the north pole  $p = (0, 0, 1)$ , we see that  $s_0 \cdot C_{l_U} = C_{l_U}$  for all  $U \subseteq \{1, \dots, m\}$ ,  $|U| = m - k$ . Moreover, for  $i \neq 0$  we have  $s_i \cdot C_{l_U} = C_{l_{s_i \cdot U}}$ , where  $s_i \cdot U \subseteq \{1, \dots, m\}$  is the set containing all elements obtained by applying the permutation  $s_i: \{1, \dots, m\} \rightarrow \{1, \dots, m\}$  switching  $i$  and  $i + 1$  to the elements in  $U$ . Since  $|s_i \cdot U| = m - k$  for  $1 \leq i \leq m - 1$ , we conclude that  $s_i \cdot (x_1, \dots, x_m) \in \text{Sk}_{2(m-k)}^m$  for all  $(x_1, \dots, x_m) \in \text{Sk}_{2(m-k)}^m$  and  $0 \leq i \leq m - 1$ . It is straightforward to check that the  $s_i$ ,  $0 \leq i \leq m - 1$ , satisfy the defining relations of  $\mathcal{W}_{C_m}$  and we obtain a well-defined  $\mathcal{W}_{C_m}$ -action on  $\text{Sk}_{2(m-k)}^m$  as claimed in Proposition 57(a).

Note that the  $\mathcal{W}_{C_m}$ -action on  $\text{Sk}_{2(m-k)}^m$  in (a) extends to an action on  $(\mathbb{S}^2)^m$ . Since the map  $\sigma_0$  is homotopic to the antipodal map it follows from standard algebraic topology that the induced action of  $s_0$  on  $H^*((\mathbb{S}^2)^m, \mathbb{C}) \cong \mathbb{C}[X_1, \dots, X_m]/(X_i^2)$  is given by sending  $X_1$  to  $-X_1$  and leaving all other  $X_i$  invariant,  $i \neq 1$ . Moreover, for  $i \neq 0$ , the induced action of  $s_i$  is given by exchanging  $X_i$  and  $X_{i+1}$  and leaving  $X_j$  invariant for  $j \neq i, i + 1$ . Since the inclusion  $\text{Sk}_{2(m-k)}^m \subseteq (\mathbb{S}^2)^m$  is clearly  $\mathcal{W}_{C_m}$ -equivariant, the above shows that the  $\mathcal{W}_{C_m}$ -action on  $\text{Sk}_{2(m-k)}^m$  defined in (a) induces the action on  $H^*(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  as claimed in (b) via the isomorphism (29).

In order to identify the irreducible representation in each cohomological degree we first note that the  $\mathcal{W}_{C_m}$ -action on  $(\mathbb{S}^2)^m$  obtained by extending the action on  $\text{Sk}_{2(m-k)}^m$  from (a) also induces an action on homology  $H_*((\mathbb{S}^2)^m, \mathbb{C}) \cong \mathbb{C}[\mathfrak{L}_m]$  which can be described explicitly using the line diagram basis. A generator  $s_i$ ,  $i \neq 0$ , acts on a line diagram by permuting the lines  $i$  and  $i + 1$ , and  $s_0$  acts by sending a line diagram to its additive inverse if the first line is undotted and leaves it invariant otherwise. By [SW16, Proposition 40] we have an isomorphism of  $\mathcal{W}_{C_m}$ -modules  $H_{2l}((\mathbb{S}^2)^m, \mathbb{C}) \cong V_{((m-l), (l))}$ . From the explicit descriptions of the  $\mathcal{W}_{C_m}$ -actions it is evident that the map  $H_{2l}((\mathbb{S}^2)^m, \mathbb{C}) \cong H^{2l}(\text{Sk}_{2(m-k)}^m, \mathbb{C})$  sending  $l_U$  to  $X_{i_1} \dots X_{i_l}$ , where  $U = \{i_1, \dots, i_l\}$ , is an isomorphism of  $\mathcal{W}_{C_m}$ -modules and we obtain the isomorphisms as claimed in Proposition 57(c).  $\square$

*Remark 58.* By Theorem 54 we found a topological space isomorphic to the exotic Springer fiber  $\mathcal{F}l^{((k), (m-k))}$  in the homotopy category of topological spaces. In contrast to the exotic Springer fiber itself (and also its homeomorphic topological model from Theorem 23), this homotopy equivalent space has the property that it admits an action of the Weyl group by Proposition 57(a). By combining Proposition 57(c) with results from [Kat17], the induced action in cohomology indeed recovers the original exotic Springer representation [Kat11] and

hence can be used as a toy model for Kato's more involved construction. Moreover, by the proof of Proposition 57 and the discussion in [SW16, §4] we can identify the monomial basis of our presentation of  $H_{2l}(\mathcal{F}l^{((k),(m-k))}, \mathbb{C}) \cong H^{2l}(\mathrm{Sk}_{2(m-k)}^m, \mathbb{C})$  from Theorem 54 as a particularly nice basis since each monomial corresponds to a basis vector of  $V_{((m-k),(k))}$  labeled by a standard bitableau of shape  $((m-k), (k))$  under the isomorphism of Proposition 57, see also [Can96].

## REFERENCES

- [AH08] P. N. Achar and A. Henderson, *Orbit closures in the enhanced nilpotent cone*, Adv. Math. **219** (2008), no. 1, 27–62.
- [AHS11] P. N. Achar, A. Henderson, and E. Sommers, *Pieces of nilpotent cones for classical groups*, Represent. Theory **15** (2011), 584–616.
- [BS11a] J. Brundan and C. Stroppel, *Highest weight categories arising from Khovanov's diagram algebra I: Cellularity*, Mosc. Math. J. **11** (2011), no. 4, 685–722.
- [BS11b] ———, *Highest weight categories arising from Khovanov's diagram algebra III: category  $\mathcal{O}$* , Represent. Theory **15** (2011), 170–243.
- [BS12a] ———, *Gradings on walled Brauer algebras and Khovanov's arc algebra*, Adv. Math. **231** (2012), no. 2, 709–773.
- [BS12b] ———, *Highest weight categories arising from Khovanov's diagram algebra IV: the general linear supergroup*, J. Eur. Math. Soc. **14** (2012), no. 2, 373–419.
- [Can96] H. Can, *Representations of the Generalized Symmetric Groups*, Beiträge Algebra Geom. **37** (1996), no. 2, 289–307.
- [CG97] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser, 1997.
- [CK08] S. Cautis and J. Kamnitzer, *Knot homology via derived categories of coherent sheaves. I. The  $\mathfrak{sl}(2)$ -case*, Duke Math. J. **142** (2008), 511–588.
- [CK14] Y. Chen and M. Khovanov, *An invariant of tangle cobordisms via subquotients of arc rings*, Fund. Math. **225** (2014), no. 1, 23–44.
- [CM93] D.H. Collingwood and W.M. McGovern, *Nilpotent Orbits in Semisimple Lie Algebras*, Mathematics Series, Van Nostrand-Reinhold, 1993.
- [dGN09] J. de Gier and A. Nichols, *The two-boundary Temperley–Lieb algebra*, J. Algebra **321** (2009), 1132–1167.
- [dGNPR05] J. de Gier, A. Nichols, P. Pyatov, and V. Rittenberg, *Magic in the spectra of the XXZ quantum chain with boundaries at  $\Delta = 0$  and  $\Delta = -1/2$* , Nucl. Phys. B **729** (2005), 387–418.
- [dGP04] J. de Gier and P. Pyatov, *Bethe ansatz for the Temperley–Lieb loop model with open boundaries*, J. Stat. Mech. **2004** (2004), P03002.
- [DR18] Z. Daugherty and A. Ram, *Two boundary Hecke algebras and combinatorics of type C*, arXiv:1804.10296, 2018.
- [Ern12] D. C. Ernst, *Diagram calculus for a type affine C Temperley–Lieb algebra, I*, J. Pure Appl. Algebra **216** (2012), 2467–2488.
- [Ern18] ———, *Diagram calculus for a type affine C Temperley–Lieb algebra, II*, J. Pure Appl. Algebra **222** (2018), 3795–3830.
- [ES16] M. Ehrig and C. Stroppel, *2-row Springer fibres and Khovanov diagram algebras for type D*, Canadian J. Math. **68** (2016), no. 6, 1285–1333.
- [FLL<sup>+</sup>18] Z. Fan, C. Lai, Y. Li, L. Luo, W. Wang, and H. Watanabe, *Quantum Schur duality of affine type C with three parameters*, arXiv:1802.01047, 2018.
- [FM10] L. Fresse and A. Melnikov, *On the singularity of the irreducible components of a Springer fiber in  $\mathfrak{sl}_n$* , Selecta Math. **16** (2010), no. 3, 393–418.
- [Fun03] F. Fung, *On the topology of components of some Springer fibers and their relation to Kazhdan–Lusztig theory*, Adv. Math. **178** (2003), no. 2, 244–276.
- [Kat09] S. Kato, *An exotic Deligne–Langlands correspondence for symplectic groups*, Duke Math. J. **148** (2009), no. 2, 305–371.
- [Kat11] ———, *Deformations of nilpotent cones and Springer correspondences*, Amer. J. Math. **133** (2011), 519–553.
- [Kat17] ———, *An algebraic study of extension algebras*, Amer. J. Math. **139** (2017), no. 3, 567–615.
- [Kho02] M. Khovanov, *A functor-valued invariant of tangles*, Alg. Geom. Top. **2** (2002), 665–741.
- [Kho04] ———, *Crossingless matchings and the  $(n, n)$  Springer varieties*, Communications in Contemporary Math. **6** (2004), 561–577.
- [KL87] D. Kazhdan and G. Lusztig, *Proof of the Deligne–Langlands conjecture for Hecke algebras*, Invent. Math. **87** (1987), no. 1, 153–215.
- [Mac95] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, Oxford University Press, 1995.
- [MS16] V. Mazorchuk and C. Stroppel,  *$G(l, k, d)$ -modules via groupoids*, J. Algebraic Combin. **43** (2016), no. 1, 11–32.
- [Nan13] V. Nandakumar, *Equivariant coherent sheaves on the exotic nilpotent cone*, Represent. Theory **17** (2013), 663–681.

- [Nic06a] A. Nichols, *Structure of the two-boundary XXZ model with non-diagonal boundary terms*, J. Stat. Mech. **2006** (2006), L02004.
- [Nic06b] ———, *The Temperley–Lieb algebra and its generalizations in the Potts and XXZ models*, J. Stat. Mech. **2006** (2006), P01003.
- [NRS17] V. Nandakumar, D. Rosso, and N. Saunders, *Irreducible Components of Exotic Springer fibres II: The Exotic Robinson-Schensted Algorithm*, arXiv:1710.08948, 2017.
- [NRS18] ———, *Irreducible components of exotic Springer fibres*, J. Lond. Math. Soc. **98** (2018), no. 3, 609–637.
- [RT11] H.M. Russell and J. Tymoczko, *Springer representations on the Khovanov Springer varieties*, Math. Proc. Cambridge Philos. Soc. **151** (2011), 59–81.
- [Rus11] H.M. Russell, *A topological construction for all two-row Springer varieties*, Pacific J. Math. **253** (2011), 221–255.
- [Sch12] G. Schäfer, *A graphical calculus for 2-block Spaltenstein varieties*, Glasgow Math. J. **54** (2012), 449–477.
- [Ser77] J.-P. Serre, *Linear representations of finite groups*, Graduate Texts in Mathematics, vol. 42, Springer-Verlag, 1977.
- [Spa76] N. Spaltenstein, *The fixed point set of a unipotent transformation on the flag manifold*, Nederl. Akad. Wetensch. Proc. Ser. A, vol. 79, 1976, pp. 452–456.
- [Spa82] ———, *Classes Unipotentes et Sous-groupes de Borel*, Lecture Notes in Mathematics, vol. 946, Springer-Verlag, 1982.
- [Spr76] T. A. Springer, *Trigonometric sums, Green functions of finite groups and representations of Weyl groups*, Invent. Math. **36** (1976), 173–207.
- [Spr78] ———, *A construction of representations of Weyl groups*, Invent. Math. **44** (1978), 279–293.
- [SS14] T. Shoji and K. Sorlin, *Exotic symmetric space over a finite field, II*, Transform. Groups **19** (2014), no. 3, 887–926.
- [Str09] C. Stroppel, *Parabolic category  $\mathcal{O}$ , perverse sheaves on Grassmannians, Springer fibres and Khovanov homology*, Compos. Math. **145** (2009), no. 4, 954–992.
- [SW12] C. Stroppel and B. Webster, *2-block Springer fibers: convolution algebras and coherent sheaves*, Comment. Math. Helv. **87** (2012), 477–520.
- [SW16] C. Stroppel and A. Wilbert, *Two-block Springer fibers of types C and D: a diagrammatic approach to Springer theory*, arXiv:1611.09828, 2016, to appear in Math. Z.
- [Var79] J.A. Vargas, *Fixed points under the action of unipotent elements of  $SL_n$  in the flag variety*, Bol. Soc. Mat. Mexicana **24** (1979), no. 1, 1–14.
- [vL89] M. van Leeuwen, *A Robinson-Schensted algorithm in the geometry of flags for classical groups*, PhD thesis, Rijksuniversiteit Utrecht, 1989.
- [Weh09] S. Wehrli, *A remark on the topology of  $(n, n)$  Springer varieties*, arXiv:0908.2185, 2009.
- [Wil18] A. Wilbert, *Topology of two-row Springer fibers for the even orthogonal and symplectic group*, Trans. Amer. Math. Soc. **370** (2018), 2707–2737.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF GREENWICH, UNITED KINGDOM; AND, HONORARY ASSOCIATE OF THE SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, AUSTRALIA  
*Email address:* n.saunders@greenwich.ac.uk

SCHOOL OF MATHEMATICS & STATISTICS, UNIVERSITY OF MELBOURNE, AUSTRALIA  
*Email address:* arik.wilbert@unimelb.edu.au