

LETTER • OPEN ACCESS

Designing multifunctional landscapes for forest conservation

To cite this article: Truly Santika *et al* 2015 *Environ. Res. Lett.* **10** 114012

View the [article online](#) for updates and enhancements.

Related content

- [Fire emissions and regional air quality impacts from fires in oil palm, timber, and logging concessions in Indonesia](#)
Miriam E Marlier, Ruth S DeFries, Patrick S Kim *et al.*
- [Spatially explicit estimates of forest carbon emissions, mitigation costs and REDD+ opportunities in Indonesia](#)
Victoria Graham, Susan G Laurance, Alana Grech *et al.*
- [The future of forests and orangutans \(*Pongo abelii*\) in Sumatra: predicting impacts of oilpalm plantations, road construction, and mechanisms for reducing carbon emissions from deforestation](#)
David L A Gaveau, Serge Wich, Justin Epting *et al.*

Recent citations

- [Changing landscapes, livelihoods and village welfare in the context of oil palm development](#)
Truly Santika *et al*
- [Does oil palm agriculture help alleviate poverty? A multidimensional counterfactual assessment of oil palm development in Indonesia](#)
Truly Santika *et al*
- [Top 100 research questions for biodiversity conservation in Southeast Asia](#)
J.L. Coleman *et al*



LETTER

Designing multifunctional landscapes for forest conservation

OPEN ACCESS

RECEIVED

25 September 2014

REVISED

29 September 2015

ACCEPTED FOR PUBLICATION

30 September 2015

PUBLISHED

5 November 2015

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Truly Santika^{1,2,3}, Erik Meijaard^{2,3} and Kerrie A Wilson^{1,2}¹ The University of Queensland, School of Biological Sciences, Brisbane, QLD 4072, Australia² ARC Centre of Excellence for Environmental Decisions (CEED), The University of Queensland, Brisbane, QLD 4072, Australia³ Borneo Futures, Country Woods 306, Jl. WR Supratman, Pondok Ranji-Rengas, Ciputat, 15412 Jakarta, IndonesiaE-mail: trulysantika@gmail.com**Keywords:** Borneo, deforestation, logging concessions, multifunctional landscapes, timber plantation concessions, tropical countries, oil-palm concessionsSupplementary material for this article is available [online](#)**Abstract**

A multifunctional landscape approach to forest protection has been advocated for tropical countries. Designing such landscapes necessitates that the role of different land uses in protecting forest be evaluated, along with the spatial interactions between land uses. However, such evaluations have been hindered by a lack of suitable analysis methodologies and data with fine spatial resolution over long time periods. We demonstrate the utility of a matching method with multiple categories to evaluate the role of alternative land uses in protecting forest. We also assessed the impact of land use change trajectories on the rate of deforestation. We employed data from Kalimantan (Indonesian Borneo) at three different time periods during 2000–2012 to illustrate our approach. Four single land uses (protected areas (PA), natural forest logging concessions (LC), timber plantation concessions (TC) and oil-palm plantation concessions (OC)) and two mixed land uses (mixed concessions and the overlap between concessions and PA) were assessed. The rate of deforestation was found to be lowest for PA, followed by LC. Deforestation rates for all land uses tended to be highest for locations that share the characteristics of areas in which TC or OC are located (e.g. degraded areas), suggesting that these areas are inherently more susceptible to deforestation due to foregone opportunities. Our approach provides important insights into how multifunctional landscapes can be designed to enhance the protection of biodiversity.

1. Introduction

Protecting forest and reducing deforestation is central to mitigating the impacts of climate change and averting the loss of biodiversity and the services provided by natural ecosystems [1]. However, for most of the developing tropics, this is challenged by socio-economic pressure to clear land, development agendas arising from government policy, and the opportunity costs of forest protection [2, 3]. Many protected areas (PA) are isolated and situated in a matrix of agricultural land uses, thus reducing their effective size and capacity to maintain the biological diversity that they were originally designated to protect [4]. Therefore, PA alone, despite their growing extent, are unlikely to be sufficient to conserve biodiversity and other ecosystem services [5, 6]. These considerations have

led to a shifted conservation paradigm from a focus solely on PA to the integration of PA within wider multifunctional landscapes [7–9]. Under this new paradigm, PA remain a cornerstone of biodiversity conservation policy, but the landscape matrix also becomes important for achieving broader conservation and development objectives [10, 11]. This approach will likely also increase the robustness and resilience of forest protection strategies, especially under a changing climate [12, 13].

A multifunctional landscape approach to forest protection is cognizant of the interrelationships among different land uses across the landscape matrix [8, 11, 13]. These land use types and spatial locations are determined by planning processes that divide a territory into zones with different rules and regulations, management practices and land covers. The locations

assigned to different land uses often have dominant characteristics. For example, in tropical regions PA have historically been located in remote and mountainous areas [14], natural forest logging concessions (LC) are generally established where timber resources are most abundant and easily extracted, and agricultural plantations are generally established in locations with suitable climate and soil conditions and accessible to human population centers. Land uses can be defined based on their use, i.e. conservation or extraction, or the users, e.g., community forests or concessions [15]. A variety of land use systems exist in tropical forested areas, although LC, agricultural plantation concessions, mining permits, and PA are the principal forms of government-controlled land use [16, 17]. Small-scale agriculture under community control predominates on remaining lands [18].

The shift in the conservation paradigm from a focus on PA to multifunctional landscapes necessitates a comprehensive approach for evaluating the performance of forest protection policies that accounts for the contribution of multiple land uses [19]. However, this presents critical methodological challenges. First, it requires evaluation of the dynamic change in both forest cover and land use configurations through time [20, 21]. The data required to support such analyses are often not available over extended time periods and at broad spatial extents. Second, past evaluations of the effectiveness of PA in abating deforestation rely on reduced-form empirical estimates. These estimates appraise whether or not PA and the surrounding landscape matrix were able to maintain forests based on an aggregated measure of effectiveness (e.g. average deforestation rate), but rarely address why and how different land uses contribute to forest protection [22]. Ideally evaluation methods should seek to identify the underlying causes and mechanisms that determine effectiveness, and appraise the associated interplay among different land uses [23].

With the lack of historical land use change data, past evaluations of the effectiveness of PA and other land uses in abating deforestation have had to rely on static land use maps that are typically based on the most current land use configurations [24–26]. This approach can potentially lead to an incomplete understanding of the effectiveness of different land uses for reducing deforestation. This is particularly the case for the developing tropics, which are characterized by rapid changes in land use and land cover in response to global demand for food, fiber and fuel [27–29], and often facilitated by a combination of decentralization of government authority [30–32] and weak and changing land tenure systems [33]. Land disputes and overlapping permits for forest use are common [34, 35], and can include overlap between LC or agriculture plantation concessions and PA [36, 37]. Ignoring the existence of such overlaps can potentially confound measures of the effectiveness of single land uses. Furthermore, recent studies from Indonesia that were


based on static land use data have shown that the deforestation rate in LC in Indonesia were generally low and comparable to PA [24, 26]. This suggests that selectively harvested LC can potentially provide a secondary source of forest protection [26]. This is contrary to other studies that have found commercial logging to be the main contributor to deforestation [37], initiating the trajectory of forest degradation that eventually leads to conversion of land to agricultural plantations [38–40]. In these earlier studies, however, the impact of timber harvest on deforestation was assessed irrespective of its legality, i.e. whether or not the removal of trees was selectively planned and performed within the boundary of an LC and within cutting limits, or illegally outside of an LC or including trees below a minimum diameter [39]. To date, there has not been a comprehensive evaluation of the contribution of LC to deforestation that explicitly accounts for both the change in land use and forest cover.

In this paper, we assessed the conservation role of different land uses to inform the design of multifunctional landscapes. We also assessed how land use change trajectories affect the pattern of deforestation. To illustrate our approach we used data from Kalimantan, the Indonesian part of Borneo, over three time periods: 2000–2004, 2004–2008, and 2008–2012. We first determined the relationship between the spatial assignments of each land use and a suite of geographical and soil variables. On the basis of these relationships we calculated the likelihood of a grid cell to be assigned to a single land use type. We then applied a matching method adapted for multiple land use categories [41, 42] and the deforestation rate for each land use was compared based on the matched datasets. Our approach offers an improved method for assessing the effectiveness of PA by enabling the contribution of multiple land uses to mitigating deforestation to be evaluated simultaneously and explicitly accounting for the spatial interplay between land uses.

2. Methods

2.1. Study area

Kalimantan covers an area of approximately 533 500 km². Between 2000 and 2012, approximately 17.6% of the total land area was allocated to LC, i.e. parcels of forest leased out to companies to selectively extract timber on a long-term basis. About 13.1% of the total land area was allocated to oil-palm (*Elaeis guineensis*) plantation concessions (OC), and 5.3% was allocated to timber plantation concessions (TC) of fast-growing species, primarily for pulp, paper and rubber production (*Acacia mangium*, *Hevea*, or *Eucalyptus spp.*). About 19.1% of Kalimantan's land area was designated as PA, including strict PA (IUCN PA Category I–III) and district-managed watershed protection forest (*Hutan Lindung*). During this period,



land use change driven by economic and political demands has accelerated the rate of deforestation [38, 43]. Gaveau *et al* [26] found that between 2000 and 2010, 1.2% of the forest area in Kalimantan's PA was deforested, while forests in areas granted to LC were reduced by 1.5% and forests in areas granted to OC were reduced by 14.1%.

2.2. Data and analysis

2.2.1. Land use

Maps of PA, including strict PA and watershed protected forest (*Hutan Lindung*), and concession types, i.e. OC, TC (IUPHHK-HT), and LC (IUPHHK-HA), for 2000–2004, 2004–2008, and 2008–2012 were obtained from the Indonesia's Ministry of Forestry (MoF) and Department of Estate Crops within the Ministry of Agriculture. The assignments of PA and concession types for the three different time periods are shown in figure 1. The study area was assigned a grid of land parcels with a spatial resolution of 1×1 km² and at least one land use was assigned to each grid cell.

2.2.2. Extent of natural forest in 2000 and annual forest loss between 2000 and 2012

The extent of natural forest in 2000 was obtained from Indonesia's MoF [44]. Natural forest constitutes undisturbed old-growth forests (primary forest in the MoF classification) and forests degraded by logging (secondary forest in the MoF classification). We overlaid the natural forest extent with forest cover data for 2000 obtained from the LANDSAT based Global Forest Change dataset [45]. The forest cover data has a spatial resolution of 30×30 m² and comprises continuous values where 0% represents bare land and 100% represents dense forest. Since we were interested in the

loss of intact forest across different land use types, we only included grid cell that were completely covered by natural forest (i.e. 100 hectares) in the beginning of each time period (i.e. 2000, 2004, and 2008). Following Hansen *et al* [45], we used a threshold of 30% to define a pixel as forested in each subsequent time period.

Data on the annual rate of forest loss between 2000 and 2012 were also derived from the Global Forest Change dataset. The annual loss data has a spatial resolution of 30×30 m² and comprises binary values, where '1' denotes clearing of forest within a 30×30 m² grid cell and '0' otherwise. We first overlaid the forest loss data with the aforementioned natural forest extent data to obtain a map of the annual rate of forest loss. We then calculated the total loss of forest in hectares in each 1×1 km² grid cell for each time period $t \in \{2000-2004, 2004-2008, 2008-2012\}$ and this was represented by the variable $DEFOR_t$. Thus, $DEFOR_t$ ranged between 0 and 100, where 0 represents the absence of forest loss and 100 represents that the entire area of forest within the grid cell had been cleared during the four year period.

2.2.3. Spatial assignment of land uses

For each time period t , where $t \in \{2000-2004, 2004-2008, 2008-2012\}$, we accounted for three categories of land use: (1) the dominant single land uses within the region SLU_t (i.e. PA, LC, TC, and OC); (2) mixed land uses MLU_t (mixed concessions where two or more concessions overlap (MC) and areas where either LC, TC, or OC overlapped with PA (PC)), (3) and all other land uses outside the single and the mixed land uses (Other).

The spatial characteristics of different land uses for each time period SLU_t were assessed using a multinomial logistic regression model incorporating a suite

of geographical and soil predictor variables [46]. The geographical predictor variables included the change in the human population density and the governing district [47, 48] and static variables of altitude, slope, travel distance to the nearest city, travel distance to the nearest arterial road and long-term mean annual rainfall (table S1). The soil predictor variables included soil type, drainage, depth, and acidity (table S1). These data are the best available soil data for Kalimantan and have been used in other recent research [49, 50]. The multinomial logistic regression model for each time period was applied using the multinom function in the nnet R-package [51] and linear, quadratic, and cubic functions were tested for each variable. Using the derived relationships for each time period, we predicted the likelihood that a grid cell would be assigned to each single land use (SLU_t) [52]. We assessed the predictive performance of the multinomial logistic models by comparing the predicted land use assignments with the actual land uses, measured using the overall percentage success (OPS) and Gwet's AC1 statistic [53]. Gwet's AC1 statistic modifies the usual Cohen's kappa agreement measure [54] and is more robust to data prevalence [53]. The AC1 value ranges between 0 and 1, where a value larger than 0.6 represents substantial agreement [55].

2.2.4. Deforestation in different land uses

The propensity score of a grid cell was determined by the probability associated with each land use as predicted by the multinomial logistic regression. We then identified grid cells with similar propensity score values to match locations in terms of their geographical and soil characteristics. We applied the nearest neighbor with calipers method to obtain the optimal matched data, by setting a caliper width of 0.2 of the standard deviation of the propensity score's logit function [56]. We used the *nn2* function in the RANN R-package [57] to perform the nearest neighbor search. For each time period, we then compared the average deforestation rate per km^2 across different land uses using the matched datasets. We also calculated the deforestation rate over the entire landscape for each land use category (referred to herein as the 'naïve' approach).

3. Results

3.1. Spatial assignment of land uses

Spatial assignments of land uses in Kalimantan for 2000–2004, 2004–2008, and 2008–2012 are shown in figure 1. PA are predominantly located in remote areas at high altitude, with a small proportion of these areas located in densely populated areas at low altitude (figure S1). Conversely, LC are largely located in areas at moderate altitude and distant from cities and arterial roads (figure S1). TC and OC are predominantly located in lowland areas in close proximity to


cities and arterial roads (figure S1). Moderately shallow (11–50 cm deep), strongly acid, and well to excessively drained soils characterizes soil found inside the boundaries of PA (figure S2). Conversely, well-drained ultisols, inceptisols or alfisols soil types characterize LC. TC have either extremely acid (pH 4.0–4.5) or neutral (pH 6.6–7.3), deep to very deep (75–150 cm) alfisols or oxisols soil types (figure S2). On the other hand, very deep (>100 cm) and extremely acid (pH 4.0–4.5) histosols, oxisols or spodosols are the dominant soil type of OC (figure S2). This suggests that OC have generally been assigned to areas with naturally infertile soil with high soil acidity, whereas TC, have generally been assigned to areas with either naturally fertile or infertile soil [58]. Oil-palm is considered moderately tolerant to a wide range of soil types and acidities, as long as it is well watered [59–61].

Of all $1 \times 1 \text{ km}^2$ grid cells within a single land use category (i.e. PA, LC, TC and OC), at least 73% were correctly predicted as the current land use assignment in each time period (figure S3). Gwet's AC1 for each time period was greater than 0.63 (figure S3), indicating strong agreement between the actual and predicted land uses and also that the land use types have been assigned in a predictable way across the landscape. The proportion of land parcels assigned to LC and OC has increased through time, with the models exhibiting better performance in the later time periods for these land uses (table S2).

3.2. Comparative deforestation rates among land uses

Based on the combined Global Forest Change datasets and the MoF natural forest extent, we estimated that in 2000 the extent of natural forest in Kalimantan was about 300 000 km^2 (table S3). However, by 2012 the extent of natural forest had been reduced to 276 000 km^2 (table S3), equating to 7.9% loss of the 2000 extent, and conforming with the findings of an earlier study [39]. Between 2000 and 2012, natural forest accounted for 81.7% of the area designated as PA and 80.8% of the area designated as LC (or 28.5% and 25.9% of Kalimantan's total natural forest area respectively, table S3). Between 2000 and 2012, 28.5% of the area designated as TC and 24.9% of the area designated as OC comprised natural forest, equating to 2.8% and 6.1% of the total area of natural forests, respectively (table S3).

Based on spatially-matched data, PA had the lowest deforestation rate per km^2 every four years (1.6 hectares on average) followed by LC (2.5 hectares), TC (5.4 hectares), and OC (9.9 hectares) (figure 2(a)). The deforestation rate inside PA where concessions have also been assigned (2.3 hectares) was higher than inside PA without overlapping concessions. In comparison, using the naive approach, the rate of deforestation for each land use was predicted to be lower for PA and LC, but higher for TC and OC. Between 2000



and 2012, the deforestation rate every four years per km² for PA was predicted to be 0.2 hectares on average, for LC the rate was 0.8 hectares, for TC the rate was 6.0 hectares, and for OC the rate was 12.0 hectares (figure 2(b)). Inside PA where concessions have also been assigned, the deforestation rate was 0.7 hectares.

The spatially-matched data reveals a shift in deforestation trends through time (figure 2(a)). Between 2000 and 2004 the deforestation rate was greatest in mixed concessions (4.4 hectares), TC (3.6 hectares), and OC (3.0 hectares). Between 2004 and 2008, the deforestation rate was greatest inside OC (7.2 hectares) and mixed concessions (5.2 hectares). Between 2008 and 2012 the deforestation rate was greatest inside OC (19.5 hectares). This suggests that deforestation in early 2000 was motivated mainly in response to demand for timber production, whereas it is now largely motivated by oil-palm establishment. Additionally, the deforestation rate has increased through time, particularly for OC, suggesting that there is a growing pressure to clear forest in this region.

For all land uses, we found that the deforestation rate for each time period was greater in locations that share characteristics typically associated with TC or OC (figure 3). For example, inside PA, in the locations that share characteristics typically associated with plantation concessions the deforestation rate per km² every four years was 3.1 hectares on average, whereas for other locations that do not share these characteristics the rate was 0.2 hectares (figure 3).

3.3. Land use change trajectories and deforestation of intact forest

Since 2000, Kalimantan has undergone a rapid expansion of concessions as well as complex alteration of land uses. In 2000, about 48.2% of the total land area had not been assigned to any concession or PA, or was assigned to the 'Other' land use category (table S3). By 2008, 14.2%, 9.2% and 1.4% of this concession-free area were granted to OC, LC and TC, respectively

(table S4(a)). The expansion of concessions continued and by 2012 about 11.9% of the concession-free area was converted to LC, TC or OC (table S4(b)). Between 2000 and 2012, some of the existing LC and TC were either converted to OC or the concession was removed. For example, about 2.8% of LC that existed in 2000 were converted to OC by 2008 (table S4(a)). At the same time, 18.0% of LC and 3.9% of TC lost their concession permit (table S4(a)). By 2012, 5.3% of LC were converted to OC (table S4(b)).

Of all locations currently assigned to OC (i.e. assigned to this land use between 2008 and 2012), 29.1% comprised intact forest in 2000 (table S5). Almost a third of this intact forest was assigned to OC as early as 2000, while a little over a half was without a concession (or concession-free) and a small proportion (3.4%) was originally allocated to LC. When the concession-free areas and the LC were later converted to OC, about 84.3% comprised intact forest (with >90% forest cover) (figure S4(a)). Similarly, of all locations currently assigned to TC (i.e. assigned to this land use between 2008 and 2012), 34.0% comprised intact forest in 2000 (table S5). About 46.3% of this intact forest was assigned TC to as early as 2000, while 53.2% were concession-free. When the concession-free area were later converted to TC, about 82.3% was intact prior to conversion (figure S4(a)). This suggests forest degradation was unlikely to be the main reason for the conversion of concession-free areas or LC to TC or OC during 2000–2012. However, among LC that were later converted to OC, the proportion of areas with excessively acid soil ($\text{pH} \leq 4.0$) was high (32%) (figure S4(b)). High soil acidity ($\text{pH} \leq 4.5$) was also found in a relatively large proportion of concession-free areas that were later converted to TC or OC (figure S4(b)). This suggests that conversion from natural forest LC or areas without concessions to plantation concessions were more likely to be motivated by the perceived inherent land capability rather than forest degradation.




Figure 3. The deforestation rate per km² for grid cells that were predicted to be other land uses based on matched data for three time periods (2000–2004, 2004–2008, and 2008–2012).

4. Discussion

In recent years, there has been an increasing awareness of the importance of multifunctional landscapes for conserving biodiversity [62]. This necessitates that evaluations of the performance of conservation policy shift from focusing on the effectiveness of PA alone (at a site level) to appraising the role of all land uses (at a landscape scale) [26]. Aggregated measures of the effectiveness of different land uses for mitigating deforestation [22], such as the overall deforestation rate, fail to reveal the underlying mechanisms of deforestation within a land use [23]. While the deforestation rate for PA and LC revealed by our analysis reflects the overall deforestation rate found by an earlier study [26], the vulnerability to deforestation varies between land uses and locations, particularly in relation to potential foregone opportunities for agriculture or timber extraction (figure 3). For example, while the deforestation rate for LC with a high potential for agriculture or plantation establishment during 2000–2010 was about 6.3 hectares per km², the deforestation rate inside PA that shared similar characteristics was half this rate (3.1 hectares per km²). This indicates that PA are mitigating some of the impacts of deforestation in this region.

TC and OC have tended to be located in areas in close proximity to cities. However, these concessions are also located in areas with naturally infertile soils, such as on peat swamps and dry lands with acidic soil. These marginal areas [63] are usually unoccupied as a result of their low value for small-holder agriculture [64]. The reclamation of peat swamps or dry land is a lucrative alternative to transporting crops from remote areas, especially if the cost of reclamation can

be subsidized from the sale of timber by clearing forest [65]. The vulnerability of such areas to deforestation, therefore, highlights the importance of appropriate management interventions, such as policies that increase the value of forest (e.g. through biodiversity and ecosystem service payments) [66] as well as enforcement of environmental regulations.

Logging activities have been viewed as the initial step in the trajectory towards forest degradation that ultimately leads to oil-palm plantation establishment [67], although we found evidence to the contrary. Based on a detailed analysis of land use change trajectories during 2000–2012, we found that among sites currently assigned as TC or OC, nearly a third comprised intact forest in early 2000, confirming an earlier report focused on forest conversion [68]. About half of the concessions assigned within this intact forest were already granted TC or OC in early 2000 and the other half were either without a concession (or concession-free) or assigned as LC. However, when these concession-free areas or LC were later converted to TC or OC in the mid to late 2000s, over 80% had >90% forest cover. This suggests that forest degradation was unlikely to be the main reason for conversion of LC or concession-free areas to plantation concessions, at least during 2000–2012. Instead, we found a large proportion of these areas were located on highly acid soils, suggesting that perceived land capability was more likely to be the key reason for conversion. This finding is consistent with the fact that the term of ‘degraded land’ is used by the Indonesian government to represent a wide variety of land conditions, including degraded forest (areas that have been severely logged [49]), critical lands (areas that have been subject to intensive agricultural practices) and marginal lands (areas

considered to be unproductive with high soil acidity, including peat swamp and dry land on acid soil) [63].

As anticipated, the deforestation rate within TC and OC was two to four times higher than the rate in LC. Our analysis also reveals that the deforestation rate within OC has increased through time. Interestingly, the deforestation rate in LC was lower than areas without a concession, revealing the impacts of illegal logging activities. The capacity for LC to reduce the rate of deforestation compared to both plantation concessions and concession-free areas in Kalimantan supports recommendations for reclassifying LC as PA under the IUCN Protected Area Category VI [26]. In addition, such reclassification could lessen the impacts of current government policy that results in the conversion of LC to OC, particularly in areas perceived to be unproductive or marginal. Instead the rehabilitation and restoration of these areas should be encouraged [69]. Such reclassification could also benefit biodiversity conservation, with many endangered species residing inside LC [70].

Our analysis has revealed locations where multiple types of concessions overlap and locations where concessions overlap with PA. In the latter, the deforestation rate was higher than in PA as a single land use type (figure 2). Therefore, this underlines the value of accounting for overlapping land use allocations when assessing the effectiveness of PA. Overlapping land uses are not unique to Indonesia and have been reported in countries in South America and Africa [32, 71, 72]. In Indonesia, overlapping land uses not only occur due to weak land use planning systems, but also as a consequence of the discrepancy in maps developed at different levels of government [73]. This is particularly evident for the province of Central Kalimantan where PA boundaries considered by the local government differ from the ones assigned by the Ministry of Environment and Forestry [74]. In an attempt to reconcile these conflicting maps, the Indonesian government has recently launched the 'One Map' initiative that aims to resolve land disputes and overlapping concession permits [75].

Our study advances past evaluations of the effectiveness of PA by accounting for the spatial interplay between alternative land uses and explicitly accounting for the dynamic change in land use through time. Our findings suggest that a mix of different forest conservation strategies is needed in Kalimantan that integrates different levels of protection and resource use by local communities [76]. Formally reclassifying LC as PA under the IUCN Protected Area Category VI could enhance biodiversity protection, increase social welfare, and provide economic opportunities for local people and businesses. Furthermore, combining PA with LC when planning for multi-functional landscapes could ensure the protection of larger areas of intact forest than possible via PA alone.

Acknowledgments

We thank Oriana Pauli and Rebecca Runting for assistance with the data and anonymous referees for comments. This study is part of 'The Borneo Futures' initiative and was supported by the Australian Research Council (Centre of Excellence and Future Fellowship programs) and The Arcus Foundation.

References

- [1] Canadell J G and Raupach M R 2008 Managing forests for climate change mitigation *Science* **320** 1456–7
- [2] Oliveira P J, Asner G P, Knapp D E, Almeyda A, Galván-Gildemeister R, Keene S, Raybin R F and Smith R C 2007 Land use allocation protects the Peruvian Amazon *Science* **317** 1233–6
- [3] Lele S, Wilshusen P, Brockington D, Seidler R and Bawa K 2010 Beyond exclusion: alternative approaches to biodiversity conservation in the developing tropics *Curr. Opin. Environ. Sustainability* **2** 94–100
- [4] Nagendra H, Paul S, Pareeth S and Dutt S 2009 Landscapes of protection: forest change and fragmentation in Northern West Bengal, India *Environ. Manage.* **44** 853–64
- [5] Tilman D, Cassman K G, Matson P A, Naylor R and Polasky S 2002 Agricultural sustainability and intensive production practices *Nature* **418** 671–7
- [6] Beier P and Brost B 2010 Use of land facets to plan for climate change: conserving the arenas, not the actors *Conserv. Biol.* **24** 701–10
- [7] Pfaff A, Robalino J, Lima E, Sandoval C and Herrera L D 2014 Governance, location and avoided deforestation from protected areas: greater restrictions can have lower impact, due to differences in location *World Dev.* **55** 7–20
- [8] Zuidema P A and Sayer J A 2003 Tropical forests in multi-functional landscapes: the need for new approaches to conservation and research *Tropical Forests in Multi-functional Landscapes* (Utrecht, The Netherlands: Prince Bernhard Centre for International Nature Conservation, Utrecht University) pp 9–19
- [9] Sayer J, Ghazoul J, Nelson P and Boedhihartono A K 2012 Oil palm expansion transforms tropical landscapes and livelihoods *Global Food Secur.* **1** 114–9
- [10] DeFries R, Hansen A, Newton A C and Hansen M C 2005 Increasing isolation of protected areas in tropical forests over the past twenty years *Ecol. Appl.* **15** 19–26
- [11] Lindenmayer D B and Franklin J F 2002 *Conserving Forest Biodiversity: A Comprehensive Multiscaled Approach* (London: Island Press)
- [12] Dawson I K et al 2011 Climate change and tree genetic resource management: maintaining and enhancing the productivity and value of smallholder tropical agroforestry landscapes: a review *Agroforestry Syst.* **81** 67–78
- [13] van Noordwijk M, Hoang M, Neufeldt H, Öborn I and Yatich T 2011 *How Trees and People Can Co-Adapt to Climate Change: Reducing Vulnerability Through Multifunctional Agroforestry Landscapes* (Nairobi, Kenya: World Agroforestry Centre (ICRAF))
- [14] Joppa L N and Pfaff A 2009 High and far: biases in the location of protected areas *PLoS One* **4** e8273
- [15] Lambin E F et al 2014 Effectiveness and synergies of policy instruments for land use governance in tropical regions *Glob. Environ. Chang.* **28** 129–40
- [16] Nasi R, Billand A and van Vliet N 2012 Managing for timber and biodiversity in the Congo Basin *Forest Ecol. Manage.* **268** 103–11
- [17] Agrawal A, Chhatre A and Hardin R 2008 Changing governance of the world's forests *Science* **320** 1460–2
- [18] White A and Martin A 2002 *Who Owns the World's Forests? Forest Tenure and Public Forests in Transition* (Washington, DC: Forest Trends Center for International Environmental Law)

- [19] Sayer J et al 2013 Ten principles for a landscape approach to reconciling agriculture, conservation, and other competing land uses *Proc. Natl Acad. Sci. USA* **110** 8349–56
- [20] Hett C, Castella J C, Heinimann A, Messerli P and Pfund J L 2012 A landscape mosaics approach for characterizing swidden systems from a REDD + perspective *Appl. Geogr.* **32** 608–18
- [21] Dewi S, van Noordwijk M, Ekadinata A and Pfund J L 2013 Protected areas within multifunctional landscapes: squeezing out intermediate land use intensities in the tropics? *Land Use Policy* **30** 38–56
- [22] Miteva D A, Pattanayak S K and Ferraro P J 2012 Evaluation of biodiversity policy instruments: What works and what doesn't? *Oxford Rev. Econ. Pol.* **28** 69–92
- [23] White H 2009 Theory-based impact evaluation: principles and practice *J. Dev. Eff.* **1** 271–84
- [24] Abood S A, Lee J S H, Burivalova Z, Garcia-Ulloa J and Koh L P 2015 Relative contributions of the logging, fiber, oil palm, and mining industries to forest loss in Indonesia *Conserv. Lett.* **8** 58–67
- [25] Brun C, Cook A R, Lee J S H, Wich S A, Koh L P and Carrasco L R 2015 Analysis of deforestation and protected area effectiveness in Indonesia: a comparison of Bayesian spatial models *Glob. Environ. Change* **31** 285–95
- [26] Gaveau D L A et al 2013 Reconciling forest conservation and logging in Indonesian Borneo *PLoS One* **8** e69887
- [27] Pfaff A, Santiago-Avila F, Carnovale M and Joppa L 2014 Protected areas' impacts upon land cover within Mexico: the need to add politics and dynamics to static land use economics *Agricultural and Applied Economics Association Annual Meeting (Minneapolis, Minnesota, 27–29 July 2014)*
- [28] Lambin E F and Meyfroidt P 2011 Global land use change, economic globalization, and the looming land scarcity *Proc. Natl Acad. Sci. USA* **108** 3465–72
- [29] Gibbs H K, Ruesch A, Achard F, Clayton M, Holmgren P, Ramankutty N and Foley J A 2010 Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s *Proc. Natl Acad. Sci. USA* **107** 16732–7
- [30] Ribot J C, Agrawal A and Larson A M 2006 Recentralizing while decentralizing: How national governments reappropriate forest resources *World Dev.* **34** 1864–86
- [31] Tacconi L 2007 Decentralization, forests and livelihoods: theory and narrative *Glob. Environ. Change* **17** 338–48
- [32] Andersson K and Gibson C C 2007 Decentralized governance and environmental change: local institutional moderation of deforestation in Bolivia *J. Policy Anal. Manag.* **26** 99–123
- [33] Schaltegger S, Burritt R, Willem van Gelder J, German L and Bailis R 2012 Biofuels investments in tropical forest-rich countries: implications for responsible finance *Sustainability Account. Manag. Policy J.* **3** 134–60
- [34] Engel S and Palmer C 2006 Who owns the right? The determinants of community benefits from logging in Indonesia *Forest Policy Econ.* **8** 434–46
- [35] Larson A M et al 2013 Land tenure and REDD+: the good, the bad and the ugly *Glob. Environ. Change* **23** 678–89
- [36] Jepson P, Jarvie J K, MacKinnon K and Monk K A 2001 The end for Indonesia's lowland forests? *Science* **292** 859–61
- [37] Curran L M, Trigg S N, McDonald A K, Astiani D, Hardiono Y M, Siregar P, Caniango I and Kasischke E 2004 Lowland forest loss in protected areas of Indonesian Borneo *Science* **303** 1000–3
- [38] Gaveau D L et al 2014 Four decades of forest persistence, clearance and logging on Borneo *PLoS One* **9** e101654
- [39] Margono B A, Potapov P V, Turubanova S, Stolle F and Hansen M C 2014 Primary forest cover loss in Indonesia over 2000–2012 *Nat. Clim. Change* **4** 730–5
- [40] Margono B A, Turubanova S, Zhuravleva I, Potapov P, Tyukavina A, Baccini A, Goetz S and Hansen M C 2012 Mapping and monitoring deforestation and forest degradation in Sumatra (Indonesia) using Landsat time series data sets from 1990 to 2010 *Environ. Res. Lett.* **7** 034010
- [41] Joffe M M and Rosenbaum P R 1999 Invited commentary: propensity scores *Am. J. Epidemiol.* **150** 327–33
- [42] Imbens G W 2000 The role of the propensity score in estimating dose-response functions *Biometrika* **87** 706–10
- [43] Stibig H J, Achard F, Carboni S, Raši R and Miittinen J 2014 Change in tropical forest cover of Southeast Asia from 1990 to 2010 *Biogeosciences* **11** 247–58
- [44] Ministry of Forestry 2012 *National Forest Monitoring System* Jakarta Indonesia
- [45] Hansen M C et al 2013 High-resolution global maps of 21st-century forest cover change *Science* **342** 850–3
- [46] Hosmer J D W, Lemeshow S and Sturdivant R X 2013 *Applied Logistic Regression* (New York: Wiley)
- [47] Linkie M, Sloan S, Kasia R, Kiswayadi D and Azmi W 2014 Breaking the vicious circle of illegal logging in Indonesia *Conserv. Biol.* **28** 1023–33
- [48] Resosudarmo I A P 2004 Closer to people and trees: Will decentralisation work for the people and the forests of Indonesia? *Eur. J. Dev. Res.* **16** 110–32
- [49] Smit H H, Meijaard E, van der Laan C, Mantel S, Budiman A and Verweij P 2013 Breaking the link between environmental degradation and oil palm expansion: a method for enabling sustainable oil palm expansion *PLoS One* **8** e68610
- [50] Fairhurst T and McLaughlin D 2009 *Sustainable Oil Palm Development on Degraded Land in Kalimantan* (Washington, USA: World Wildlife Fund)
- [51] Ripley B 2015 nnet: Feed-forward Neural Networks and Multinomial Log-Linear Models R package version 7.3–11
- [52] Rosenbaum P R and Rubin D B 1983 The central role of the propensity score in observational studies for causal effects *Biometrika* **70** 41–55
- [53] Gwet K 2002 Kappa statistic is not satisfactory for assessing the extent of agreement between raters *Stat. Methods Inter-rater Reliab. Assess.* **1** 1–6
- [54] Cohen J 1968 Weighted kappa: nominal scale agreement provision for scaled disagreement or partial credit *Psychol. Bull.* **70** 213–20
- [55] Landis J R and Koch G G 1977 The measurement of observer agreement for categorical data *Biometrics* **33** 159–74
- [56] Wang Y, Cai H, Li C, Jiang Z, Wang L, Song J and Xia J 2013 Optimal caliper width for propensity score matching of three treatment groups: a Monte Carlo study *PLoS One* **8** e81045
- [57] Arya S, Mount D, Kemp S E and Jefferis G 2015 RANN: Fast Nearest Neighbour Search (Wraps Arya and Mount's ANN Library) R package version 2.5
- [58] Vitousek P M and Sanford R 1986 Nutrient cycling in moist tropical forest *Annu. Rev. Ecol. Evol. Syst.* **17** 137–67
- [59] Corley R H V and Tinker P 2008 *The Oil Palm* (New York: Wiley)
- [60] Nelson P, Rhebergen T, Berthelsen S, Webb M, Banabas M, Oberthur T, Donough C R, Rahmadsyah, Indrasuara K and Lubis A 2011 Soil acidification under oil palm: rates and effects on yield *Better Crops Plant Food* **95** 22–5
- [61] Auxtero E A and Shamshuddin J 1991 Growth of oil palm (*Elaeis guineensis*) seedlings on acid sulfate soils as affected by water regime and aluminium *Plant Soil* **137** 243–57
- [62] Bray D B, Duran E, Ramos V H, Mas J F, Velazquez A, McNab R B, Barry D and Radachowsky J 2008 Tropical deforestation, community forests, and protected areas in the Maya forest *Ecol. Soc.* **13** 56
- [63] Suprpto A 2002 Land and water resources development *Indonesia Investment in Land and Water* (Bangkok: Food and Agriculture Organization of the United Nations (FAO))
- [64] Tan A K 2015 Can't we even share our maps? Cooperative and unilateral mechanisms to combat forest fires and transboundary 'haze' in Southeast Asia *Transboundary Pollution: Evolving Issues of International Law and Policy* ed S Jayakumar et al (Singapore: NUS Centre for International Law) pp 327–61
- [65] Dillon H S, Laan T and Dillon H S 2008 *Biofuels, at What Cost? Government Support for Ethanol and Biodiesel in Indonesia* (Winnipeg Canada: International Institute for Sustainable Development)
- [66] van Noordwijk M, Leimona B, Jindal R, Villamor G B, Vardhan M, Namirembe S, Catacutan D, Kerr J,

- Minang P A and Tomich T P 2012 Payments for environmental services: evolution toward efficient and fair incentives for multifunctional landscapes *Annu. Rev. Env. Resour.* **37** 389–420
- [67] Carlson K M, Curran L M, Ratnasari D, Pittman A M, Soares-Filho B S, Asner G P, Trigg S N, Gaveau D A, Lawrence D and Rodrigues H O 2012 Committed carbon emissions, deforestation, and community land conversion from oil palm plantation expansion in West Kalimantan, Indonesia *Proc. Natl Acad. Sci. USA* **109** 7559–64
- [68] Fairhurst T, McLeish M and Prasodjo R 2010 *Conditions Required by the Private Sector for Oil Palm Expansion on Degraded Land in Indonesia* (Wye: Tropical Crop Consultants Limited)
- [69] Edwards D P, Tobias J A, Sheil D, Meijaard E and Laurance W F 2014 Maintaining ecosystem function and services in logged tropical forests *Trends Ecol. Evol.* **9** 511–20
- [70] Wich S A et al 2012 Understanding the impacts of land use policies on a threatened species: Is there a future for the Bornean orang-utan? *PLoS One* **7** e49142
- [71] Carodenuto S and Cerutti P O 2014 Forest law enforcement, governance and trade (FLEGT) in Cameroon: perceived private sector benefits from VPA implementation *Forest Policy Econ.* **48** 55–62
- [72] Osti M, Coad L, Fisher J B, Bomhard B and Hutton J M 2011 Oil and gas development in the World Heritage and wider protected area network in sub-Saharan Africa *Biodivers. Conserv.* **20** 1863–77
- [73] Sunderlin W D et al 2014 The challenge of establishing REDD + on the ground: insights from 23 subnational initiatives in six countries *CIFOR Occasional paper 104* p 44
- [74] Caldecott J, Indrawan M, Rinne P and Halonen M 2011 Indonesia–Norway REDD + Partnership: first evaluation of deliverables *Final Report* (Jakarta: Gaia Consulting)
- [75] Samadhi N 2013 Indonesia one map: assuring better delivery of national development goals *Geospatial World Forum* pp 12–3
- [76] Abram N K, Meijaard E, Ancrenaz M, Runting R K, Wells J A, Gaveau D, Pellier A S and Mengersen K 2014 Spatially explicit perceptions of ecosystem services and land cover change in forested regions of Borneo *Ecosystem Serv.* **7** 116–27