

# Cordycepin, a metabolite of *Cordyceps militaris*, reduces immune-related gene expression in insects

Victoria C. Woolley<sup>1</sup>, Graham R. Teakle<sup>1</sup>, Gillian Prince<sup>1</sup>, Cornelia H. de Moor<sup>2</sup>, David Chandler<sup>1</sup>

<sup>1</sup>Warwick Crop Centre, School of Life Sciences, University of Warwick, Wellesbourne, Warwick, CV35 9EF UK

<sup>2</sup>School of Pharmacy, University of Nottingham, University Park, Nottingham, NG7 2RD UK

## Abstract

Hypocrealean entomopathogenic fungi (EPF) (Sordariomycetes, Ascomycota) are natural regulators of insect populations in terrestrial environments. Their obligately-killing life-cycle means that there is likely to be strong selection pressure for traits that allow them to evade the effects of the host immune system. In this study, we quantified the effects of cordycepin (3'-deoxyadenosine), a secondary metabolite produced by *Cordyceps militaris* (Hypocreales, Cordycipitaceae), on insect susceptibility to EPF infection and on insect immune gene expression. Application of the immune stimulant curdlan (20 µg ml<sup>-1</sup>, linear beta-1,3-glucan, a constituent of fungal cell walls) to *Drosophila melanogaster* S2r+ cells resulted in a significant increase in the expression of the immune effector gene *metchnikowin* compared to a DMSO-only control, but there was no significant increase when curdlan was co-applied with 25 µg ml<sup>-1</sup> cordycepin dissolved in DMSO. Injection of cordycepin into larvae of *Galleria mellonella* (Lepidoptera: Pyralidae) resulted in dose-dependent mortality (LC<sub>50</sub> of cordycepin = 2.1 mg per insect 6 days after treatment). Incubating conidia of *C. militaris* and *Beauveria bassiana* (Hypocreales, Cordycipitaceae; an EPF that does not synthesize cordycepin) with 3.0 mg ml<sup>-1</sup> cordycepin had no effect on the numbers of conidia germinating *in vitro*. Co-injection of *G. mellonella* with a low concentration of cordycepin (3.0 mg ml<sup>-1</sup>) plus 10 or 100 conidia per insect of *C. militaris* or *B. bassiana* caused a significant decrease in insect median survival time compared to injection with the EPF on their own. Analysis of predicted vs. observed mortalities indicated a synergistic interaction between cordycepin and the EPF. The injection of *C. militaris* and *B. bassiana* into *G. mellonella* resulted in increased expression of the insect immune effector genes *lysozyme*, *IMPI* and *gallerimycin* at 72 h post injection, but this did not occur when the EPF were co-injected with 3.0 mg ml<sup>-1</sup> cordycepin. In addition, we observed

increased expression of *IMPI* and *lysozyme* at 48 h after injection with *C. militaris*, *B. bassiana* and sham injection (indicating a wounding response), but this was also prevented by application of cordycepin. These results suggest that cordycepin has potential to act as a suppressor of the immune response during fungal infection of insect hosts.

## 1. Introduction

In order to successfully grow and reproduce in its host, a pathogen must evade the effects of the host immune system. This can be done through passive mechanisms (e.g. avoiding immune detection) or by actively interfering with host immune responses (Schmid-Hempel, 2009). Understanding the strategies used by pathogens for immune evasion provides valuable insights into the evolution of virulence in host-parasite relationships (Schmid-Hempel, 2009) but our present understanding of immune evasion in insect pathogens is not well developed.

Hypocrealean entomopathogenic fungi (EPF) (Sordariomycetes, Ascomycota) are obligately-killing parasites that infect a wide range of insects and are common in terrestrial ecosystems (Vega et al., 2009). Infection occurs by percutaneous growth of fungal spores into the haemocoel, from where the fungus starts to proliferate and consume host tissues, resulting in insect death (Chandler, 2016). In laboratory assays insect death usually occurs 3-7 days after infection, however this can take longer in a natural situation and varies between EPF species (Chandler, 2016). If environmental conditions are favourable, the fungus grows back out of the insect cadaver to produce ascospores from fruiting bodies (for sexually reproducing forms) or vegetative conidia (for asexual reproduction), which are released into the environment to be acquired by naïve hosts. This particular life cycle, with transmission being dependent upon the pathogen killing its host, imposes a strong selection pressure for traits that confer high virulence in order to maximise pathogen fitness (Roy et al., 2006; Boomsma et al., 2014). Hypocrealean EPF produce a range of secondary metabolites during infectious growth (de Bekker et al., 2013), and there is experimental evidence that some of these disrupt cellular and systemic immune defences, such as the destruxins, a group of cyclic hexadepsipeptides produced by *Metarhizium* species (Vilcinskis et al., 1997a, 1997b; Pal et al., 2007; Zimmermann, 2007a, 2007b). Indirect evidence on the anti-immune function of some other EPF metabolites comes from their use as immune suppressors in human medicine. Examples include cyclosporine, which is used to treat autoimmune diseases and to prevent rejection in organ transplantation, and which is a natural product from *Tolyocladium inflatum*

(Hypocreales, Ophiocordycipitaceae) (Survase et al., 2011), as well as the immune-modulating drug fingolimod, used in the treatment of multiple sclerosis, and which is a synthetic derivative of myriocin, produced by *Cordyceps / Isaria cicadae* s.l. (previously known as *Isaria sinclairii*, currently undergoing taxonomic revision; Hypocreales, Clavicipitaceae) (Fujita et al., 1994; Strader et al., 2011; Kepler et al., 2017). The homologies between the innate immune systems of mammals and insects (Stokes et al., 2015) probably explains why immunosuppressing metabolites from EPF are also able to affect the human immune system. The innate immune systems are sufficiently similar that insects are also being used increasingly as *in vivo* models of pathogen virulence in medical research (Sheehan et al., 2018).

Here, we present the results of a study on insect immune interference by cordycepin (3'-deoxyadenosine), a secondary metabolite produced by the teleomorphic hypocrealean EPF *Cordyceps militaris* (Hypocreales, Cordycipitaceae) (Cunningham et al., 1950; Shih et al., 2007; Tuli et al., 2014). This fungus occurs naturally in temperate and sub-tropical areas within Asia, Europe, and North and South America, although it is considered to be rare (Shrestha et al., 2012). It is a host specialist, causing natural infections predominantly in lepidopteran larvae and pupae (Shrestha et al., 2012). Cordycepin has been investigated previously for its pharmacological potential, particularly in connection with the use of *C. militaris* fruiting bodies as a traditional herbal medicine (Paterson, 2008). It has been reported to have anti-inflammatory (Kim et al., 2006; Jeong et al., 2010; Kondrashov et al., 2012; Zhang et al., 2014; Yang et al., 2015; Ashraf et al., 2019) as well as anti-tumour (Nakamura et al., 2006) and anti-angiogenic properties (Lu et al., 2014) in studies with mammalian cells, but little is known about its effect on insects. For the present study, we addressed the topic using *Drosophila melanogaster* (Diptera: Drosophilidae) cell culture and *Galleria mellonella* (Lepidoptera: Pyralidae) live caterpillars as experimental systems (Lemaitre et al., 1995; Mylonakis et al., 2005; Bergin et al., 2006; O'Neill, 2006; Ramarao et al., 2012; Harding et al., 2013). Firstly, we quantified the effect of cordycepin on the expression of immunity-related genes in a *D. melanogaster* S2r+ cell line treated with activators of the immune response. Secondly, we measured the survival of *G. mellonella* larvae, together with the expression of immunity-related genes, following injection with cordycepin on its own and in combination with *C. militaris* and *Beauveria bassiana* (Hypocreales, Cordycipitaceae), a related, asexually-reproducing EPF species that does not synthesize cordycepin (Xia et al., 2017). The aim was to provide new information on the effect of cordycepin on insect immune gene expression under the controlled

conditions achievable using a cell-based assay, as well as the more biologically-complex conditions of a whole-animal environment.

## **2. Materials and Methods**

### *2.1 Experimental material*

*D. melanogaster* S2r+ cells (Schneider, 1972; Rämét et al., 2001) were cultured in 15 ml cell culture flasks (VWR, Radnor, USA) with 12 ml of Schneider Drosophila Insect Medium (Thermo Fisher Scientific, Waltham, USA) supplemented with 10% foetal bovine serum (FBS; Sigma-Aldrich) and 10% penicillin-streptomycin (Sigma-Aldrich, St Louis, USA). Cell density was determined by trypan blue staining (Strober, 2001). Cells were subcultured when their density reached  $1 \times 10^7$  cells ml<sup>-1</sup>. All cells used in experiments were between passages 5-15.

Final instar, non-diapausing *G. mellonella* larvae were purchased from Wazp Brand UK Ltd. (Yorkshire, UK). They were maintained at 15°C in darkness and larvae between 200-300 mg were selected for experiments. Cordycepin (Sigma-Aldrich, St Louis, USA) was prepared as a stock solution (100 mg ml<sup>-1</sup> in DMSO; Thermo Fisher Scientific, Waltham, USA) and diluted with sterile 0.01% Triton X-100 (Merck, Darmstadt, Germany) wetting agent in DEPC-treated water plus 0.1 mg ml<sup>-1</sup> pentostatin (2'-deoxycoformycin ; Sigma-Aldrich, St Louis, USA) (Johnston, 2011).

Laboratory bioassays with entomopathogenic fungi were done using *B. bassiana* strain 433.99 and *C. militaris* strain ARSEF 11703 (Supplementary Table 1). Stock cultures of the strains were stored in cryopreservation (Chandler, 1994). Laboratory cultures were grown from stock cultures on Sabouraud Dextrose Agar (SDA; Thermo Fisher Scientific, Waltham, USA) slopes and maintained at 5°C for up to six months. Subcultures for laboratory experiments were grown on SDA from slope cultures and incubated in darkness at 23°C for 10-14 days. Conidia were harvested in sterile 0.01% Triton X-100 and suspensions filtered through milk filters (Lantor Ltd, Bolton, UK) to remove hyphal fragments. Conidia were counted using an Improved Neubauer haemocytometer and aliquots prepared at different concentrations as required.

## 2.2 Effect of cordycepin treatment on the response of *D. melanogaster* S2r+ cells to simulated immune challenge

*D. melanogaster* S2r+ cells were cultured as described above. Cells were collected by centrifugation at 1500 x g for 5 min at 4°C, the supernatant removed and cells resuspended in Schneider Drosophila Insect Medium with 10% FBS. Aliquots (3 ml) were pipetted into each well of a 6-well plate (TPP tissue culture plate, Sigma-Aldrich, St Louis, USA) at a density of approximately  $8 \times 10^5$  cells ml<sup>-1</sup>. To determine the effect of cordycepin on gene expression, cells were treated with the immune response stimulants curdlan (linear beta-1,3-glucan; 20 µg ml<sup>-1</sup>; Carbosynth Ltd., Compton, UK) or crude LPS (lipopolysaccharide; 20 µg ml<sup>-1</sup>; Sigma-Aldrich, St Louis, USA), followed by addition of 25 µg ml<sup>-1</sup> (100 µM) cordycepin solution in DMSO, or DMSO only. An untreated control was also included. Curdlan is structurally similar to fungal cell wall polysaccharides and is used to stimulate an anti-fungal immune response (Kumar et al., 2009), whereas crude LPS (which also contains peptidoglycan) is found in the outer membrane of gram-negative bacteria and is used to stimulate an anti-(gram-negative) bacterial immune response (Tanji and Ip, 2005). Cells were incubated for 4 h at 25°C, then RNA was extracted ahead of RT-qPCR (see below).

## 2.3 Effect of cordycepin treatment on the survival of *G. mellonella* larvae infected with the EPF *C. militaris* and *B. bassiana*

### 2.3.1 Dose-response bioassays

Dose response bioassays were done with cordycepin and EPF alone against *G. mellonella* larvae. For the cordycepin dose-response bioassay, 10 final instar *G. mellonella* larvae were cooled on ice for 5 min, then injected in the right front proleg with 30 µl cordycepin at 1.0, 1.8, 3.3, 6.0 or 11 mg ml<sup>-1</sup> (equivalent to 30, 55, 100, 180, and 330 µg per insect respectively, with a sham injection control) using a 0.3 ml microfine insulin syringe (BD, Franklin Lakes, USA). Where the suspension bled from larvae following injection, the larvae were discarded and not included in the experiment. Immediately after treatment, insects were placed individually in Petri dishes on damp filter paper, sealed with Parafilm, and maintained in darkness at 20°C. Survival of the larvae was monitored every 24 h for seven days. The bioassay was repeated on three separate occasions, resulting in a total of 30 larvae in each treatment. The DRC package in R (Ritz et al., 2015) was used to estimate lethal concentration (LC) and lethal dose (LD) values at day 6. The EPF dose response bioassay was done in the same way, with final instar *G. mellonella* larvae being injected with 30 µl of conidia suspensions of *C. militaris* and *B.*

*bassiana* at concentrations of  $1 \times 10^3$ ,  $1 \times 10^4$ ,  $1 \times 10^5$ ,  $1 \times 10^6$  and  $1 \times 10^7$  conidia  $\text{ml}^{-1}$  (equivalent to 30, 300,  $3 \times 10^3$ ,  $3 \times 10^4$ ,  $3 \times 10^5$  conidia per insect respectively) in sterile 0.01% Triton X-100 (10 insects per treatment, sham injection controls, three independent repeats, LC and LD estimations at day 6).

### 2.3.2 EPF spore germination

The effect of cordycepin on EPF spore germination was measured as follows: for both *B. bassiana* and *C. militaris*, 100  $\mu\text{l}$  conidia suspension ( $1 \times 10^7$  conidia  $\text{ml}^{-1}$  in sterile 0.01% Triton X-100) was combined with 30  $\mu\text{l}$  of cordycepin (100  $\text{mg ml}^{-1}$ ) and made up to a final volume of 1 ml using Sabouraud Dextrose Broth (Sigma-Aldrich, St Louis, USA). This gave final concentrations of  $1 \times 10^6$  conidia  $\text{ml}^{-1}$  and 3.0  $\text{mg ml}^{-1}$  (12 mM) cordycepin (equivalent to the  $\text{LC}_{15}$  in the *G. mellonella* dose response bioassay). The mixture was incubated in darkness at  $23^\circ\text{C}$  for 24 h, after which the germination of approximately 100 conidia was recorded by examination under a microscope. A conidium was considered germinated if the germ tube was longer than the length of the conidium. The experiment was performed on three occasions. Percentage conidia germination was analysed using a one-way ANOVA following a logit transformation (Warton and Hui, 2011) and statistical normality testing (Shapiro and Wilk, 1965) in SPSS (IBM Corp. 2016).

### 2.3.3 Effect of cordycepin application on *G. mellonella* susceptibility to EPF

A bioassay was done to measure insect survival following EPF infection supplemented with cordycepin. Batches of 20 final instar *G. mellonella* larvae were injected, as described above, with 30  $\mu\text{l}$  conidia suspension of *B. bassiana* or *C. militaris* at concentrations of  $3.3 \times 10^2$  and  $3.3 \times 10^3$  conidia  $\text{ml}^{-1}$  (equivalent to 10 and 100 conidia per insect respectively) in 0.01% Triton X-100 with / without 3.0  $\text{mg ml}^{-1}$  cordycepin. Where the suspension bled from larvae following injection, the larvae were discarded and not included in the experiment. Assessment of larval survival was done as described above for 11 days and the experiment was repeated on three separate occasions, resulting in a total of 30 larvae for each treatment. For each insect batch, 10 individuals were used for survival analysis, while 5 individuals were removed at 0 h (to), 48 h and 72 h, and snap frozen in liquid nitrogen for RNA extraction and quantification of gene expression (Section 2.4). Changes in insect survival were visualised using a Kaplan-Meier estimator, and analysed using a Cox proportional-hazards regression model with replicate and treatment as factors, in which the median survival time (MST) of the insect populations of each

treatment and their 95% confidence intervals were calculated, and pairwise comparisons were done using a log-rank  $\chi^2$  test (Cox, 1972; IBM SPSS Statistics Version 24; Bewick et al., 2004). The outcome of combining cordycepin with *B. bassiana* or *C. militaris* on total percentage mortality (synergism, antagonism, or additive effect) at day 6 and day 11 was investigated using the fractional product method for combination treatments, where the effect of the combination is given as  $(1-X)$ , where  $X = (1-A)(1-B)$  and where A = the proportional effect of agent A on its own, and B is the proportional effect of agent B (Webb, 1963).

#### 2.4 Quantification of expression of insect immune-associated genes

RNA was extracted from *D. melanogaster* S2r+ cells using the ReliaPrep RNA Cell Miniprep system (Promega, Madison, USA) following manufacturer's instructions. For RNA extraction from *G. mellonella*, individual snap frozen larvae were ground in liquid nitrogen using an autoclaved mortar and pestle. RNA was extracted from 50 mg of material using the phenol: chloroform method with Tri-reagent (Sigma-Aldrich, St Louis, USA) then treated with DNase I (Sigma-Aldrich, St Louis, USA) following manufacturer's instructions. After washing with 75% ethanol, air dried RNA pellets were resuspended in 50  $\mu$ l DEPC-treated water and stored at -80 °C. RNA concentrations were measured using a NanoDrop® ND-100 spectrophotometer (Thermo Fisher Scientific, Waltham, USA). cDNA was synthesized using Superscript kits (Thermo Fisher Scientific, Waltham, USA): Superscript III was used to reverse transcribe *D. melanogaster* RNA (100  $\mu$ g) and Superscript II was used for *G. mellonella* RNA (500  $\mu$ g), both with random hexamers. Manufacturer's instructions were followed, with the exception that 0.5  $\mu$ l (100 units) SuperScript III was used for reactions. The expression of insect immune associated genes was quantified by RT-qPCR using a Lightcycler 480 (Roche Holding AG, Basel, Switzerland). For *D. melanogaster* S2r+ cells, expression levels were quantified for genes encoding the ribosomal protein RP49 (used as the reference), and the antimicrobial peptides (AMPs) metchnikowin and attacin. For *G. mellonella*, expression levels were quantified for genes encoding the following proteins: the ribosomal protein S7e (reference); the AMPs gallerimycin and galiomicin; lysozyme; and the insect metalloproteinase inhibitor (IMPI). Gene expression in *G. mellonella* was measured for larvae injected with EPF + / - exogenous cordycepin, as well as for sham injections and uninjected controls. Primers for all genes are shown in Table 1. SensiFAST SYBR No-ROX (Bioline, London, UK) was used for RT-qPCR, each reaction contained: 5  $\mu$ l SYBR Green, 2  $\mu$ l DEPC treated water, 1  $\mu$ l forward primer (10  $\mu$ M), 1  $\mu$ l reverse primer (10  $\mu$ M) and 1  $\mu$ l cDNA. The conditions were: 95°C 2

min, 40x (95°C 5 s, 60°C 10 s and 72°C 20s). The comparative  $C_T$  method (also known as the  $2^{-\Delta\Delta CT}$  method) (Schmittgen and Livak, 2008) was used to calculate fold changes in gene expression compared to untreated samples with correction for the internal control. Each gene of interest was analysed independently. For *D. melanogaster*, data on relative gene expression were tested for normality (Shapiro and Wilk, 1965) before being subject to ANOVA, while for *G. mellonella*, the data were tested for normality and then analysed using Kruskal-Wallis H tests (IBM SPSS Statistics Version 24). Where overall significant differences were observed, post-hoc analysis was performed using a Tukey HSD post hoc test (following ANOVA) or the Dunn-Bonferroni post hoc method (following Kruskal-Wallis tests).

### 3. Results

#### 3.1 Effect of cordycepin treatment on gene expression in *D. melanogaster* S2r+ cells following immune challenge

The relative expression of the immune effector genes *metchnikowin* and *attacin* was quantified in a *D. melanogaster* S2r+ cell line treated separately with two activators of the immune response, curdlan and LPS, with and without the addition of 25  $\mu\text{g ml}^{-1}$  cordycepin (Fig. 1). There was significant variation among treatments in the relative expression for both *metchnikowin* ( $F_{5,17} = 5.93$ ,  $P < 0.01$ ) and *attacin* A ( $F_{5,17} = 3.48$ ,  $P < 0.05$ ). Application of curdlan significantly elevated the mean relative expression of *metchnikowin* compared to the DMSO-only control or DMSO plus cordycepin (Tukey HSD,  $P < 0.05$ , approximately 6-fold increase). When cordycepin was co-applied with curdlan, there was no significant difference in mean relative gene expression compared to DMSO-only or DMSO plus cordycepin ( $P > 0.05$ , approximately 2.5-fold change), nor was there a significant difference between the mean relative expression of *metchnikowin* for the curdlan-only treatment vs. curdlan + cordycepin ( $P > 0.05$ , 2.5-fold change). Treatment with LPS, or LPS + cordycepin, did not have a significant effect on relative expression of *metchnikowin* compared to the DMSO-only control ( $P > 0.05$ , 2-fold increase for LPS only, no fold change for LPS + cordycepin) (Fig. 1A). For *attacin* A, post hoc comparisons showed that relative expression was significantly higher following treatment with curdlan compared to the DMSO plus cordycepin- treatment ( $P < 0.05$ , approximately 7-fold increase) (Fig. 1B).



### 3.2 Effect of cordycepin treatment on the survival of *G. mellonella* larvae infected with *C. militaris* and *B. bassiana*

In this set of experiments we investigated whether application of exogenous cordycepin would affect the susceptibility of *G. mellonella* larvae to fungal infection and impact on the expression of host immune genes. We used a low dose of cordycepin that did not inhibit fungal activity and did not cause excessive insect mortality on its own.

#### 3.2.1 Dose-response bioassays of *B. bassiana* and *C. militaris* against *G. mellonella* larvae

Larvae of *G. mellonella* were treated with either EPF or cordycepin to established the impact of these treatments individually on insect survival. There was a positive relationship between *G. mellonella* mortality and dose of cordycepin applied (Fig. 2, Weibull type 1 model), the  $LC_{15}$ ,  $LC_{50}$ ,  $LD_{15}$  and  $LD_{50}$  were also calculated on day 6 after treatment and found to be 2.91 ( $\pm 0.35$ ) mg ml<sup>-1</sup>, 6.96 ( $\pm 1.03$ ) mg ml<sup>-1</sup>, 87.95 ( $\pm 10.79$ )  $\mu$ g per larvae and 209.34 ( $\pm 30.70$ )  $\mu$ g per larvae, respectively. Individual larvae receiving a lethal dose of cordycepin became distinctly grey in colour between 24-48 h prior to death (Supplementary Fig. 1) and their movement declined markedly. After death, cadavers turned black and their abdomens were fluid-filled. *G. mellonella* larvae were susceptible to lethal infections by both *B. bassiana* and *C. militaris* (Fig. 3; estimated *B. bassiana*  $LC_{50}$  (day 6) =  $3.6 \times 10^3$  ( $\pm 9.4 \times 10^2$ ) conidia ml<sup>-1</sup>,  $LD_{50}$  =  $1.1 \times 10^2$  ( $\pm 2.8 \times 10^1$ ) conidia per insect; *C. militaris*  $LC_{50}$  =  $4.0 \times 10^4$  ( $\pm 9.1 \times 10^3$ ) conidia ml<sup>-1</sup>,  $LD_{50}$  =  $1.2 \times 10^3$  ( $\pm 2.8 \times 10^2$ ) conidia per insect; two-parameter log-logistic model).

#### 3.2.2 EPF conidia germination

Conidia of *C. militaris* and *B. bassiana* were incubated with cordycepin to identify any inhibitory effects on germination. There was no significant effect of cordycepin treatment (3 mg ml<sup>-1</sup>) on the numbers of *B. bassiana* and *C. militaris* conidia germinating *in vitro* after 24 h ( $F_{3,11} = 1.045$ ,  $P > 0.05$ ). The mean germination (back-transformed) of *B. bassiana* conidia populations was 75% and 79% at 0 and 3 mg ml<sup>-1</sup> of cordycepin respectively, while for *C. militaris* conidia it was 82% and 79% respectively (Fig. 4).

#### 3.2.3 Effect of cordycepin application on *G. mellonella* susceptibility to fungal infection

A laboratory bioassay was done to quantify the effects on *G. mellonella* survival of co-applying 3 mg ml<sup>-1</sup> cordycepin with *C. militaris* or *B. bassiana* at doses of 10 and 100 conidia per insect. Mortality of *G. mellonella* larvae in the sham injection control was 3.3% at 11 days post injection (dpi) (Supplementary Table 2). Injection of larvae with 3 mg ml<sup>-1</sup> cordycepin resulted

in a mortality of 13.3%, which was in keeping with the results from the earlier cordycepin dose response bioassay (Fig. 2) but was not statistically significantly different from the sham injection control (Supplementary Table 2). Compared to injection of EPF on their own, co-injection of 3 mg ml<sup>-1</sup> cordycepin with *B. bassiana* or *C. militaris* caused a significant decrease in median survival time of c. 24 h in all cases (Supplementary Table 2, log rank chi squared > 3.841, P < 0.05; also see survival curves, Fig. 5) and this was reflected by an increase in the hazards ratio (Supplementary Table 2). A dose of 10 conidia per insect of *C. militaris* resulted in 36.7% mortality at 11 dpi, meaning that median survival time could not be estimated (Supplementary Table 2). At day 6 the predicted mean mortality of *B. bassiana* + cordycepin combination, calculated from the fractional product of *B. bassiana* and cordycepin mortality individually, was 0% at a dose of 10 conidia per insect (observed value = 26.7%) and 46.7% at a dose of 100 conidia per insect (observed value = 80%). At day 11 the predicted mean mortality of the *C. militaris* + cordycepin combination was 45.1% at a dose of 10 conidia per insect (observed value = 80%), and 65.3% at a dose of 100 conidia per insect (observed value = 80%). Both the results on days 6 and 11 after treatment indicate that there was a synergistic effect of the combination treatment with EPF and cordycepin (Webb, 1963).

### 3.3 Effect of cordycepin treatment on the expression of immune-related genes in *G. mellonella* larvae infected with *B. bassiana* and *C. militaris*

In order to provide information on the possible mechanism by which cordycepin affects fungal infection, the expression of four *G. mellonella* immune-related genes (*gallerimycin*, *gallerimycin*, *IMPI* and *lysozyme*) was quantified at 48 h and 72 h in larvae injected with *B. bassiana* or *C. militaris* +/- cordycepin (Fig. 6). The experiment also included a sham injection control and an untreated (= un-injected) control. Pairwise comparisons (Dunn-Bonferroni method) showed that, at 48 h, injection with *C. militaris*, *B. bassiana*, and sham injection resulted in a significantly larger increase (P < 0.05) in relative expression of *IMPI* and *lysozyme* compared to the untreated control. At 72 h, injection with *C. militaris* and *B. bassiana* resulted in a significantly larger increase (P < 0.05) in relative expression of *gallerimycin*, *IMPI*, and *lysozyme* compared to the untreated control, but there was no significant difference between sham injection and the untreated control for any of the four genes (P > 0.05). In contrast, there was no significant increase (P > 0.05) in relative immune gene expression when cordycepin was co-applied with *C. militaris*, *B. bassiana* or on its own.

Addition of cordycepin caused between 2-fold to 7-fold reduction in relative gene expression depending on the EPF species and immune gene (Fig. 6). Compared to injection with *B. bassiana* only, co-injection of *B. bassiana* + cordycepin resulted in a reduction in expression of: (i) *galiomicin* by ~3.5x (P < 0.01) at 48h and ~3x (P < 0.01) at 72 h; (ii) *gallerimycin* by ~3x at both 48h (P < 0.05) and 72 h (P < 0.05); (iii) *IMPI* by ~7.5x (P < 0.001) at 48h and ~5x (P < 0.001) at 72h; and (iv) *lysozyme* by ~6.5x (P < 0.001) at 48h and ~3.5x (P < 0.001) at 72h. A similar pattern of reduced gene expression was observed with *C. militaris* and cordycepin. Compared to injection with *C. militaris* only, co-injection of *C. militaris* + cordycepin resulted in a reduction in expression of: (i) *galiomicin* by ~5x (P < 0.001) at 48h and ~4.5x (P < 0.001) at 72 h; (ii) *gallerimycin* by ~2x at both 48h (P < 0.01) and 72 h (P < 0.05); (iii) *IMPI* by ~4x (P < 0.005) at 48h and ~6x (P < 0.001) at 72h; and (iv) *lysozyme* by ~6.5x (P < 0.01) at 48h and ~4x (P < 0.005) at 72h. Compared to the sham injection control, injection of cordycepin resulted in a reduction in expression of: (i) *galiomicin* by ~3x (P < 0.01) at 72h ; (ii) *gallerimycin* by ~2x (P < 0.02) at 48h; (iii) *IMPI* by ~3x (P < 0.001) at 48h and ~3x (P < 0.001) at 72h; and (iv) *lysozyme* by ~2.5x (P < 0.001) at 48h.

#### 4. Discussion

In order to complete their parasitic life cycle, obligately-killing EPF such as *C. militaris* must nullify the effects of host immune defences (Vey et al., 2002; Wang and St Leger 2006; Pal et al 2007; Butt et al., 2016). The inference from our findings is that the *C. militaris* secondary metabolite cordycepin is capable of acting as an insect immune modulator, dampening down the upregulation of defence genes in response to fungal infection. This suppression of the immune response may facilitate infection of insects by *C. militaris*. To our knowledge, this is the first report of the effects of cordycepin on immune gene expression in insects.

The statistical variation that we observed in the *D. melanogaster* S2r+ cell assay means that some caution is needed in interpretation of the results, but overall, they suggest that cordycepin may have an inhibitory effect on expression of the anti-fungal AMP gene *metchnikowin*, despite the downregulation in expression of this gene not being significant following curdlan treatment. Specifically, while immune challenge with the anti-fungal elicitor curdlan resulted in increased expression of *metchnikowin* compared to the DMSO-only control, this did not occur when curdlan was applied in the presence of cordycepin, suggesting that cordycepin was able to inhibit upregulation of this immune gene (post hoc comparisons showed no significant difference between treatment with curdlan vs. curdlan + cordycepin, but we expect that higher

doses of cordycepin are likely to have resulted in a greater effect on gene expression). *Metchnikowin* (Levashina et al., 1995), which is specific to members of the Drosophilidae, is synthesized via the Toll and Imd pathways in response to fungal or bacterial infection (De Gregorio et al., 2002; Imler and Bulet, 2005). As expected, treatment with the anti-(gram negative) bacterial elicitor LPS (Tanji and Ip, 2005) did not result in increased *metchnikowin* expression compared to the control. However, unexpectedly, we observed no significant increase in expression of the AMP gene *attacin A* in response to LPS in our study. Attacins are conserved across different species in the Lepidoptera and Diptera, they are broadly considered to be active against gram-negative bacteria (Lemaitre and Hoffmann, 2007; Yi et al., 2014), although in *D. melanogaster* there are multiple *attacin* genes (A, B1, D) which are expressed in response to gram-negative bacteria and fungi. The expression of *attacin D* is controlled by the IMD pathway, which is activated by gram-negative bacteria (De Gregorio et al., 2002). The expression of *attacins A* and B1 is controlled by both the Toll and IMD pathways and responds to infections by fungi and bacteria (De Gregorio et al., 2002). Previous experiments with *Aedes aegypti* and *D. melanogaster* have found that *attacin* is expressed in response to *B. bassiana* infection (Lemaitre et al., 1997; De Gregorio et al., 2002; Dong et al., 2012; Yi et al., 2014). Our finding of significantly increased *attacin A* expression in the *D. melanogaster* S2r+ cell assay in the curdlan vs. cordycepin-only treatment, provides some tentative support that *attacin A* expression is responsive to fungal infection.

The inhibitory effects of cordycepin on the anti-fungal immune insect immune response was further supported by the results from our *G. mellonella* experiment, where application of cordycepin prevented increased expression of different immune effector genes in response to fungal infection, and resulted in faster host insect death. Injection of *G. mellonella* larvae with *B. bassiana*, *C. militaris*, or a sham injection, resulted in increased expression of *lysozyme* and *IMPI* at 48 h after treatment. Meanwhile, at 72 h we observed elevated expression of *lysozyme*, *IMPI*, and *gallerimycin* (but not *galiomycin*) in response to *B. bassiana* and *C. militaris* injection, but there was no elevated immune gene expression in response to sham injection compared to the untreated control, suggesting that the gene expression response to wounding was short lived. Applying exogenous cordycepin alongside *B. bassiana* or *C. militaris*, or as an addition to the sham injection, dampened down the expression of these immune genes so that relative expression was no different from the untreated control. *Gallerimycin* and *galiomycin* are both anti-fungal defensins (Schuhmann et al., 2003; Lee et al., 2004; Langen et al., 2006) synthesized via the Toll pathway in response to EPF infection (Vertyporokh and

Wojda 2017). Lysozyme has both anti-bacterial and anti-fungal activity (Jollès and Jollès, 1984; Wojda et al., 2009; Sowa-Jasilek et al., 2016 ), while IMPI is a *G. mellonella*-specific inhibitor of EPF metalloproteinases that hydrolyses insect proteins in the cuticle and haemolymph (Vilcinskas and Wedde, 2002; Altincicek et al., 2007; Wedde et al., 2007), and is itself a target for EPF proteinases synthesized as a fungal countermeasure to the *G. mellonella* immune response (Mukherjee and Vilcinskas, 2018). Expression of *IMPI* has been shown previously to be up-regulated within 48 h of fungal infection (Vertyporokh and Wojda, 2017). Expression of both *IMPI* and *lysozyme* in *G. mellonella* is reported to be upregulated by a range of stimuli including fungal and bacterial infection (Vertyporokh and Wojda, 2017), metamorphosis and wounding (Griesch et al., 2000; Altincicek and Vilcinskas 2006; Vilcinskas, 2019). The signalling pathway(s) leading to *IMPI* and *lysozyme* expression in *G. mellonella* have yet to be fully elucidated, although they are likely to be regulated by Toll, Imd or a related pathway as their transcription is controlled by c-Rel proteins Relish, DIF and Dorsal (Altincicek and Vilcinskas, 2006; Vilcinskas, 2019).

We observed no upregulation of *galiomycin* in response to fungal infection in *G. mellonella*, which may have been due to the route of infection (injection rather than topical application) or because gene expression was measured using whole insects rather than specific tissues, such as the fat body, which is a site of AMP synthesis (Vertyporokh and Wojda, 2017). We chose injection as the way to deliver cordycepin to *G. mellonella* as it mimics the normal route of exposure, since host tissues are exposed to it after fungal hyphae have grown into the insect haemocoel. Injection of EPF also allowed us to precisely control the dose of pathogen, and to ensure that cordycepin and EPF were applied at the same time and location.

To date, studies on the effects of cordycepin have focused on the mammalian innate response rather than the insect immune system (for example see Chu and Edelman, 1972; Penman et al., 1970; Kim et al., 2006; Jeong et al., 2010; Kondrashov et al., 2012; Ren et al., 2012; Ashraf et al., 2019). The close similarities between aspects of the mammalian innate immune system and insect immunity - such as the homology between TLR (mammal) and Toll (insect) immune signalling (Sheehan et al., 2018) - would explain why cordycepin is able to affect both mammalian and insect immune responses. Cordycepin treatment is reported to reduce inflammatory mRNAs in primary human airway smooth muscle cells and mouse macrophage cells treated with an inflammatory stimulus (Kondrashov et al., 2012; Ashraf et al., 2019). The precise mechanism(s) of action of cordycepin on mammalian cells is not yet known, although

there is evidence that it can prevent nuclear translocation of NFκB in mammalian microglia, embryonic kidney and macrophage cells (Jeong et al., 2010; Ren et al., 2012; Ashraf et al., 2019), while it has also been shown to affect a subset of polyadenylated mRNAs in a variety of human cell types, (Penman et al., 1970; Chu and Edelman, 1972; Kondrashov et al., 2012).

Secondary metabolites play an important role in the pathogenicity of EPF (de Bekker et al., 2013) and - in principle - the deployment of metabolites for active suppression of the insect immune response would be beneficial for fungal fitness. The array of secondary metabolites produced by non-specialist EPF has been shown to vary between species, insect tissue type, and whether fungal growth is infectious or saprotrophic (i.e. in living or dead tissue) (de Bekker et al., 2013). It is possible that some of these compounds function as immunosuppressors, but may also have additional roles. For example, injection of destruxin A into *D. melanogaster*, mimicking infection by *Metarhizium* species, reduced expression of AMPs and made them susceptible to infection by normally non-pathogenic *E. coli* by inhibition of the IMD pathway (Pal et al., 2007). However destruxin A (as well as destruxins B and E) has also been found to have insecticidal activity against *Spodoptera litura* (Lepidoptera) and, in the same study, a crude destruxin preparation was found to have synergistic effects on the pathogenicity of *Cordyceps javanica* (Hypocreales, Cordycipitaceae) (formerly *Paecilomyces javanicus* and *Isaria javanica*) to *S. litura* (Hu et al., 2007), which echoes our finding of a synergistic interaction between co-application of exogenous cordycepin together with *C. militaris* or *B. bassiana*.

Few studies have been done on the mechanisms of pathogenesis by *C. militaris*, but there is evidence that passive avoidance of immune detection is less important for successful infection compared to generalist EPF species (Kryukov et al., 2020). Our finding that cordycepin was able to suppress the expression of different antimicrobial effectors is likely to be of benefit to *C. militaris* as an immune interference strategy, as insect defence compounds are known to operate synergistically against pathogen infections (Kaur et al., 2015; Butt et al., 2016). Therefore, the ability to suppress the activity of a range of defence compounds at once might be a more efficient - and therefore selectively advantageous - strategy than targeting individual compounds with different suppressors. The production of cordycepin by *C. militaris* has been found to vary between 750-16500 µg g<sup>-1</sup> depending on fungal strain, cultivation method and extraction method (Huang et al., 2003; Li et al., 2004; Yang and Li, 2008). Therefore, the

amount of cordycepin used in our experiments probably represents a higher dose than that received by a *G. mellonella* larva (which weighed 200 - 300 mg) when infected with *C. militaris* on its own: the evidence for this is that injection of *C. militaris* into *G. mellonella* still resulted in an increase in expression of immune related genes, even though this fungus is known to synthesize cordycepin. This raises a question of why, if cordycepin is able to suppress the immune response and increase the virulence of the fungus, it is not produced at higher levels in natural infections by *C. militaris*? It has been proposed that there is a trade-off between metabolite production and EPF growth, as highly active toxins could result in host mortality before the fungus has grown sufficient biomass to produce structures needed for spore production, and which would also leave the host cadaver vulnerable to fast growing, saprotrophic microbes (Boomsma et al., 2014). In our study, cordycepin was lethal to *G. mellonella* larvae depending on the dose applied. Larvae killed by cordycepin injection exhibited the symptoms of a bacterial infection (Tanada and Kaya, 2012) and we were able to recover opportunist bacterial pathogens from them including *Bacillus* sp., *Enterobacter* sp., *Enterococcus* sp. *Pantoea* sp. and *Serratia* sp. (Woolley, 2018). Similar symptoms were observed by Kim et al., (2002), who found that larvae of the diamondback moth *Plutella xylostella* fed leaf discs treated with cordycepin (25-500 mg l<sup>-1</sup>) became dark brown and then lysed after death. A methanolic extract of the *C. militaris* fruiting body has been found to exhibit antibacterial and antifungal properties (Reis et al., 2013), which suggests that *in vivo*, *C. militaris* produces metabolites that may prevent opportunistic infections of the fruiting body by bacteria or fungi.

Further work needs to be conducted to elucidate the mode of action of cordycepin during insect infection by *C. militaris*. In particular, it would be important to determine how cordycepin inhibits AMP expression, including effects on the activity of transcription factors in the Toll and Imd pathways (Kondrashov et al., 2012; Ren et al., 2012) and polyadenylation of immune genes (Kondrashov et al., 2012). It would also be worth investigating whether cordycepin has an effect on the cellular immune response, including the prophenol oxidase cascade and wounding response. Although our study suggests that cordycepin may play a role in the virulence of *C. militaris*, conclusive proof would be needed by construction of knock outs of the genes responsible for cordycepin production (Xia et al., 2017). Improved knowledge of the EPF-host immune interaction may help us understand the selection pressures affecting fungal pathogenicity and host immune defences (Boomsma et al., 2014; Mukherjee and Vilcinskis, 2018), as well as practical applications such as the selection and development of

EPF strains for biological pest control or the identification of secondary metabolites as novel pesticides, efficacy enhancing agents, or pharmaceuticals (Butt et al., 2016).

### **Declaration of Competing Interest**

The authors declare that they have no competing interests that could have influenced the work reported in this paper.

### **Acknowledgements**

We would like to thank Dr Sonia Lopez de Quinto of Cardiff University UK for providing the *Drosophila* cell line used in this study, and the USDA ARS Collection of Entomopathogenic Fungal Cultures (ARSEF) for providing EPF cultures. This work was funded by the Biotechnology and Biosciences Research Council (BBSRC) as part of the Midlands Integrative Biosciences Training Partnership (MIBTP).

### **References**

- Altincicek, B., Vilcinskas, A., 2006. Metamorphosis and collagen-IV-fragments stimulate innate immune response in the greater wax moth, *Galleria mellonella*. *Dev. Comp. Immunol.* 30, 1108-1118.
- Altincicek, B., Linder, M., Linder, D., Preissner, K.T., Vilcinskas, A., 2007. Microbial metalloproteinases mediate sensing of invading pathogens and activate innate immune responses in the lepidopteran model host *Galleria mellonella*. *Infect. Immune.* 75(1), 175-183.
- Ashraf, S., Radhi, M., Gowler, P., Burston, J.J., Gandhi, R.D., Thorn, G.J., Piccinini, A.M., Walsh, D.A., Chapman, V., De Moor, C.H., 2019. The polyadenylation inhibitor cordycepin reduces pain, inflammation and joint pathology in rodent models of osteoarthritis. *Sci. Rep.* 9, 4696.
- Bergin, D., Murphy, L., Keenan, J., Clynes, M., Kavanagh, K., 2006. Pre-exposure to yeast protects larvae of *Galleria mellonella* from a subsequent lethal infection by *Candida albicans* and is mediated by the increased expression of antimicrobial peptides. *Microbes Infect.* 8, 2105-2112.
- Bewick, V., Cheek L., Ball, J., 2004. Statistics review 12: Survival analysis. *Crit.Care* 8, 389-394.
- Boomsma, J.J., Jensen, A.B., Meyling, N.V., Eilenberg, J., 2014. Evolutionary interaction networks of insect pathogenic fungi. *Annu. Rev. Entomol.* 59, 467-485.
- Butt, T.M., Coates, C.J., Dubovskiy, I.M., Ratcliffe, N.A., 2016. Entomopathogenic fungi: new insights into host-pathogen interactions. In *Genetics and Molecular Biology of Entomopathogenic Fungi* (Eds. Lovett, B., St Leger, R. J.), Academic Press. pp. 307-364.



- Castillo, J.C., Shokal, U., Eleftherianos, I., 2013. Immune gene transcription in *Drosophila* adult flies infected by entomopathogenic nematodes and their mutualistic bacteria. *J. Insect Physiol.* 59, 179-185.
- Chandler, D., 1994. Cryopreservation of fungal spores using porous beads. *Mycol. Res.* 98, 525-526.
- Chandler, D., 2016. Basic and applied research on entomopathogenic fungi. In *Microbial Agents of Insect and Mite Pests: from theory to practice* (Ed. L. Lacey), Academic Press, pp. 42-60.
- Chu, L.L., Edelman, I.S., 1972. Cordycepin and  $\alpha$ -amanitin: Inhibitors of transcription as probes of aldosterone action. *J. Membrane Biol.* 10, 291-310.
- Cox, D.R., 1972. Regression models and life-tables. *J. R. Stat. Soc. B.* 34, 187-220.
- Cunningham, K.G., Manson, W., Spring, F.S., Hutchinson, S.A., 1950. Cordycepin, a metabolic product isolated from cultures of *Cordyceps militaris* (Linn.) Link. *Nature* 166, 949.
- De Bekker, C., Smith, P.B., Patterson, A.D., Hughes, D.P., 2013. Metabolomics reveals the heterogeneous secretome of two entomopathogenic fungi to *ex vivo* cultured insect tissues. *PLoS ONE* 8, e70609.
- De Gregorio, E., Spellman, P.T., Tzou, P., Rubin, G.M., Lemaitre, B., 2002. The Toll and Imd pathways are the major regulators of the immune response in *Drosophila*. *EMBO J.* 21, 2568-2579.
- Dong, Y., Morton, J.C., Ramirez, J.L., Souza-Neto, J.A., Dimopoulos, G., 2012. The entomopathogenic fungus *Beauveria bassiana* activate toll and JAK-STAT pathway-controlled effector genes and anti-dengue activity in *Aedes aegypti*. *Insect Biochem. Mol.* 42, 126-132.
- Fujita, T., Inoue, K., Yamamoto, S., Ikumoto, I., Sasaki, S., Toyama, R., Chiba, K., Hoshino, Y., Okumoto, T., 1994. Fungal metabolites. Part 11. A potent immunosuppressive activity found in *Isaria sinclairii* metabolite. *J. Antibiot.* 47, 208-215.
- Gobert, V., Gottar, M., Matskevich, A.A., Rutschmann, S., Royet, J., Belvin, M., Hoffmann, J.A., Ferrandon, D., 2003. Dual activation of the *Drosophila* toll pathway by two pattern recognition receptors. *Science* 302, 2126-2130.
- Griesch, J., Wedde, M., Vilcinskas, A., 2000. Recognition and regulation of metalloproteinase activity in the haemolymph of *Galleria mellonella*: a new pathway mediating induction of humoral immune responses. *Insect Biochem. Molec.* 30, 461-472.
- Harding, C.R., Schroeder, G.N., Collins, J.W., Frankel, G., 2013. Use of *Galleria mellonella* as a model organism to study *Legionella pneumophila* infection. *J. Vis. Exp.* 81, 50964.
- Hu, Q.B., Ren, S.X., An, X.C., Qian, M.H., 2007. Insecticidal activity influence of destruxins on the pathogenicity of *Paecilomyces javanicus* against *Spodoptera litura*. *J. Appl. Entomol.* 131, 262-268.

- Huang, L.F., Liang, Y.Z., Guo, F.Q., Zhou, Z.F., Cheng, B.M., 2003. Simultaneous separation and determination of active components in *Cordyceps sinensis* and *Cordyceps militaris* by LC/ESI-MS. *J. Pharmaceut. Biomed.* 33, 1155-1162.
- IBM Corp. Released 2016. IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.
- Imler, J.L., Bulet, P., 2005. Antimicrobial peptides in *Drosophila*: structures, activities and gene regulation. In *Mechanisms of epithelial defense, Vol. 86*. Karger Publishers. pp. 1-21.
- Jeong, J.W., Jin, C.Y., Kim, G.Y., Lee, J.D., Park, C., Kim, G.D., Kim, W.J., Jung, W.K., Seo, S.K., Choi, I.W., Choi, Y.H., 2010. Anti-inflammatory effects of cordycepin via suppression of inflammatory mediators in BV2 microglial cells. *Int. Immunopharmacol.* 10, 1580-1586.
- Jin, L.H., Shim, J., Yoon, J.S., Kim, B., Kim, J., Kim-Ha, J., Kim, Y.J., 2008. Identification and functional analysis of antifungal immune response genes in *Drosophila*. *PLoS Pathog.* 4, e1000168.
- Johnston, J.B., 2011. Mechanism of action of pentostatin and cladribine in hairy cell leukemia. *Leuk. Lymphoma*, 52(Suppl 2), 43-45.
- Jollès, P., Jollès, J., 1984. What's new in lysozyme research?. *Mol. Cell. Biochem.* 63, 165-189.
- Kaur, H.P., Singh, B., Thakur, A., Kaur, A., Kaur, S., 2015. Studies on immunomodulatory effect of endophytic fungus *Alternaria alternata* on *Spodoptera litura*. *J. Asia-Pac. Entomol.* 18, 67-75.
- Kepler, R.M., Luangsa-ard, J.J., Hywel-Jones, N.L., Quandt, C.A., Sung, G.H., Rehner, S.A., Aime, M.C., Henkel, T.W., Sanjuan, T., Zare, R., Chen, M.J., Li, Z.Z., Rossman, A. Y., Spatafora, J. W., Shrestha, B., 2017. A phylogenetically-based nomenclature for Cordycipitaceae (Hypocreales). *IMA Fungus* 8, 335-353.
- Kim, H.G., Shrestha, B., Lim, S.Y., Yoon, D.H., Chang, W.C., Shin, D.J., Han, S.K., Park, S.M., Park, J.H., Park, H.I., Sung, J.M., 2006. Cordycepin inhibits lipopolysaccharide-induced inflammation by the suppression of NF- $\kappa$ B through Akt and p38 inhibition in RAW 264.7 macrophage cells. *Eur. J. Pharmacol.* 545, 192-199.
- Kim, J.R., Yeon, S.H., Kim, H.S., Ahn, Y.J., 2002. Larvicidal activity against *Plutella xylostella* of cordycepin from the fruiting body of *Cordyceps militaris*. *Pest Man. Sci.* 58, 713-717.
- Kondrashov, A., Meijer, H.A., Barthet-Barateig, A., Parker, H.N., Khurshid, A., Tessier, S., Sicard, M., Knox, A.J., Pang, L., De Moor, C.H., 2012. Inhibition of polyadenylation reduces inflammatory gene induction. *RNA* 18, 2236-2250.
- Kryukov, V.Y., Kryukova, N.A., Tomilova, O.G., Vorontsova, Y., Chertkova, E., Pervushin, A.L., Sepneva, I., Glupov, V.V., Yaroslavtseva, O.N., 2020. Comparative analysis of the immune response of the wax moth *Galleria mellonella* after infection with the fungi *Cordyceps militaris* and *Metarhizium robertsii*. *Microb. Pathog.* 141, 103995.


- Kumar, H., Kumagai, Y., Tsuchida, T., Koenig, P.A., Satoh, T., Guo, Z., Jang, M.H., Saitoh, T., Akira, S., Kawai, T., 2009. Involvement of the NLRP3 inflammasome in innate and humoral adaptive immune responses to fungal  $\beta$ -glucan. *J. Immunol.* 183, 8061-8067.
- Langen, G., Imani, J., Altincicek, B., Kieseritzky, G., Kogel, K.H., Vilcinskas, A., 2006. Transgenic expression of gallerimycin, a novel antifungal insect defensin from the greater wax moth *Galleria mellonella*, confers resistance to pathogenic fungi in tobacco. *Biol. Chem.* 387, 549-557.
- Lee, Y.S., Yun, E.K., Jang, W.S., Kim, I., Lee, J.H., Park, S.Y., Ryu, K.S., Seo, S.J., Kim, C.H., Lee, I.H., 2004. Purification, cDNA cloning and expression of an insect defensin from the great wax moth, *Galleria mellonella*. *Insect Mol. Biol.* 13, 65-72.
- Lemaitre, B., Hoffmann, J., 2007. The Host Defense of *Drosophila melanogaster*. *Annu. Rev. Immunol.* 25, 697-743.
- Lemaitre, B., Kromer-Metzger, E., Michaut, L., Nicolas E, Meister, M., Georgel, P., Reichhart, J-M., Hoffman, J.A., 1995. A recessive mutation, immune deficiency (Imd), defines two distinct control pathways in the *Drosophila* host defense. *Proc. Nat. Acad. Sci. USA* 92, 9465-9469.
- Lemaitre, B., Reichhart, J. M., Hoffmann, J.A., 1997. *Drosophila* host defense: differential induction of antimicrobial peptide genes after infection by various classes of microorganisms. *Proc. Nat. Acad. Sci. USA* 94, 14614-14619.
- Levashina, E.A., Ohresser, S., Bulet, P., Reichhart, J.M., Hetru, C., Hoffmann, J.A., 1995. Metchnikowin, a novel immune-inducible proline rich mpeptide from *Drosophila* with antibacterial and antifungal properties. *Eur. J. Biochem.* 233, 694-700.
- Li, S., Li, P., Lai, C., Gong, Y., Kan, K.K., Dong, T.T., Tsim, K.W., Wang, Y., 2004. Simultaneous determination of ergosterol, nucleosides and their bases from natural and cultured *Cordyceps* by pressurised liquid extraction and high-performance liquid chromatography. *J. Chromatogr. A* 1036, 239-243.
- Lu, H., Li, X., Zhang, J., Shi, H., Zhu, X., He, X., 2014. Effects of cordycepin on HepG2 and EA.hy926 cells: Potential antiproliferative, antimetastatic and anti-angiogenic effects on hepatocellular carcinoma. *Oncol. Lett.* 7, 1556-1562.
- Mukherjee, K., Vilcinskas, A., 2018. The entomopathogenic fungus *Metarhizium robertsii* communicates with the insect host *Galleria mellonella* during infection. *Virulence* 9, 402-413.
- Mylonakis, E., Moreno, R., El Khoury, J.B., Idnurm, A., Heitman, J., Calderwood, S.B., Ausubel, F.M., Diener, A., 2005. *Galleria mellonella* as a model system to study *Cryptococcus neoformans* pathogenesis. *Infect. Immun.* 73, 3842-3850.
- Nakamura, K., Yoshikawa, N., Yamaguchi, Y.U., Kagota, S., Shinozuka, K., Kunitomo, M., 2006. Antitumor effect of cordycepin (3'-deoxyadenosine) on mouse melanoma and lung carcinoma cells involves adenosine A3 receptor stimulation. *Anticancer Res.* 26, 43-47.

- O'Neill, L.A., 2006. How Toll-like receptors signal: what we know and what we don't know. *Curr. Opin. Immunol.* 18, 3-9.
- Pal, S., Leger, R.J.S., Wu, L.P., 2007. Fungal peptide Destruxin A plays a specific role in suppressing the innate immune response in *Drosophila melanogaster*. *J. Biol. Chem.* 282, 8969-8977.
- Paterson, R.R.M., 2008. Cordyceps—a traditional Chinese medicine and another fungal therapeutic biofactory?. *Phytochemistry* 69, 1469-1495.
- Penman, S., Rosbash, M., Penman, M., 1970. Messenger and heterogeneous nuclear RNA in HeLa cells: differential inhibition by cordycepin. *Proc. Nat. Acad. Sci. USA* 67, 1878-1885.
- Ramarao, N., Nielsen-Leroux, C., Lereclus, D., 2012. The insect *Galleria mellonella* as a powerful infection model to investigate bacterial pathogenesis. *J. Vis. Exp.* 70, 4392.
- Rämet, M., Pearson, A., Manfrulli, P., Li, X., Koziel, H., Göbel, V., Chung, E., Krieger, M., Ezekowitz, R.A.B., 2001. *Drosophila* scavenger receptor CI is a pattern recognition receptor for bacteria. *Immunity* 15, 1027-1038.
- Reis, F.S., Barros, L., Calhela, R.C., Ćirić, A., Van Griensven, L.J., Soković, M., Ferreira, I.C., 2013. The methanolic extract of *Cordyceps militaris* (L.) Link fruiting body shows antioxidant, antibacterial, antifungal and antihuman tumor cell lines properties. *Food Chem. Toxicol.* 62, 91-98.
- Ren, Z., Jianhua, C., Huo, Z., Xue, J., Cui, H., Luo, B., Jiang, L., Yang, R., 2012. Cordycepin suppresses TNF- $\alpha$ -induced NF $\kappa$ B activation by reducing p65 transcriptional activity, inhibiting I $\kappa$ B $\alpha$  phosphorylation, and blocking IKK $\gamma$  ubiquitination. *Int. Immunopharmacol.* 14, 698-703.
- Ritz, C., Baty, F., Streibig, J.C., Gerhard, D., 2015. Dose-response analysis using R. *PloS One* 10, e0146021.
- Roy, H.E., Steinkraus, D.C., Eilenberg, J., Hajek, A.E., Pell, J.K., 2006. Bizarre interactions and endgames: entomopathogenic fungi and their arthropod hosts. *Annu. Rev. Entomol.* 51, 331-357.
- Schmid-Hempel, P., 2009. Immune defence, parasite evasion strategies and their relevance for 'macroscopic phenomena' such as virulence. *Philos. T. Roy. Soc. B* 364, 85-98.
- Schmittgen, T.D., Livak, K.J., 2008. Analyzing real-time PCR data by the comparative C<sub>T</sub> method. *Nat. Protoc.* 3, 1101.
- Schneider, I., 1972. Cell lines derived from late embryonic stages of *Drosophila melanogaster*. *Development* 27, 353-365.
- Schuhmann, B., Seitz, V., Vilcinskas, A., Podsiadlowski, L., 2003. Cloning and expression of gallerimycin, an antifungal peptide expressed in immune response of greater wax moth larvae, *Galleria mellonella*. *Arch. Insect Biochem. Physiol.* 53, 125-133.


- Shapiro, S.S., Wilk, M.B., 1965. An analysis of variance test for normality (complete samples). *Biometrika* 52, 591-611.
- Sheehan, G., Garvey, A., Croke, M., Kavanagh, K., 2018. Innate humoral immune defences in mammals and insects: The same, with differences? *Virulence* 9, 1625-1639.
- Shih, L., Tsai, K.L., Hsieh, C., 2007. Effects of culture conditions on the mycelial growth and bioactive metabolite production in submerged culture of *Cordyceps militaris*. *Biochem.Eng. J.* 33, 193-201.
- Shrestha, B., Zhang, W., Zhang, Y., Liu, X., 2012. The medicinal fungus *Cordyceps militaris*: research and development. *Mycol. Prog.* 11, 599-614.
- Sowa-Jasiłek, A., Zdybicka-Barabas, A., Stączek, S., Wydrych, J., Skrzypiec, K., Mak, P., Deryło, K., Tchórzewski, M., Cytryńska, M., 2016. *Galleria mellonella* lysozyme induces apoptotic changes in *Candida albicans* cells. *Microbiol. Res.* 193, 121-131.
- Stokes, B.A., Yadav, S., Shokal, U., Smith, L.C., Eleftherianos, I., 2015. Bacterial and fungal pattern recognition receptors in homologous innate signalling pathways of insects and mammals. *Front. Microbiol.* 6, 19.
- Strader, C.R., Pearce, C.J., Oberlies, N.H., 2011. Fingolimod (FTY720): a recently approved multiple sclerosis drug based on a fungal secondary metabolite. *J. Nat. Prod.* 74, 900-907.
- Strober, W., 2001. Trypan blue exclusion test of cell viability. *Curr. Protoc. Immunol.* Appendix 3B.
- Survase, S.A., Kagliwal, L.D., Annapure, U.S., Singhal, R.S., 2011. Cyclosporin A—a review on fermentative production, downstream processing and pharmacological applications. *Biotechnol. Adv.* 29, 418-435.
- Tanada, Y., Kaya, H.K., 1993. *Insect pathology*. Academic press.
- Tanji, T., Ip, Y.T., 2005. Regulators of the Toll and Imd pathways in the *Drosophila* innate immune response. *Trends Immunol.* 26, 193-198.
- Tuli, H.S., Sandhu, S.S., Sharma, A.K., 2014. Pharmacological and therapeutic potential of *Cordyceps* with special reference to Cordycepin. *3 Biotech*, 4, 1-12.
- U.S Department of Agriculture 2016. Agricultural Research Service, Biological Integrated Pest management Research Unit. *ARS Collection of Entomopathogenic Fungal Cultures (ARSEF)*. U.S. Department of Agriculture, Agricultural Research Service.
- Vega, F.E., Goettel, M.S., Blackwell, M., Chandler, D., Jackson, M.A., Keller, S., Koike, M., Maniania, N.K., Monzón, A., Ownley, B.H., Pell, J.K., Rangel, D.E.N., Roy, H.E., 2009. Fungal entomopathogens: new insights on their ecology. *Fungal Ecol.* 2, 149-159.
- Vertyporokh, L., Wojda, I., 2017. Expression of the insect metalloproteinase inhibitor IMPI in the fat body of *Galleria mellonella* exposed to infection with *Beauveria bassiana*. *Acta Biochim. Pol.* 64, 273-278.

- Vey, A., Matha, V., Dumas, C., 2002. Effects of the peptide mycotoxin destruxin E on insect haemocytes and on dynamics and efficiency of the multicellular immune reaction. *J. Invertebr. Pathol.* 80, 177-187.
- Vilcinskas, A., 2019. Evolutionary ecology of parasitic fungi and their host insects. *Fungal Ecol.* 38, 12-20.
- Vilcinskas, A., Matha, V., Gotz, P., 1997a. Effects of the entomopathogenic fungus *Metarhizium anisopliae* and its secondary metabolites on morphology and cytoskeleton of plasmatocytes isolated from *Galleria mellonella*. *J. Insect Physiol.* 43, 1149-1159.
- Vilcinskas, A., Matha, V., Gotz, P., 1997b. Inhibition of phagocytic activity of plasmatocytes isolated from *Galleria mellonella* by entomopathogenic fungi and their secondary metabolites. *J. Insect Physiol.* 43, 475-483.
- Vilcinskas, A., Wedde, M., 2002. Insect inhibitors of metalloproteinases. *IUBMB Life* 54, 339-343.
- Wang, C., Leger, R.J.S., 2006. A collagenous protective coat enables *Metarhizium anisopliae* to evade insect immune responses. *Proc. Nat. Acad. Sci. USA* 103, 6647-6652.
- Warton, D.I., Hui, F.K., 2011. The arcsine is asinine: the analysis of proportions in ecology. *Ecology* 92, 3-10.
- Webb J.L., 1963. General Principles of Inhibition. In *Enzyme and Metabolic Inhibitors, Volume 1*. Academic Press: New York, USA. p. 949.
- Wedde, M., Weise, C., Nuck, R., Altincicek, B., Vilcinskas, A., 2007. The insect metalloproteinase inhibitor gene of the lepidopteran *Galleria mellonella* encodes two distinct inhibitors. *Biol. Chem.* 388, 119-127.
- Wojda, I., Jakubowicz, T., 2007. Humoral immune response upon mild heat-shock conditions in *Galleria mellonella* larvae. *J. Insect Physiol.* 53, 1134-1144.
- Wojda, I., Kowalski, P., Jakubowicz, T., 2009. Humoral immune response of *Galleria mellonella* larvae after infection by *Beauveria bassiana* under optimal and heat-shock conditions. *J. Insect Physiol.* 55, 525-531.
- Woolley, V.C., 2018. Elucidating the natural function of cordycepin, a secondary metabolite of the fungus *Cordyceps militaris*, and its potential as a novel biopesticide in Integrated Pest Management. Ph.D. Thesis. University of Warwick.
- Xia, Y., Luo, F., Shang, Y., Chen, P., Lu, Y., Wang, C., 2017. Fungal cordycepin biosynthesis is coupled with the production of the safeguard molecule pentostatin. *Cell Chem. Biol.* 24, 1479-1489.
- Yang, F., Li, S., 2008. Effects of sample preparation methods on the quantification of nucleosides in natural and cultured *Cordyceps*. *J. Pharmaceut. Biomed.* 48, 231-235.


- Yang, X., Li, Y., He, Y., Li, T., Wang, W., Zhang, J., Wei, J., Deng, Y., Lin, R., 2015. Cordycepin alleviates airway hyperreactivity in a murine model of asthma by attenuating the inflammatory process. *Int. Immunopharmacol.* 26, 401-408.
- Yi, H.Y., Chowdhury, M., Huang, Y.D., Yu, X.Q., 2014. Insect antimicrobial peptides and their applications. *Appl. Microbiol. Biot.* 98, 5807-5822.
- Zhang, D.W., Wang, Z.L., Qi, W., Lei, W., Zhao, G.Y., 2014. Cordycepin (3'-deoxyadenosine) down-regulates the proinflammatory cytokines in inflammation-induced osteoporosis model. *Inflammation* 37, 1044-1049.
- Zimmermann, G., 2007a. Review on safety of the entomopathogenic fungi *Beauveria bassiana* and *Beauveria brongniartii*. *Biocontrol Sci. Technol.* 17, 553-596.
- Zimmermann, G., 2007b. Review on safety of the entomopathogenic fungus *Metarhizium anisopliae*. *Biocontrol Sci. Technol.* 17, 879-920.



**Fig. 1- Expression of AMPs in *D. melanogaster* S2r+ cells in response to cordycepin treatment.** S2r+ cells (three biological replicates) were treated with DMSO (dark grey bars) or DMSO plus 25  $\mu\text{g ml}^{-1}$  (100  $\mu\text{M}$ ) cordycepin (light grey bars). Immune challenge was simulated using 20  $\mu\text{g ml}^{-1}$  LPS or curdlan. Cells were sampled 4 h after treatment. Fold changes in gene expression were calculated using the  $2^{-\Delta\Delta\text{CT}}$  method (Schmittgen and Livak, 2008) compared to untreated samples with correction for the internal control (*RP49*). Relative expression of A) *metchnikowin* and B) *attacin* A is shown. Error bars show  $\pm$  SEM and different lowercase letters indicate significant differences ( $p < 0.05$ ; ANOVA, Tukey HSD post hoc tests).




**Fig. 2- Proportion mortality of *G. mellonella* larvae following cordycepin treatment.** Mean proportional mortality of *G. mellonella* larvae (N=30 for each dose) injected with 1, 1.82, 3, 6 and 11 mg ml<sup>-1</sup> of cordycepin and a DMSO control on day 6 after treatment. A type 1 Weibull model is fitted to the response.







**Fig. 3- Proportion mortality of *G. mellonella* larvae following treatment with EPF.** Mean proportional mortality of *G. mellonella* larvae (N=30 for each dose) injected with *B. bassiana* and *C. militaris* conidia at six concentrations on day 6 after treatment. A two-parameter log-logistic model is fitted to the responses.



**Fig. 4- The impact of cordycepin on EPF germination.** Mean (back-transformed) percentage germination of *B. bassiana* 433.99, *C. militaris* 11703, and *M. brunneum* 275.86 conidia  $\pm$ SE following incubation for 24 h with 0 (dark grey) and 3 (light grey) mg ml<sup>-1</sup> of cordycepin.



**Fig. 5 – Survival curves of *G. mellonella* treated with cordycepin and EPF.** Survival of *G. mellonella* larvae injected with 10 or 100 *B. bassiana* conidia or *C. militaris* conidia. Solid lines show observed mortalities, those lines ending in ‘+’ indicate censored populations. Dashed lines indicate the expected population decline (shaded area illustrates the standard error) estimated by fitting a binomial generalized logistic model. Pale grey lines indicate treatment with cordycepin (3 mg ml<sup>-1</sup>) and dark grey with a DMSO control. N=30 per treatment. P values are the result of a log-rank test.



**Fig. 6- Expression of immune-related genes in response to cordycepin and *C. militaris* or *B. bassiana* 48 h and 72 h after treatment.** The expression of *galiomycin*, *gallerimycin*, *IMPI* and *lysozyme* in response to stimulation by *C. militaris* or *B. bassiana* injection and sham (Triton X-100) injection 48 h and 72 h. Fold changes in gene expression were calculated using the  $2^{-\Delta\Delta CT}$  method (Schmittgen and Livak, 2008) compared to untreated samples with correction for the internal control (*S7e*). Treatment without cordycepin is shown by dark grey bars and with cordycepin ( $3 \text{ mg ml}^{-1}$ ) by light grey bars, expression in uninjected insects is also included.  $N=13-15$  per treatment. Different letters above bars indicate statistically significant differences ( $p < 0.05$ ) identified using the Dunn-Bonferroni post hoc method following a Kruskal-Wallis test.

**Table 1- Primers used to monitor immune-related gene expression in *G. mellonella* and *D. melanogaster*.**

<b>Gene</b>	<b>Primer sequence (5'-3')</b>	<b>Reference</b>
<i>gallerimycin</i>	TATCATTGGCCTTCTTGGCTG GCACTCGTAAAATACACATCCGG	Wojda and Jakubowicz, 2007
<i>galiomicin</i>	TCGTATCGTCACCGCAAATG GCCGCAATGACCACCTTTATA	Wojda et al., 2009
<i>lysozyme</i>	TCCCAACTCTTGACCGACGA AGTGGTTGCGCCATCCATAC	Altincicek and Vilcinskas, 2006
<i>IMPI</i>	AGATGGCTATGCAAGGGATG AGGACCTGTGCAGCATTCT	Altincicek and Vilcinskas, 2006
<i>S7e</i>	ATGTGCCAATGCCCAAGTTG GTGGCTAGGCTTGGGAAGAAT	Wojda and Jakubowicz, 2007
<i>attacin A</i>	AGGTCCTTAACCTCCAATC CATGACCAGCATTGTTGTAG	Jin et al., 2008
<i>metchnikowin</i>	TCTTGGAGCGATTTTTCTGG AATAAATTGGACCCGGTCTTG	Castillo et al., 2013
<i>RP49</i>	GACGCTCAAGGGACAGTATCTG AAACGCGGTTCTGCATGAG	Gobert et al., 2003