

# New Phytologist

## **Pollen sterols are associated with phylogenetics and environment but not with pollinators**

Journal:	<i>New Phytologist</i>
Manuscript ID	NPH-MS-2020-34747.R1
Manuscript Type:	MS - Regular Manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	Zu, Pengjuan; University of Zurich, Geology; Royal Botanic Gardens Kew, Natural Capital and Plant Health Koch, Hauke; Royal Botanic Gardens Kew, Natural Capital and Plant Health Schwery, Orlando; New Mexico Consortium, Plant biology Pironon, Samuel; Royal Botanic Gardens Kew, Biodiversity Informatics and Spatial Analysis Phillips, Charlotte; Royal Botanic Gardens Kew, Natural Capital and Plant Health Ondo, Ian; Royal Botanic Gardens Kew, Biodiversity Informatics and Spatial Analysis Farrell, Iain; Royal Botanic Gardens Kew, Natural Capital and Plant Health Nes, David; Texas Tech University, Dept of Chemistry & Biochemistry Moore, Elynor; Oxford University, Department of Zoology Wright, Geraldine; Oxford University, Department of Zoology Farman, Dudley; University of Greenwich, Natural Resources Institute Stevenson, Phillip; Royal Botanic Gardens, Jodrell Laboratory,; University of Greenwich, Natural Resources Institute
Key Words:	Phytosterol diversity, Pollen nutrient, Pollinator assemblages, Environmental factors, Phylogenetic, Plant-insect interactions, Chemical ecology, Chemotaxonomy

SCHOLARONE™  
Manuscripts

1 **Pollen sterols are associated with phylogenetics and environment but not with**  
2 **pollinators**

3

4 Pengjuan Zu<sup>1,2\*</sup>, Hauke Koch<sup>1</sup>, Orlando Schwery<sup>3</sup>, Samuel Pironon<sup>1</sup>, Charlotte  
5 Phillips<sup>1</sup>, Ian Ondo<sup>1</sup>, Iain W. Farrell<sup>1</sup>, W. David Nes<sup>4</sup>, Elynor Moore<sup>5</sup>, Geraldine A.  
6 Wright<sup>5</sup>, Dudley I. Farman<sup>6</sup>, Philip C. Stevenson<sup>1,6</sup>

7

8 <sup>1</sup> Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, UK

9 <sup>2</sup> Swiss Federal Institute of Aquatic Science and Technology, Seestrasse 79, CH-  
10 6047, Kastanienbaum, Switzerland

11 <sup>3</sup> New Mexico Consortium, 4200 W. Jemez Rd, Suite 301, Los Alamos, NM 87544,  
12 USA

13 <sup>4</sup> Texas Tech University, Lubbock, Texas, USA

14 <sup>5</sup> University of Oxford, Oxford, UK

15 <sup>6</sup> Natural Resources Institute, University of Greenwich, Chatham, Kent, ME4 4TB, UK

16 \*Correspondence author: [Pengjuan.zu@gmail.com](mailto: Pengjuan.zu@gmail.com)

17

## 18 **Summary**

- 19 • Phytosterols are primary plant metabolites that have fundamental structural and  
20 regulatory functions. They are also essential nutrients for phytophagous insects,  
21 including pollinators, that cannot synthesize sterols. Despite the well-described  
22 composition and diversity in vegetative plant tissues, few studies have examined  
23 phytosterol diversity in pollen.
- 24 • We quantified 25 pollen phytosterols in 122 plant species (105 genera, 51 families)  
25 to determine their composition and diversity across plant taxa. We searched  
26 literature and databases for plant phylogeny, environmental conditions, and  
27 pollinator guilds of the species to examine the relationships with pollen sterols.
- 28 • 24-methylenecholesterol, sitosterol and isofucosterol were the most common and  
29 abundant pollen sterols. We found phylogenetic clustering of twelve individual  
30 sterols, total sterol content and sterol diversity, and of sterol groupings that reflect  
31 their underlying biosynthesis pathway (24 carbon alkylation, ring B desaturation).  
32 Plants originating in tropical-like climates (higher mean annual temperature, lower  
33 temperature seasonality, higher precipitation in wettest quarter) were more likely  
34 to record higher pollen sterol content. However, pollen sterol composition and  
35 content showed no clear relationship with pollinator guilds.
- 36 • Our study is the first to show that pollen sterol diversity is phylogenetically  
37 clustered and that pollen sterol content may adapt to environmental conditions.
- 38
- 39

## 40 **Introduction**

41 Phytosterols are a class of lipids with key metabolic and ecological functions for plants  
42 (Nes & McKean, 1977; Vanderplanck *et al.*, 2020a). For example, they regulate  
43 membrane fluidity and permeability (Grunwald, 1971; Schuler *et al.*, 1991; Hartmann,  
44 1998), and act as precursors for metabolic signals such as brassinosteroid growth  
45 hormones (Grove *et al.*, 1979; Chung *et al.*, 2010) that promote cell division, and  
46 mediate reproduction in plants and protect them against environmental stresses  
47 (Khripach *et al.*, 2000). Phytosterols may also modulate plant defence against  
48 bacterial pathogens (Posé *et al.*, 2009; Wang *et al.*, 2012; Ferrer *et al.*, 2017) and  
49 pollen sterols accelerate germination and tube growth and protect against desiccation  
50 (Kumar *et al.*, 2015; Rotsch *et al.*, 2017).

51 Phytosterols show considerable diversity with more than 250 structures reported (Nes,  
52 2011) with notable variation at the methine substitution (double bond) in ring B and  
53 methyl or ethyl substitutions at C-24 (Fig. 1). The structural variation and composition  
54 of sterols in plant tissues is important for phytophagous insects since they cannot  
55 synthesize sterols *de novo*, and therefore depend upon specific plants to obtain the  
56 required sterols from their diet to sustain their development (Behmer & Elias, 1999,  
57 2000; Lang *et al.*, 2012). This may be especially important for pollen feeding insects  
58 that require specific sterols. Honeybees, for example, require 24-methylenecholesterol  
59 (Herbert *et al.* 1980; Chakrabati *et al.*, 2020) so must collect pollen from plant species  
60 that produce this sterol to rear brood. Bee sterols are similar to those occurring in the  
61 pollen on which they feed (Vanderplanck *et al.*, 2020a) but differ across bee taxa  
62 suggesting bees are what they eat with respect to sterols. Wild pollinators range from  
63 pollen generalists to specialists (Rasmussen *et al.*, 2020) and for some species, this  
64 specialism may be mediated by pollen sterols. Therefore, a landscape of flowers that  
65 does not provide the sterols required for a specific bee may be nutritionally deficient  
66 for that species. In general, however, the relationships between pollen sterols and the  
67 nutritional needs of pollination insects has not yet been evaluated.

68 Conversely, plant sterol composition may evolve with antagonists as well as mutualists  
69 since the pathways for the synthesis of sterols overlap with that for some defence  
70 compounds against herbivores (Qi *et al.*, 2006). A range of naturally occurring insect  
71 toxins occur in pollen (Arnold *et al.*, 2014; Rivest & Forrest, 2019) with the likely role  
72 of protecting the male gamete and since some sterols can also act as defensive

73 compounds against arthropod herbivores (Jing & Behmer, 2020). They could also be  
74 toxic to pollen feeding insects to reduce damage to or excessive loss of pollen grains.

75 Abiotic conditions may affect phytosterol structural variations at different levels. At  
76 plant individual level, a 24-ethyl substitution (e.g., sitosterol and stigmasterol), for  
77 example, reinforces membrane cohesion (Piironen, 2000; Dufourc, 2008), and  
78 therefore sterol structures may be altered in response to temperature variations. At  
79 population level, from limited heritability studies on phytosterols in plant seeds (Amar  
80 *et al.*, 2008, Velasco *et al.*, 2013), environmental factors also contribute to sterol  
81 phenotypic variation, although much less compared to the contribution from genetic  
82 factors (heritabilities above 0.8 were documented). At species level, pollen sterol  
83 composition seems to be highly variable between different species (Villette *et al.*,  
84 2015; Vanderplanck *et al.*, 2020a) and can differ from vegetative tissues (Nes, 1990;  
85 Nes *et al.*, 1993). However, no study has investigated whether pollen sterol variations  
86 at species level can be the consequences of evolutionary adaptation to environmental  
87 conditions.

88 Moreover, due to the limited number of studies on pollen sterol profiles, we lack a  
89 comprehensive and fundamental understanding of the patterns of pollen sterol  
90 diversity across plant taxa. It is still controversial whether pollen phytosterols are  
91 phylogenetically structured. For example, Standifer *et al.* (1968) suggested a lack of  
92 phylogenetic constraints of pollen sterol composition based on the evidence of large  
93 variation in three Salicaceae species. Vanderplanck *et al.* (2020a), in contrast, found  
94 similar pollen sterol composition within the genus *Salix* and our interpretation of the  
95 data published by Villette *et al.* (2015) suggested the occurrence of some pollen  
96 sterols was phylogenetically constrained. Since most studies focused on a few plant  
97 species, they were insufficient to reach a general overview of the patterns of pollen  
98 sterol diversity across plant taxa and their drivers.

99 In this study, we analysed pollen sterols including saturated stanols in 122  
100 angiosperms representing 51 plant families and 25 plant orders. We further compiled  
101 data from literature and databases on plant phylogeny, pollinators, and environmental  
102 conditions within native geographic regions for each plant species to examine  
103 relationships between these factors and pollen sterol composition and diversity.  
104 Specifically, we ask 1) Are pollen sterols phylogenetically clustered? 2) Are pollen

105 sterols correlated with abiotic environments? 3) Are pollen sterols associated with  
106 pollinator guilds?

107

## 108 **Materials and methods**

### 109 **Pollen collection**

110 From March to November 2018, we collected pollen from fresh flowers growing in the  
111 Royal Botanic Gardens (RBG), Kew, UK and nearby areas (see Table S1 for details  
112 of collection dates and locations for each species). RBG Kew supports a diverse  
113 collection of living plant species from across the world. Prior to pollen collection, we  
114 used a fine-meshed bag to cover flower buds whenever possible to prevent potential  
115 contamination or removal due to pollinator visitation. When flowers were fully open,  
116 we gently shook the flower and collected pollen into a weighed 2 mL microcentrifuge  
117 tube (Eppendorf®, Safe-Lock™). For species for which pollen was more difficult to  
118 harvest, such as in the cases of *Lamium purpureum* L. and *Ulex europaeus* L., we  
119 used small forceps to help push the pollen out or trigger pollen ejection, respectively.  
120 Pilot studies carried out in our laboratory (with *Helleborus foetidus*, *Prunus avium*,  
121 *Prunus spinosa*, *Salix cinerea* and *Symphytum officinale*) showed a conserved pattern  
122 of pollen sterol composition: within species variation was significantly lower than  
123 between species variation (all p-values < 0.001 under multi-variate distribution tests  
124 e.g., Hotelling test, Pillai test, and Wilks' lambda distribution test), consistent with  
125 findings on within vs. between species variation of other pollen metabolites (Palmer-  
126 Young *et al.*, 2019). Therefore, we collected 2 to 5 replicates per species (details see  
127 Table S1) and used the average quantities across replicates of each species for  
128 analyses. In total, we collected 308 samples from 122 species, representing 105  
129 genera, 51 families and 25 orders across the major groups of seed plants  
130 (Gymnosperms, Nymphaeales, Monocots, Ranunculids, Caryophyllales, Asterids and  
131 Rosids; Table S1). Our selection of species was guided by a combination of practical  
132 considerations (feasibility to collect sufficient pollen for analysis, availability of species  
133 at Kew) while attempting to maximize phylogenetic and ecological diversity of plants  
134 (pollinator guilds, ecological niches). Pollen weight (to 0.1 mg accuracy) and collection  
135 date were recorded for each sample. Pollen samples were stored in a freezer (-20°C)  
136 before extracting sterols.

## 137 **Sterol content analysis**

138 To extract sterols and stanols (from here referred to as phytosterols or pollen sterols)  
139 from the pollen, we added 500  $\mu\text{l}$  10% KOH in MeOH to the microcentrifuge tubes  
140 containing a weighed pollen sample. Then, an internal standard (20  $\mu\text{l}$  of 0.5  $\text{mg ml}^{-1}$   
141 epicoprostanol) was added prior to incubating the tube for 2 h at 80°C for  
142 saponification. Phytosterols were then recovered into 1 mL hexane. After complete  
143 evaporation of hexane, phytosterols remained in the tube. We derivatized these with  
144 20  $\mu\text{l}$  Tri-Sil (Sigma, Gillingham, Dorset UK) and then briefly vortexed and injected  
145 directly into an Agilent Technologies (Palo Alto, CA, USA) 7890A gas chromatograph  
146 connected to an Agilent Technologies 5975C MSD mass spectrometer (GC-MS) and  
147 eluted over an Agilent DB5 column using a splitless injection at 250°C with a standard  
148 GC program at 170°C for 1 minute ramped to 280°C at 20°C per minute and monitoring  
149 between 50 and 550 amu.

150 All 25 phytosterols were identified by comparison of their retention time relative to  
151 cholesterol and mass spectra from authentic standards (David W Nes collection,  
152 details see Fig. S4 for mass spectra of each sterol) either directly through co-analysis  
153 or using existing data and confirmed where data was available with the NIST (National  
154 Institute of Standards and Technology) mass spectral library (Guo *et al.*, 1995; Heupel  
155 and Nes, 1984; Nes *et al.*, 1977; Xu *et al.*, 1988; Zhou *et al.*, 2009; Nes *et al.*, 2003).

156 To quantify the amount of each phytosterol, we used its relative peak area by  
157 calculating the ratio of the peak area of the targeted sterol to that of the internal  
158 standard. Then, by multiplying the ratio with the mass of the internal standard, we  
159 obtained the mass of each sterol in the sample. Compound identification (using target  
160 ion) and quantification were carried out with ChemStation Enhanced Data Analysis  
161 (Version E.01.00). In total, we identified 25 phytosterols in pollen (Table S1).

162 For each plant species, we calculated each phytosterol amount ( $\mu\text{g}$  per mg sampled  
163 pollen), total sterol content ( $\mu\text{g}$  per mg sampled pollen), and the percentage of each  
164 sterol in total phytosterol content. In addition, we calculated the chemical diversity  
165 index using Shannon entropy: where  $S$  is the total number of phytosterols,  $p_i$  is the  
166 percentage of the  $i^{\text{th}}$  phytosterol. Note that we used the total phytosterol number  $S$  as  
167 the base of log (instead of the natural base  $e$ ) to scale the range of diversity index  
168 values to  $[0, 1]$  with 1 indicating the highest diversity. This equates to calculating  
169 Shannon's equitability. Finally, for each phytosterol, we calculated its commonness

170 and abundance across all plant species. Commonness is given by the proportion of  
171 plant species that contained that specific phytosterol (i.e., present/absent). Relative  
172 abundance was given by the average proportion of a specific sterol across all species.

173 Additionally, to understand how different phytosterol in pollen co-varied, we performed  
174 a factor analysis using the R package *stats* (R Core Team, 2020) on the data for the  
175 absolute weight of phytosterols measured in pollen across the entire data set. We set  
176 a criterion of eigenvalue  $> 1$  for inclusion of extracted factors. A varimax rotation was  
177 used to adjust the fit of the factor analysis to variance in the data.

178 Moreover, based on biosynthetic reasoning as discussed by Benveniste (2004), we  
179 arranged these phytosterols identified in our pollen samples into alternate hypothetical  
180 biosynthetic pathways to cholesterol and 24-alkyl phytosterols.

### 181 **Phylogenetic tree construction and analyses**

182 We used the R package *rotl* (Michonneau *et al.*, 2016) to download the induced  
183 subtree of only our focal taxa from the Open Tree of Life (OTL) synthetic tree (Hinchliff  
184 *et al.*, 2015; Rees *et al.*, 2017). If only the genus was known, OTL used the root of the  
185 genus for the subtree wherever possible. Name synonyms and corrections suggested  
186 by OTL for genus and species were adopted in our analyses (see Table S2). Taxa  
187 with subspecies or other epithets beyond species level were reduced to genus and  
188 species only (*Amaryllis belladonna* L., *Campanula fragilis* Cirillo, *Campanula isophylla*  
189 Moretti, *Euphorbia milii* Des Moul., *Hieracium umbellatum* L.). Only one terminal was  
190 retained to represent the two differently coloured varieties of *Hymenocallis littoralis*  
191 (Jacq.) Salisb.

192 We estimated divergence times with penalised likelihood using nine secondary  
193 calibration points. Using the R package *ape* (Paradis *et al.*, 2004), we randomly  
194 resolved polytomies and computed branch lengths using Grafen's method. We looked  
195 up the inferred ages of seven clades from the large phylogeny of spermatophytes by  
196 Zanne *et al.*, (2014): Angiospermae (243 million years ago [mya]), Monocotyledoneae  
197 (171 mya), Eudicotyledoneae (137 mya), Superrosidae (118 mya), Rosidae (117  
198 mya), Superasteridae (117 mya), and Asteridae (108 mya). The age of Nymphaea  
199 (78.07 mya) was obtained from DateLife (Sanchez-Reyes, 2019), and we took the  
200 estimated origin of Spermatophyta at 327 mya (Smith *et al.*, 2010) to calibrate the root  
201 age. We used those times as minimal age constraints for a penalized likelihood

202 analysis using *chronopl* in *ape* (Paradis *et al.*, 2004). Monophyly of families was  
203 checked using *MonoPhy* (Schwery & O'Meara, 2016).

204 To determine whether there is phylogenetic structure in the pollen sterol data, we used  
205 the function *phyloSignal* from the R package *phyloSignal* (Keck *et al.*, 2016) to  
206 calculate Pagel's  $\lambda$  (Pagel, 1999) and Blomberg's K (Blomberg *et al.*, 2003), each with  
207 999 iterations for *p*-value estimation. Phylogenetic signal was estimated this way for  
208 each of the individual sterol compounds (based on their percentage value), for sums  
209 of compounds belonging to each C-24 substitution (C0, C1, C2 indicating substitution  
210 with no carbon, a methyl and an ethyl), and position of the olefinic or methine moiety  
211 in ring B ( $\Delta^0$ ,  $\Delta^5$ ,  $\Delta^7$ ,  $\Delta^8$ ), for the sterol diversity index H, and for the total phytosterol  
212 content (absolute sterol amount per mg pollen). The output of these analyses was  
213 visualized using the R packages *phytools* (Revell 2012) and *picante* (Kembel *et al.*  
214 2010).

#### 215 **Plant occurrence records and abiotic environmental data**

216 To investigate whether species-level variations in pollen sterols are partially the  
217 consequences of evolutionary adaptation to environmental conditions, we retrieved  
218 environmental information of the native geographic ranges of each species. Note that  
219 here we focused on "long-term" prevailing abiotic conditions (e.g., climate) capable of  
220 shaping evolutionary changes of sterol composition at species level, as opposed to  
221 "short-term" abiotic variables (e.g., stresses, weather) affecting traits via phenotypic  
222 plasticity at the individual level. For each species, we extracted geographic occurrence  
223 records from several global and continental databases: GBIF (Global Biodiversity  
224 Information Facility; <https://www.gbif.org/>) using the *rgbif* package in R, BIEN  
225 (Botanical Information and Ecology Network; <http://bien.nceas.ucsb.edu/bien/>) using  
226 the *BIEN* R package, BioTIME (Dornelas *et al.*, 2018) and Rainbio (Dauby *et al.*,  
227 2016). Because raw occurrence data from these databases contain taxonomic, spatial  
228 and temporal inconsistencies (Meyer *et al.*, 2016), we applied different cleaning filters  
229 using the *CoordinateCleaner* package in R (Zizka *et al.*, 2019). We discarded non-  
230 georeferenced records, records with latitude and longitude given as zero and having  
231 equal longitude and latitude, points recorded before 1950, as well as fossil data,  
232 records corresponding to centroids of countries, capitals, known botanical institutions  
233 and GBIF headquarters, occurrences falling in the sea, cultivated records, and points  
234 indicated as having high coordinate uncertainty (>20 km). For each species, we used

235 the World Geographical Scheme for Recording Plant Distribution (WGSRPD)  
236 database (Brummitt, 2001), developed by the International Working Group on  
237 Taxonomic Databases for Plant Sciences (TDWG), to only retain occurrence records  
238 located inside its native range defined at subcontinental scale (level-2 of TDWG).  
239 Finally, we removed duplicates and thinned each species' occurrence dataset by  
240 keeping only one record by 10x10 arc-min grid cell to limit spatial autocorrelation.  
241 Finally, we discarded records from species introduced ranges not intersecting native  
242 regions at the level-2 of the World Geographical Scheme for Recording Plant  
243 Distribution (regional or sub-continental level) (Brummitt, 2001) from the World  
244 Checklist of Vascular Plants (WCVP, 2020). In total, 355,912 occurrence records were  
245 retrieved across all species (Table S1).

246 We quantified species environmental niches based on a set of 13 climate, soil, and  
247 topography variables. Eight of them were bioclimatic variables (BIO1, BIO4, BIO10,  
248 BIO11, BIO12, BIO15, BIO16 and BIO17) extracted from the CHELSA database  
249 (Karger *et al.*, 2017), representing annual mean, seasonality, minimum and maximum  
250 temperature and precipitation (full list of variables and descriptions see Table S3).  
251 Four soil variables were extracted from the SoilGrids database (ISRIC, 2013;  
252 <http://www.data.isric.org>) and averaged across a 0-60 cm depth gradient: depth to  
253 bedrock, mean soil organic carbon stock, pH and water capacity. Land slope was  
254 calculated using the Slope function in the Spatial Analyst toolbox of ARC/INFO GIS  
255 based on the Global Multi-resolution Terrain Elevation Database (GMTED) (Danielson  
256 *et al.*, 2011). To match the resolution of the occurrence records, all environmental  
257 variables were upscaled to 10 arc-min (ca. 20 km) using the *resample* function of the  
258 *raster* package in R.

259 We extracted each of the 13 environmental variables at each occurrence point of each  
260 species using the *extract* function of the *raster* package in R. Mean environmental  
261 conditions were then calculated for each of the 13 variables across all occurrences of  
262 each species (i.e., environmental niche position along individual environmental  
263 gradients). We also created an environmental space summarizing the variation in the  
264 13 environmental variables across the world using a Principal Component Analysis  
265 (PCA) and the function *princomp* in the *stats* package in R. We kept the first three  
266 component axes that explained 74% of the variation in the 13 variables: PC1 being  
267 mainly positively correlated with mean temperatures and negatively correlated with

268 temperature seasonality and soil carbon content, PC2 being positively correlated with  
269 soil pH and negatively with precipitation, and PC3 being positively correlated with soil  
270 depth to bedrock and negatively with land slope (see Table S3 for variable  
271 contributions to PCA axes). To quantify the niche breadth of each species, we first  
272 drew three-dimensional alpha shapes around each set of occurrence points of each  
273 species in the environmental space defined by the PCA with an alpha value of two  
274 using the *ashape3d* function in the *alphashape3d* package in R (Capinha & Pateiro-  
275 López, 2014). The alpha-shape is a profile technique used to compute species  
276 environmental niche envelopes using a flexible envelope fitting procedure that does  
277 not make any assumption about the shape of the niche (Capinha & Pateiro-López,  
278 2014). We then calculated the volume of each species' alpha shape as a measure of  
279 their environmental niche breadth using the *volume\_ashape3d* function from the latter  
280 package. We also calculated the mean position of each species' alpha shape on the  
281 three retained main axes of the PCA (i.e., niche position, individual variable  
282 contributions see Table S3). Because three-dimensional alpha shapes require at least  
283 five occurrence points to be drawn, species with fewer records were discarded. We  
284 also discarded those species lacking sufficient and reliable geographic data or  
285 taxonomic uncertainty (i.e., we did not extract occurrence records for genera,  
286 subspecies and hybrids). In the end, we quantified niche breadth for 90 species, while  
287 32 taxa were discarded, and niche position for 100 species (22 taxa discarded; details  
288 see Table S1).

### 289 **Pollinator data collection**

290 To study whether there is a relationship between plants' pollen sterols and their  
291 pollinators, we categorized plants in two different ways. Firstly, based on pollinator  
292 guilds, as 1) Bee, 2) Fly, 3) Lepidoptera, 4) Thrips, 5) Generalist insect, 6) Bird, or 7)  
293 Wind pollinated. Secondly, we grouped plants by whether or not pollen acts as a  
294 reward for bee pollinators. On the one hand, bees depend on pollen as larval food and  
295 require pollen sterols as essential nutrients. Plants could therefore hypothetically  
296 attract bee pollinators with pollen sterol profiles of high nutritional quality to them. On  
297 the other hand, if pollen does not play a role as bee reward (i.e., in non-bee pollinated  
298 plants, and/or where nectar is the sole reward), sterol profiles could be expected that  
299 are of low quality or even toxic to bees to prevent pollen robbery (as shown for some  
300 other chemical compounds in pollen, Rivest & Forrest 2019).

301 To classify plant pollinator guilds and groups, we conducted literature searches for  
302 each plant species on Google Scholar, using the scientific name (including common  
303 synonyms) and “pollinat\*”, OR “pollen”, OR “flower” as search terms. We examined  
304 relevant cited or citing references of publications found in this way for additional  
305 records, and consulted Knuth (1908, 1909) and Westrich (2018), or personal  
306 observations. If no sources on pollination and flower visitation were available, the  
307 pollinator guild was classified as “unknown” (10 species in data set). We included  
308 plant species in the “pollen as bee reward” group that both receive pollination services  
309 by bees (including some plants in the “generalist insect pollination” category) and have  
310 records of bees collecting pollen. Plants were classified as not producing pollen as  
311 bee reward if they were either not pollinated by bees, or, in case of bee pollination,  
312 had clear evidence of pollen not being collected by bees (e.g., pollen contained in  
313 pollinia of bee-pollinated orchids). Plants for which data on pollinator guild and  
314 collection of pollen by bees was missing were classified as “unknown” (34 species in  
315 data set). A full list of relevant references and the assigned pollinator guilds is provided  
316 in Table S1.

### 317 **Analyses on relationships between phytosterols and (a)biotic factors**

318 To assess the association of sterol composition with environmental variables and  
319 pollinator guilds, we first calculated a Bray-Curtis distance matrix for sterol profiles of  
320 pairwise plant species comparisons, using absolute weights ( $\mu\text{g}$ ) of each sterol per mg  
321 pollen. Then we related this distance matrix to environmental factors and pollinator  
322 guild to study their relationships. Specifically, for abiotic environmental factors  
323 (continuous values), we ran MRM (multiple regression on distance matrices) analyses  
324 (Lichstein, 2007) using an additive linear model with the Bray-Curtis distance matrix  
325 of pollen sterol composition dissimilarity as response, and environmental niche  
326 distance matrices for PC1, PC2, PC3 (see “abiotic environmental data” above,  
327 calculated from pairwise Euclidean distances for all plants for their position on each of  
328 the PCs) and a phylogenetic distance matrix (pairwise phylogenetic distance in mya,  
329 phylogeny see above) as independent variables. We used Pearson correlations with  
330 10000 permutations to test for significant associations. Calculations of distance  
331 matrices and MRM analyses were done with the R package *ecodist* (Goslee & Urban,  
332 2007). For pollinator modes (categorical variables), we conducted ANOSIMs  
333 (analyses of similarities; Clarke, 1993) to test for significant differences of pollen sterol

334 profiles between different pollinator groups (excluding pollinator groups with only one  
335 representative, i.e. wind, thrips, and fly) or between plant groups where pollen is used  
336 as reward by bees or not. ANOSIMs were conducted in PAST 4.03 (Hammer *et al.*,  
337 2001) with 10000 permutations. We illustrated the relationship of these factors to sterol  
338 profile similarity with 2D non-metric multidimensional scaling (NMDS) ordination plots  
339 in PAST 4.03 based on Bray-Curtis dissimilarities.

340 We furthermore examined associations of environmental variables and niche breadth  
341 with total sterol content and diversity. We calculated phylogenetic independent  
342 contrasts (Felsenstein, 1985; implemented in R package *ape* (Paradis *et al.*, 2004))  
343 with the phylogeny outlined above for sterol contents, Shannon diversity H, positions  
344 on environmental principal component axes (PC1, PC2, PC3), and environmental  
345 niche breadth. Associations between contrasts of sterol content or diversity (as  
346 dependent variable) with contrasts of environmental principal components, niche  
347 breadth, or the 13 separate environmental factors were then individually evaluated by  
348 linear models in R (fitting the regression through the origin).

349 Finally, as 24-methylenecholesterol is a key sterol nutrient for honey bee development  
350 (Svoboda *et al.*, 1980; Herbert *et al.*, 1980), and could therefore have been selected  
351 for as an attracting reward in bee pollinated plants, we tested for differences in 24-  
352 methylenecholesterol content for plants that offer pollen as reward for bees or not (or  
353 for which this interaction was unknown) with a phylogenetic ANOVA (Garland *et al.*,  
354 1993), implemented in the R package *phytools* (Revell, 2012), with 1000 simulations,  
355 and “Holm” post-hoc testing. The same test was also conducted for total sterol content.  
356 Only species with phylogenetic information were included (pollen as bee reward: n =  
357 54; pollen not bee reward: n = 22; unknown: n = 24).

358

## 359 **Results**

### 360 **Pollen sterol composition and diversity across taxa**

361 We profiled 25 phytosterols in pollen of 122 plant species from 51 families including  
362 representatives of Gymnosperms, Nymphaeales, Monocots, Ranunculids,  
363 Caryophyllales, Asterids and Rosids (Fig. 2, Table S1). These phytosterols can be  
364 arranged into biosynthetic pathways with three main distinct branches (i.e., 24C-0,  
365 24C-methyl and 24C-ethyl groups, Fig 3. See Fig. 1 for structure-illustration of the

366 groups). Pollen phytosterols varied qualitatively and quantitatively across taxa with  
367 each species exhibiting a distinctive sterol profile (Fig. 2).

368 Across all the sampled species, the most common pollen sterols (labelled  
369 “commonness”, Fig. 2) were sitosterol (recorded in 97.5% sampled species),  
370 campesterol (88.5%), isofucosterol (82.0%), cholesterol (82.0%), cycloartenol  
371 (81.1%), 24-methylenecholesterol (ostreasterol) (73.0%) and stigmasterol (59.0%).  
372 The most abundant sterol dominating pollen sterol profiles (labelled “abundance”, Fig.  
373 2) was 24-methylenecholesterol (on average accounting for 23% of total sterol  
374 content), followed by isofucosterol (21.5%), sitosterol (20.7%), and cycloartenol  
375 (17.7%). The first three are all  $\Delta^5$  sterols, of which 24-methylenecholesterol belongs to  
376 the 24C-methyl group, whereas sitosterol and isofucosterol belong to the 24C-ethyl  
377 group. Cholesterol, the primary sterol in animals, only represented a small portion  
378 (<1%) of pollen sterol content, despite being common.

379 The pollen sterol diversity of plants varied dramatically with a mean of  $9.98 \pm 4.46$   
380 (mean  $\pm$  s.d.) different phytosterols. For example, the carnivorous plant *Drosera regia*  
381 Stephens had almost exclusively 24-methylenecholesterol in pollen, whereas pollen  
382 from ivy (*Hedera helix* L.) contained 23 different sterols, tea pollen (*Camellia sinensis*  
383 L.) had 22 sterols, and pollen from the bellflowers *Campanula fragilis* Cirillo and  
384 *Campanula isophylla* Moretti had 23 and 19 sterols respectively. However, in all these  
385 species, only one to two sterol compounds were typically major components  
386 (contributing >50% of total sterol content). The variation in the total weights of sterols  
387 led to a Shannon diversity index for pollen sterol composition ranging from 0 in *Drosera*  
388 *regia* to 0.64 in *Hedera helix* (Fig. 2, Table S1), with a mean of 0.34 (note that we  
389 standardized the maximum value of the Shannon diversity index to be 1.0, details see  
390 method).

### 391 **Covariance of pollen sterols**

392 The factor analysis reduced the data to 12 independent latent factors that explained  
393 73% of sterol variation (Table 1). Overall, phytosterols that have close positions in their  
394 biosynthetic pathways (Fig. 3) or use the same enzyme (e.g., reductase) for production  
395 tend to align together with the same factors. For example, iso-obtusifoliol is the  
396 precursor of 24-methylenelophenol, then it branches to either epifungisterol or  
397 avenasterol via episterol (Fig. 3). These four sterol compounds (not including

398 episterol) largely aligned together with factor 1 which accounted for ~9% of the  
399 variance (Table 1). Similar patterns also applied to factor 3 and factor 4 whose main  
400 contributing sterols represented the early cyclopropyl pathway intermediates. Factor  
401 5 represented a strong positive correlation among the stanols (saturated in ring B),  
402 campestanol and sitostanol. Factors 6 and 7 represent products of  $\Delta$ -24 reduction. In  
403 addition, we found one inverse relationship between four of the most common  
404 phytosterols (in factor 2, accounting for 8% of the variance), where 24-  
405 methylenecholesterol was aligned in the opposite direction as the presence of three  
406 other phytosterols: sitosterol, campesterol and stigmasterol.

### 407 **Phylogenetic patterns**

408 We found significant phylogenetic signal in 12 out of 25 phytosterols (percentages of  
409 individual compounds), of which 7 were significant for both Pagel's  $\lambda$  and Blomberg's  
410 K, and 5 for only one of the tests (Fig.2, Table 2). When grouping phytosterols based  
411 on the substitution at C-24 (24C-methyl-, 24C-ethyl-, or 24C-0) or based on the  
412 position of methine in ring B ( $\Delta^0$ ,  $\Delta^5$ ,  $\Delta^7$ ,  $\Delta^8$ ), we found a significant phylogenetic signal  
413 (both Pagel's  $\lambda$  and Blomberg's K) for all groups except the  $\Delta^8$  sterols (Fig.2, Table 2).  
414 Additionally, we found a significant signal for the Shannon diversity index and total  
415 sterol content ( $\mu\text{g}$  sterol per mg pollen; Fig.2, Table 2). These results remain largely  
416 consistent when excluding all taxa which are only identified to genus level. Note that  
417  $\lambda$  and K are largely agreeing on which phytosterols showed significant signal (Table  
418 2), although the significant estimates for  $\lambda$  are relatively high (0.585 to 1, mean = 0.847  
419 for individual compounds; 0.668 to 0.906, mean = 0.79 for categories), whereas those  
420 for K are comparatively low (0.183 to 0.505, mean = 0.332 for individual compounds;  
421 0.158 to 0.201, mean = 0.182 for categories). Some phylogenetic clustering of plants  
422 by overall sterol compositional similarity was also apparent in the NMDS plot, with for  
423 example plants in the Asteraceae, Asparagaceae, or Cactaceae sharing similar sterol  
424 profiles (Fig. S1).

### 425 **Sterols and abiotic environmental factors**

426 How similar pollen sterol profiles are between plants was neither significantly  
427 associated with the similarity between native environmental niches (represented by  
428 environmental principal component axes PC1-PC3) nor with phylogenetic distances

429 ( $r^2 = 0.013$ ,  $p = 0.17$  for additive model in MRM analysis, for individual factors see  
430 Table 3).

431 Total pollen sterol content of plant species was positively correlated with some of the  
432 environmental variables in their native range, but in general the explained variance  
433 ( $r^2$ ) was low (Fig. 4, Table S4). Specifically, total sterol content correlated with  
434 environmental PC1 (associated with high mean temperatures, low temperature  
435 seasonality and low soil carbon content;  $p = 0.015$ ,  $r^2 = 0.060$ ; Fig. 4). For linear  
436 models of individual environmental variables, species with higher total pollen sterol  
437 content tended to occur in locations with higher annual mean temperature, higher  
438 temperatures in the coldest quarter, higher precipitation in the wettest quarter, and  
439 lower temperature seasonality ( $p$ -values  $< 0.05$  for linear models of phylogenetic  
440 independent contrasts,  $r^2$  between 0.05 to 0.08, Table S4), as is the case in tropical  
441 conditions. For Shannon's H diversity of pollen sterol profiles, the only significant  
442 association with environmental variables was a weak negative correlation with  
443 temperature seasonality ( $p = 0.014$ ;  $r^2 = 0.06$ ) (Table S4). None of the other  
444 environmental variables or principal components were significantly correlated with  
445 sterol content or diversity, nor was the total environmental niche breadth (Fig. 4, Table  
446 S4).

#### 447 **Sterols and pollinator guilds**

448 We found overall pollen sterol profiles were largely overlapping between plant groups  
449 with different pollinator guilds (bee, Lepidoptera, generalist insect, bird, unknown; Fig.  
450 5a; ANOSIM: among group  $R = -0.0069$ ,  $p = 0.57$ ; no significant difference for any  
451 pairwise group comparison), and between plants with or without pollen as reward for  
452 bee pollinators (Fig. 5b; ANOSIM:  $R = 0.033$ ,  $p = 0.15$ ). This suggests that pollinator  
453 guilds or the use of pollen as reward by bees do not explain differences in pollen sterol  
454 composition. We note that wind pollinated Angiosperms were not part of this dataset.

455 Neither 24-methylenecholesterol nor total sterol content differed between plants  
456 with/without pollen as reward for bees (phylogenetic ANOVA:  $p = 0.46$  and  $0.66$   
457 respectively; Fig. S2, S3).

458

## 459 Discussion

460 Phytosterols are primary metabolites in plants and are also essential nutrients for  
461 phytophagous insects, making them an important functional trait that provides a  
462 mechanistic link between plants and insects. Our study focused on the pollen sterol  
463 patterns across plant taxa, aiming to provide a more comprehensive overview of pollen  
464 sterol diversity and its relationship with plant phylogeny, abiotic environmental  
465 conditions, and pollinator guilds. We analysed 25 phytosterols in the pollen of more  
466 than 120 angiosperms representing 51 plant families and identified covariance  
467 patterns these phytosterols. Our data are the first to show significant phylogenetic  
468 signal for pollen phytosterols. Although environmental factors and pollinator guilds  
469 showed either weak or no relationships with pollen sterol content, future studies with  
470 more stratified sampling based on more finely defined pollinator species and  
471 environmental conditions could bring more insights about the drivers and importance  
472 of pollen sterol diversity.

### 473 Phylogenetic structure of pollen sterols

474 Similar pollen sterol profiles in related taxa are ultimately due to shared evolutionary  
475 history and proximately due to shared genes for the enzymes involved in their  
476 biosynthesis. Indeed, we show that phylogenetic patterns in pollen sterols in part  
477 reflect their relations in the underlying biosynthesis pathway. For example, we  
478 observed significant phylogenetic clustering of plant species whose pollen sterol  
479 profiles are dominated by 24-methyl (C1-group), 24-ethyl (C2-group) or non-  
480 substituted (C0-group) phytosterols (Fig. 2, Table 2), reflecting the bifurcation of  
481 biosynthesis pathways (Fig. 3). Key enzymes (SMTs, sterol methyltransferases) that  
482 bifurcate the phytosterol pathways are SMT1, which methylates C0 sterol cycloartenol  
483 to 24-methylenecycloartanol, and SMT2, the key and effective enzyme to methylate  
484 24-methyl to 24-ethyl sterols (Akihisa et al., 1991; Nes, 2000; Schaeffer *et al.*, 2001;  
485 Neelakandan *et al.*, 2009). Based on our findings, it would seem likely that the  
486 expression of these enzymes follows phylogenetically conserved patterns in different  
487 clades. Similarly, phylogenetic clusters of the main sterol groups based on the  
488 presence or absence of and position of the double bond in ring B (e.g.,  $\Delta^5$ ,  $\Delta^7$ , or  
489 saturated ring B) suggest conserved expressions of specific desaturases (e.g. STE1  
490 or  $\Delta^7$  and  $\Delta^5$ -sterol-C5-desaturases which convert saturated carbon bonds to

491 methines) and reductase (e.g. DWF5, sterol-  $\Delta^7$  and  $\Delta^5$ -reductase which reduce  
492 methines to saturated bonds) (Benveniste, 2004; Villette *et al.*, 2015).

493 Our factor analysis (Table 1) further revealed an inverse relationship between the  
494 abundance of the major  $\Delta^{24,28}$  sterol (24-methylenecholesterol) and the C24,28-  
495 saturated sterols: campesterol, sitosterol and stigmasterol. This suggests an overall  
496 trade-off of these two groups, the balance of which may be governed by DWF1 (sterol-  
497  $\Delta^{24}$ -reductase) activity. Data from previous studies (Villette *et al.*, 2015; Vanderplanck  
498 *et al.*, 2020a) suggests a similar inverse correlation between 24-methylenecholesterol  
499 and the 24C-ethyl sterols, although this has not been explicitly stated. A high ratio of  
500 24-methylenecholesterol to C24,28-saturated sterols is evident in Cactaceae,  
501 Droseraceae, Rosaceae, Onagraceae and Paeoniaceae. Conversely, C24,28-  
502 saturated sterols are more abundant than 24-methylenecholesterol in Ericaceae,  
503 Primulaceae, Salicaceae and Amaryllidaceae. These families are not closely related,  
504 suggesting convergent evolution of sterol composition. Overall, this indicates an  
505 interplay of environmental selection pressures for particular structural groups and  
506 phylogenetic constraints of sterol biosynthesis enzyme expression.

507 The composition of phytosterols appears to be tissue-dependent (Nes, 1990; Nes *et*  
508 *al.*, 1993). For example, 24-methylenecholesterol has been identified as the main  
509 pollen sterol in many Cactaceae (Fig. 2, Table S1) but is not abundant in their  
510 photosynthetic tissue (Lusby *et al.*, 1993; Standifer *et al.*, 1968; Salt *et al.*, 1987; Li  
511 1996). The unique functional roles in pollen development when compared to the  
512 sporophyte may contribute to the distinct sterol profiles in pollen. We observed strong  
513 correlations among early, cyclopropyl sterol intermediates of the sterol pathway,  
514 particularly 9b,19-cyclopropyl sterols (Table 1). Cycloartenol, 31-norcycloartenol and  
515 24-methylenecycloartanol are correlated with each other: 31-norcycloartenol and 24-  
516 methylenecycloartanol are both derived from cycloartenol. 24,25-  
517 Dehydropollinastanol and 31-norcycloartanol also show high correlation and both are  
518 derived from 31-norcycloartenol. Co-occurrence of cyclopropyl sterols suggests a  
519 reduction in CPI1 (cyclopropyl isomerase) activity and truncation of the sterol pathway,  
520 either within the pollen grain or in the surrounding tapetum cells from which pollen coat  
521 sterols are derived. 9b,19-Cyclopropyl sterols have been identified as key components  
522 of the pollen coat in *Brassica napus* (Villette *et al.*, 2015; Wu *et al.*, 1999). In addition,

523 cycloeucanol is the main sterol synthesised in the growing pollen tube of *Nicotiana*  
524 *tabacum* (Villette *et al.*, 2015; Rotsch *et al.*, 2017).

### 525 **Correlations of phytosterols with abiotic factors**

526 The presence of different phytosterols could be evolutionary adaptations to  
527 environmental conditions. We detected a positive relationship between sterol content  
528 and temperature (particularly mean annual temperature and mean temperature of the  
529 coldest quarter), and a negative correlation with temperature seasonality, even though  
530 the overall association strength was low (Table S4). This indicates that plants found  
531 in cool and temperate climatic conditions were likely to have less pollen sterol than  
532 those found in areas of the world with warm climates with little seasonal fluctuations  
533 (e.g., tropical climates). The association between warmer climates and higher total  
534 amounts of pollen sterols may have evolved as protection against membrane heat  
535 stress, since the role of phytosterols in adaptation to high temperature stress is  
536 established (Dufourc, 2008, Narayanan *et al.* 2016). Phytosterols including  
537 campesterol, sitosterol and avenasterol degrade in stored grain more rapidly at higher  
538 temperatures (Wawrzyniak *et al.*, 2019), so higher sterol content in warmer climates  
539 may avert the risk of their rapid breakdown and limited availability. Besides this, other  
540 pollen sterol characteristics (e.g., sterol diversity and the overall pollen sterol  
541 composition) were not notably associated with abiotic factors. Our sampling, however,  
542 was biased towards plants of temperate regions (the predominant species available  
543 to us for sampling). Limited sampling towards extremes of the environmental gradients  
544 may have reduced our scope to detect associations between abiotic factors and pollen  
545 sterol characteristics. Future work should therefore be targeted at sampling additional  
546 plant species of more extreme environments to fill this gap. Note that our species were  
547 sampled at glasshouses (e.g., tropical glasshouse, alpine glasshouse) or outdoors at  
548 Royal Botanic Gardens Kew and nearby areas (sampling details see Table S1) to get  
549 a first estimate of pollen sterol diversity across a broad range of species. Future in-  
550 depth studies on how abiotic conditions affect pollen sterol variation within-species  
551 deserve further attention to build a more complete overview of pollen sterol diversity  
552 at different taxonomic levels.

### 553 **Impact of sterol diversity on pollinators**

554 Pollen sterol amount and composition did not differ significantly between bee  
555 pollinated and non-bee pollinated plant species. This could indicate that pollen sterols  
556 have generally not been under selection by bee pollinators although we acknowledge  
557 that our analysis combined all bee pollinated plants into one group. Therefore, it  
558 remains possible that pollen sterols play a role in finer scale interactions between  
559 different bee species of varying levels of pollen specialization and their host plants.  
560 We also note that, although we based our assessment of pollinator guilds on the best  
561 available literature data, the quality of evidence for the effective pollinators of the  
562 plants in our data set varied. This calls for further in-depth studies of the relationships  
563 between pollen sterols and pollinators, also including wind-pollinated Angiosperm taxa  
564 missing in this work as points of comparison to animal pollinated plants.

565 A major knowledge gap exists in understanding how important specific phytosterols  
566 are for bees, particularly wild bee species, since many of them are pollen specialists.  
567 Plants adapt nectar chemistry to suit the specific needs of pollinators (Vandeloos *et al.*,  
568 2019) and could similarly alter nutritional chemistry of pollen to optimize its  
569 nutritional suitability for flower visitors. Bee pollinators require a dietary source of  
570 sterols (Wright *et al.*, 2018) and for this they must use the phytosterols found in pollen.  
571 Therefore, determining how lipid components of pollen vary qualitatively and  
572 quantitatively across different plant taxa is important in understanding how nutritionally  
573 limiting landscapes might be for bees, especially where they are not botanically  
574 diverse. For example, honeybee colony growth benefits from 24-methylenecholesterol  
575 (Herbert *et al.*, 1980). Thus, honeybees may be nutritionally limited in landscapes  
576 where floral resources do not provision 24-methylenecholesterol. Our data suggested  
577 that many Asteraceae (e.g., *Achillea ptarmica* L., *Tanacetum vulgare* L. *Achillea*  
578 *millefolium* L., *Jacobaea vulgaris* Gaertn., *Centaurea nigra* L. and *Cirsium vulgare*  
579 (Savi.) Ten) are rich in  $\Delta^7$ -sterols (Fig. 2, Table S1) and lack the common honeybee-  
580 favourable  $\Delta^5$ -sterols (e.g., 24-methylenecholesterol).  $\Delta^7$ -sterols are known to be toxic  
581 to non-specialist herbivores and can only be utilized by some insect species (Behmer  
582 & Nes, 2003; Lang *et al.*, 2012). Thus, plant species that produce unusual phytosterols  
583 in pollen may produce these as defence against pollen herbivory, but some specialist  
584 bee species may have developed mechanisms overcoming this defense. Indeed,  
585 pollen foraging bees on Asteraceae plants are mostly specialized oligolectic bees,  
586 while polylectic bee species avoid the pollen despite the ubiquitous distribution of

587 Asteraceae species and their substantial amount of pollen provision (known as the  
588 Asteraceae paradox, Müller & Kuhlmann, 2008). While the reasons for this Asteraceae  
589 paradox remain unresolved, the abundance of  $\Delta^7$ -sterols we found in the pollen of  
590 Asteraceae species could provide a potential explanation (see also Vanderplanck *et*  
591 *al.*, 2018, 2020b).

592

### 593 **Acknowledgements**

594 We thank Wenxu Zhou for his guidance in analysing sterols, Amy Kendal-Smith,  
595 Richard Moore and James Woodward for their help with pollen collection, Geoffrey  
596 Kite for support with the GC-MS, Ellen Baker for helping revise the manuscript. We  
597 also thank the Horticultural staff of the Royal Botanic Gardens, Kew for maintenance  
598 of the outstanding living collections used to provide pollen for this study. The project  
599 was funded by a BBSRC grant to GAW (BB/P007449/1) and PCS (BB/P005276/1)  
600 and a Peter Sowerby Grant to PCS and HK. PZ was supported to work with DN at  
601 Texas Tech on a BBSRC International Travel Award Scheme grant (BB/S004653/1)  
602 to PCS.

603

### 604 **Author contributions**

605 PZ, HK, and PCS designed the research. PZ collected the pollen and extracted and  
606 analysed sterols. DN, IWF and DIF helped with sterol identification and  
607 quantification. OS collected phylogenetic information on studied species and  
608 conducted phylogenetic analyses. SP, CP, and IO collected species geographic and  
609 environmental information and performed abiotic niche analyses. HK and PZ  
610 collected pollinator records on studied species, and HK conducted analyses with  
611 environmental factors and pollinator guilds. EM and GW conducted factor analysis.  
612 WDN generated sterol biosynthesis pathways. PZ and PCS drafted the manuscript.  
613 HK, OS, SP, GAW and all other authors contributed in writing and revising the  
614 manuscript.

615

### 616 **References**

- 617 **Akihisa T, Kokke WCMC, Tamura T. 1991.** Naturally occurring sterols and related  
618 compounds from plants. In *Physiology and Biochemistry of Sterols* (Ed. G. W.  
619 Patterson and W. D. Nes). AOCS press, Champaign, pp 172–228.
- 620
- 621 **Amar, S., Becker, H. C., & Möllers, C. 2008.** Genetic variation and genotype×  
622 environment interactions of phytosterol content in three doubled haploid populations  
623 of winter rapeseed. *Crop science* **48**(3): 1000-1006.
- 624
- 625 **Arnold SEJ, Peralta Idrovo ME, Lomas Arias LJ, Belmain SR, Stevenson PC 2014.**  
626 Herbivore Defence Compounds Occur in Pollen and Reduce Bumblebee Colony Fitness.  
627 *Journal of Chemical Ecology* **40**: 878–881.
- 628
- 629 **Behmer ST, Elias DO. 1999.** The nutritional significance of sterol metabolic  
630 constraints in the generalist grasshopper *Schistocerca americana*. *Journal of Insect*  
631 *Physiology* **45**: 339–348.
- 632
- 633 **Behmer ST, Elias DO. 2000.** Sterol metabolic constraints as a factor contributing to  
634 the maintenance of diet mixing in grasshoppers (Orthoptera: Acrididae).  
635 *Physiological and Biochemical Zoology* **73**: 219–230.
- 636
- 637 **Behmer ST, Nes WD. 2003.** Insect Sterol Nutrition and Physiology: A global  
638 overview. *Advances in Insect Physiology* **31**: 1–72.
- 639
- 640 **Benveniste P. 2004.** Biosynthesis and accumulation of sterols. *Ann. Rev. Plant Biol*  
641 **55**: 429–457.
- 642
- 643 **Blomberg SP, Garland Jr. T, Ives AR, Crespi B. 2003.** Testing for phylogenetic  
644 signal in comparative data: behavioral traits are more labile. *Evolution* **57**: 717–745.
- 645

- 646 **Brummitt RK. 2001.** World Geographical Scheme for Recording Plant Distributions,  
647 Edition 2. Biodiversity Information Standards (TDWG)  
648 <http://www.tdwg.org/standards/109>  
649
- 650 **Capinha C, Pateiro-López B. 2014.** Predicting species distributions in new areas or  
651 time periods with alpha-shapes. *Ecological Informatics* **24**: 231–237.  
652
- 653 **Clarke KR. 1993.** Non-parametric multivariate analyses of changes in community  
654 structure. *Australian Journal of Ecology* **18**: 117–143.  
655
- 656 **Chung HY, Fujioka S, Choe S, Lee S, Lee YH, Baek NI, Chung IS. 2010.**  
657 Simultaneous suppression of three genes related to brassinosteroid (BR)  
658 biosynthesis altered campesterol and BR contents, and led to a dwarf phenotype in  
659 *Arabidopsis thaliana*. *Plant Cell Reports* **29**: 397–402  
660
- 661 **Danielson JJ, Gesch DB. 2011.** Global multi-resolution terrain elevation data 2010  
662 (gmted2010). Technical report, US Geological Survey  
663
- 664 **Dauby G, Zaiss R, Blach-Overgaard A, Catarino L, Damen T, Deblauwe V,**  
665 **Dessein S, Dransfield J, Droissart V, Duarte MC. et al. 2016.** Rainbio: a mega-  
666 database of tropical African vascular plants distributions. *PhytoKeys* **74**: 1-18.  
667
- 668 **Dornelas M, Antao LH, Moyes F, Bates AE, Magurran AE, Adam D,**  
669 **Akhmetzhanova, AA, Appeltans W, Arcos JM, Arnold H, et al. 2018.** Biotime: A  
670 database of biodiversity time series for the anthropocene. *Global Ecology and*  
671 *Biogeography* **27**: 760–786  
672
- 673 **Dufourc EJ. 2008.** The role of phytosterols in plant adaptation to temperature.  
674 *Plant Signalling & Behavior* **3**: 133–134.

675

676 **Felsenstein J. 1985.** Phylogenies and the comparative method. *The American*  
677 *Naturalist* **125**: 1–15.

678

679 **Ferrer A, Altabella T, Arr' o M, Boronat A. 2017.** Emerging roles for conjugated  
680 sterols in plants. *Progress In Lipid Research* **67**: 27–37.

681

682 **Garland T, Dickerman AW, Janis CM, Jones JA. 1993.** Phylogenetic analysis of  
683 covariance by computer simulation. *Systematic Biology* **42**: 265-292.

684

685 **Goslee SC, Urban DL. 2007.** The ecodist package for dissimilarity-based analysis of  
686 ecological data. *Journal of Statistical Software* **22**: 1–19.

687

688 **Grove MD, Spencer, GF, Rohwedder WK, Mandava N, Worley JF, Warthen JD,**  
689 **Steffens GL, Flippen-Anderson JL, Cook JC. 1979.** Brassinolide, a plant growth-  
690 promoting steroid isolated from *Brassica napus* pollen. *Nature* **281**: 216–217.

691

692 **Grunwald C. 1971.** Effects of free sterols, steryl ester, and steryl glycoside on  
693 membrane permeability. *Plant Physiology* **48**: 653–655.

694

695 **Guo D, Venkatramesh, M, Nes WD. 1995.** Developmental Regulation of Sterol  
696 Biosynthesis in *Zea mays*. *Lipids* **30**: 203-219.

697

698 **Hammer Ø, Harper DAT, Ryan PD 2001.** PAST: Paleontological statistics software  
699 package for education and data analysis. *Palaeontologia Electronica* **4**: 9pp.

700

701 **Hartmann M-A. 1998.** Plant sterols and the membrane environment. *Trends in Plant*  
702 *Science* **3**: 170–175.

703

- 704 **Heupel, R. C. and Nes, W. D.** 1994. Evidences for Differences in Sterol  
705 Biosynthesis and Derivatization in Sorghum. *J. Nat. Prod.* 47, 292-299  
706
- 707 **Herbert EW, Svoboda JA, Thompson MJ, Shimanuki H.** 1980. Sterol utilization in  
708 honey bees fed a synthetic diet: Effects on brood rearing. *Journal of Insect*  
709 *Physiology* 26: 287–289.  
710
- 711 **Hinchliff CE, Smith SA, Allman JF, Burleigh JG, Chaudhary R, Coghill LM,**  
712 **Crandall KA, Deng J, Drew BT, Gazis R, et al.** 2015. Synthesis of phylogeny and  
713 taxonomy into a comprehensive tree of life. *Proceedings of the National Academy of*  
714 *Sciences* 112: 12764–12769.  
715
- 716 **ISRIC.** 2013. Soilgrids: an automated system for global soil mapping  
717
- 718 **Jing X, Behmer ST.** 2020. Insect sterol nutrition: physiological mechanisms,  
719 ecology, and applications. *Annual Review of Entomology* 65: 251–271.  
720
- 721 **Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW,**  
722 **Zimmermann NE, Linder HP, Kessler M** 2017. Climatologies at high resolution for  
723 the earths land surface areas. *Scientific Data* 4: 170122.  
724
- 725 **Keck F, Rimet F, Bouchez A, Franc A.** 2016. phylosignal: an r package to  
726 measure, test, and explore the phylogenetic signal. *Ecology and Evolution* 6: 2774–  
727 2780.  
728
- 729 **Kembel S, Cowan P, Helmus M, Cornwell W, Morlon H, Ackerly D, Blomberg S,**  
730 **Webb C** 2010. Picante: R tools for integrating phylogenies and ecology.  
731 *Bioinformatics*, 26, 1463–1464.  
732

- 733 **Khripach V, Zhabinsk V, de Groot A. 2000.** Twenty Years of Brassinosteroids:  
734 Steroidal Plant Hormones Warrant Better Crops for the XXI Century. *Annals of*  
735 *Botany* **86**: 441–447.
- 736
- 737 **Knuth P. 1908.** Handbook of Flower Pollination. Volume 2. Oxford, At the Clarendon  
738 Press, UK.
- 739
- 740 **Knuth P. 1909.** Handbook of Flower Pollination. Volume 3. Oxford, At the Clarendon  
741 Press, UK.
- 742
- 743 **Kumar MS, Ali K, Dahuja A, Tyagi A. 2015.** Role of phytosterols in drought stress  
744 tolerance in rice. *Plant Physiology and Biochemistry* **96**: 83-89.
- 745
- 746 **Lang M, Murat S, Clark AG, Gouppil G, Blais C, Matzkin LM, Guittard E,**  
747 **Yoshiyama-Yanagawa T, Kataoka H, Niwa R, et al. 2012.** Mutations in the  
748 neverland gene turned *Drosophila pachea* into an obligate specialist species.  
749 *Science* **337**: 1658–1661.
- 750
- 751 **Li, Y. 1996.** Stereochemical Studies on the Metabolism of Sterols by  
752 *Saccharomyces cerevisiae* strain GL7. PhD thesis, Texas Tech University, TX, USA.
- 753
- 754 **Lichstein JW. 2007.** Multiple regression on distance matrices: a multivariate spatial  
755 analysis tool. *Plant Ecology* **188**: 117-131.
- 756
- 757 **Lusby WR, Buchmann SL, Feldlaufer MF. 1993.** Pollen sterols from three species  
758 of Sonoran cacti. *Lipids* **28**: 469-470.
- 759
- 760 **Meyer C, Weigelt P, Kreft H. 2016.** Multidimensional biases, gaps and  
761 uncertainties in global plant occurrence information. *Ecology Letters* **19**: 992–1006.

762

763 **Michonneau F, Brown JW, Winter DJ. 2016.** rotl: an r package to interact with  
764 the open tree of life data. *Methods in Ecology and Evolution* **7**: 1476–1481.

765

766 **Moreau RA, Nyström L, Whitaker BD, Winkler-Moser JK, Baer DJ, Gebauer SK,**  
767 **Hicks KB. 2018.** Phytosterols and their derivatives: Structural diversity, distribution,  
768 metabolism, analysis, and health-promoting uses. *Progress in Lipid Research* **80**:  
769 35–61.

770

771 **Müller A, Kuhlmann M. 2008.** Pollen hosts of western palaeartic bees of the  
772 genus *Colletes* (Hymenoptera: Colletidae): the Asteraceae paradox. *Biological*  
773 *Journal of the Linnean Society* **95**: 719–733.

774

775 **Narayanan S, Tamura PJ, Roth MR, Prasad PV, Welti R. 2016.** Wheat leaf lipids  
776 during heat stress: I. High day and night temperatures result in major lipid  
777 alterations. *Plant, Cell & Environment* **39**: 787–803.

778

779 **Neelakandan AK, Song Z, Wang J, Richards MH, Wu X, Valliyodan B, Nguyen**  
780 **HT, Nes WD. 2009.** Cloning, Functional Expression, and Phylogenetic Analysis of  
781 Plant 24C-Methyltransferases Involved in Sitosterol  
782 Biosynthesis. *Phytochemistry* **70**: 1982–1998.

783

784 **Nes WD. 2011.** Biosynthesis of cholesterol and other sterols. *Chemical Reviews*  
785 **111**: 6423–6451.

786

787 **Nes WD, Zhou W, Ganapathy K, Liu J, Vatsyayan R, Chamala S, Hernandez K,**  
788 **Miranda, M. 2009.** Sterol 24-C-methyltransferase: An enzymatic target for the  
789 disruption of ergosterol biosynthesis and homeostasis in *Cryptococcus*  
790 *neoformans*. *Archives of Biochemistry and Biophysics*. **481**: 210–218

791

792 **Nes WD, Song Z, Dennis AL, Zhou W, Nam J, Miller M. 2003.** Biosynthesis of  
793 Phytosterols: Kinetic mechanism for the Enzymatic C-Methylation of Sterols. *Journal*  
794 *of Biological Chemistry*. **278**: 34505–34516.

795

796 **Nes WD. 2000.** Sterol Methyl Transferase: Enzymology and Inhibition. *Biochimica*  
797 *Biophysica Acta*. **1529**: 63–88.

798

799 **Nes WD, Parker SR, Crumley FG, Ross SA. 1993.** Regulation of phytosterol biosynthesis. In *Lipid*  
800 *Metabolism in Plants*. Ed. T. S. Moore. CRC Press, 1993. 389–426

801

802 **Nes WD. 1990.** Control of sterol biosynthesis and its importance to developmental  
803 regulation and evolution. *Recent Advances in Phytochemistry*. **24**: 283–327.

804

805 **Nes WR, Nes WD, 1980.** *Lipids in evolution*. Plenum Press, New York.

806

807 **Pagel M. 1999.** Inferring the historical patterns of biological evolution. *Nature* **401**:  
808 877–884.

809

810 **Palmer-Young EC, Farrell IW, Adler, IW, Milano NJ, Egan PA, Junker RR,**  
811 **Irwin RE, Stevenson PC. 2019.** Chemistry of floral rewards: intra- and interspecific  
812 variability of nectar and pollen secondary metabolites across taxa. *Ecological*  
813 *Monographs* **89**: e01335.

814

815 **Paradis E, Claude J, Strimmer K. 2004.** Ape: analyses of phylogenetics and  
816 evolution in R language. *Bioinformatics* **20**: 289–290.

817

818 **Piironen V, Lindsay DG, Miettinen TA, Toivo J, Lampi AM. 2000.** Plant sterols:  
819 biosynthesis, biological function and their importance to human nutrition. *Journal of*  
820 *the Science of Food and Agriculture* **80**: 939–966.

821

822 **Poisot T, Kéfi S, Morand S, Stanko M, Marquet PA, Hochberg ME 2015.** A  
823 Continuum of Specialists and Generalists in Empirical Communities. *PLoS ONE* **10**:  
824 e0114674.

825

826 **Posé D, Castanedo I, Borsani O, Nieto B, Rosado A, Taconnat L, Ferrer A,**  
827 **Dolan L, Valpuesta V, Botella MA. 2009.** Identification of the Arabidopsis  
828 dry2/sqe1-5 mutant reveals a central role for sterols in drought tolerance and  
829 regulation of reactive oxygen species. *Plant Journal* **59**: 63–76.

830

831 **Qi X, Bakht S, Qin B, Leggett M, Hemmings A, Mellon F, Eagles J, Werck-**  
832 **Reichhart, D, Schaller H, Lesot A, et al. 2006.** A different function for a member  
833 of an ancient and highly conserved cytochrome p450 family: from essential sterols  
834 to plant defense. *Proceedings of the National Academy of Sciences* **103**: 18848–  
835 18853,

836

837 **R Core Team 2020.** R: A language and environment for statistical computing. R  
838 Foundation for Statistical Computing, Vienna, Austria. URL [https://www.R-](https://www.R-project.org/)  
839 [project.org/](https://www.R-project.org/)

840

841 **Rasmussen C, Engel MS, Vereecken NJ. 2020.** A primer of host-plant  
842 specialization in bees. *Emerging Topics in Life Sciences* **4**: 7-17.

843

844 **Rees JA, Cranston K. 2017.** Automated assembly of a reference taxonomy for  
845 phylogenetic data synthesis. *Biodiversity Data Journal* **5**: e12581.

846

847 **Revell LJ. 2012.** Phytools: an R package for phylogenetic comparative biology (and  
848 other things). *Methods in Ecology and Evolution* **3**: 217-223.

849

850 **Rivest, S., Forrest, J. R. K. 2020.** Defence compounds in pollen: why do they occur  
851 and how do they affect the ecology and evolution of bees? *New Phytologist*, **225**(3),  
852 1053-1064.

853

854 **Rotsch AH, Kopka J, Feussner I, Ischebeck T. 2017.** Central metabolite and  
855 sterol profiling divides tobacco male gametophyte development and pollen tube  
856 growth into eight metabolic phases. *The Plant Journal* **92**: 129–146.

857

858 **Salt TA, Tocker JE, Adler JH. 1987.** Dominance of  $\Delta^5$ -sterols in eight species of the  
859 Cactaceae. *Phytochemistry* **26**: 731-733.

860

861 **Sanchez-Reyes LL, O'Meara BC. 2019.** Datelife: leveraging databases and  
862 analytical tools to reveal the dated Tree of Life. <https://t.co/ovO81Fh0pn> #bioRxiv

863

864 **Schaeffer A, Bronner R, Benveniste P, Schaller H. 2001.** The ratio of campesterol  
865 to sitosterol that modulates growth in Arabidopsis is controlled by sterol  
866 methyltransferase 2;1. *Plant Journal* **25**: 605–615

867

868 **Schuler I, Milon A, Nakatani Y, Ourisson G, Albrecht A-M, Benveniste P,**  
869 **Hartman M-A. 1991.** Differential effects of plant sterols on water permeability and on  
870 acyl chain ordering of soybean phosphatidylcholine bilayers. *Proceedings of the*  
871 *National Academy of Sciences* **88**: 6926–6930.

872

873 Schwery O, **O'Meara BC** 2016. MonoPhy: a simple R package to find and visualize  
874 monophyly issues. *PeerJ Computer Science* 2:e56.

875

876 **Smith SA, Beaulieu JM, Donoghue MJ. 2010.** An uncorrelated relaxed-clock  
877 analysis suggests an earlier origin for flowering plants. *Proceedings of the National*  
878 *Academy of Sciences* **107**: 5897–5902.

879

880 **Standifer LN, Devys M, Barbier M. 1968.** Pollen sterols—a mass spectrographic  
881 survey. *Phytochemistry* **7**: 1361-1365.

882

883 **Svoboda JA, Thompson MJ, Herbert EW, Shimanuki H. 1980.** Sterol utilization in  
884 honey bees fed a synthetic diet: Analysis of prepupal sterols. *Journal of Insect*  
885 *Physiology* **26**: 287-289.

886

887 **Vandelook F, Janssens SB, Gijbels P, Fischer E, Van den Ende W, Honnay O,**  
888 **Abrahamczyk S. 2019.** Nectar traits differ between pollination syndromes in  
889 Balsaminaceae. *Annals of Botany* **124**: 269–279.

890

891 **Vanderplanck M, Decleves S, Roger N, Decroo C, Caulier G, Glauser G, Michez**  
892 **D. 2018.** Is non-host pollen suitable for generalist bumblebees? *Insect Science* **25**:  
893 259–272.

894

895 **Vanderplanck M, Zerck P-L, Lognay G, Michez D. 2020a.** Generalized host-plant  
896 feeding can hide sterol-specialized foraging behaviors in bee–plant interactions.  
897 *Ecology and Evolution* **10**: 150–162.

898

899 **Velasco, L., Fernández-Cuesta, Á., García-Ruiz, J. R., Fernández-Martínez, J.**  
900 **M., & Domínguez-Giménez, J. 2013.** Genetic variation and genotype× environment  
901 interactions for seed phytosterols in sunflower. *Crop Science*, **53**(4): 1589-1593.

902

903 **Vanderplanck M, Gilles H, Nonclercq D, Duez P, Gerbaux P. 2020b.** Asteraceae  
904 Paradox: Chemical and Mechanical Protection of *Taraxacum* Pollen. *Insects* **11**: 304.

905

- 906 **Villette C, Berna A, Compagnon V, Schaller H. 2015.** Plant sterol diversity in  
907 pollen from angiosperm. *Lipids* **50**: 749–760.
- 908
- 909 **Wang K, Senthil-Kumar M, Ryu C-M, Kang L, Mysore KS. 2012.** Phytosterols Play  
910 a Key Role in Plant Innate Immunity against Bacterial Pathogens by Regulating  
911 Nutrient Efflux into the Apoplast. *Plant Physiology* **158**: 1789–1802.
- 912
- 913 **Wawrzyniak J, Gawrysiak-Witulska M, Rudzińska M. 2019.** Dynamics of  
914 phytosterol degradation in a bulk of rapeseed stored under different temperature and  
915 humidity conditions. *Journal of Stored Products Research* **83**: 292–304.
- 916
- 917 **WCVP 2020.** World Checklist of Vascular Plants, version 2.0. Facilitated by the  
918 Royal Botanic Gardens, Kew. Published on the Internet; <http://wcvp.science.kew.org/>
- 919
- 920 **Westrich P. 2018.** Die Wildbienen Deutschlands. Germany: Verlag Eugen Ulmer.
- 921
- 922 **Wright GA, Nicolson SW, Shafir S. 2018.** Nutritional physiology and ecology of  
923 honey bees. *Annual Review of Entomology* **63**: 327–344.
- 924
- 925 **Wu SS, Moreau RA, Whitaker BD, Huang AH. 1999.** Steryl esters in the elaioplasts  
926 of the tapetum in developing Brassica anthers and their recovery on the pollen  
927 surface. *Lipids* **34**: 517-523.
- 928
- 929
- 930 **Xu S, Norton RA, Crumley FG, Nes WD. 1988.** Comparison of the  
931 Chromatographic Properties of Sterols, Select Additional Steroids and  
932 Triterpenoids: Gravity-flow Liquid Chromatography, Thin-layer Chromatography,  
933 Gas-liquid Chromatography, and High Performance Liquid Chromatography. *Journal*  
934 *of Chromatography.* **4** 377–398.
- 935

936 **Yeagle PL. 1985.** Cholesterol and the cell membrane. *Biochimica et Biophysica Acta*  
937 **822:** 267–287.

938

939 **Zanne AE, Tank DC, Cornwell WK, Eastman JM, Smith SA, FitzJohn RG,**  
940 **McGlenn DJ, O'Meara BC, Moles AT, Reich PB, et al. 2014.** Three keys to the  
941 radiation of angiosperms into freezing environments. *Nature* **506:** 89–92.

942

943 **Zizka A, Silvestro D, Andermann T, Azevedo J, Duarte Ritter C, Edler D, Farooq**  
944 **H, Herdean A, Ariza M, Scharn R, et al. 2019.** Coordinate cleaner: Standardized  
945 cleaning of occurrence records from biological collection databases. *Methods in*  
946 *Ecology and Evolution* **10:** 744–751.

947

948 **Figure legends:**

949 **Fig. 1.** Chemical structure of 24-methylenecholesterol as an illustration of  
 950 phytosterols showing a) carbon numbering; b) different substitutions in ring B; and c)  
 951 different substitutions at C-24.

952 **Fig. 2.** Pollen sterol profiles of plant species. Phylogenetic relationships are given on  
 953 the left, bold numbers indicate families. Relative contributions of individual sterols to  
 954 each species' total sterol content are given in the centre; commonness (proportion of  
 955 plants containing an individual sterol) and relative abundance (average proportion of  
 956 individual sterol in each species) are given at the bottom; deeper reds indicate  
 957 values closer to 1. Shannon diversity index (H), 24th carbon groups, delta groups,  
 958 and total sterol content are given on the right; circle size represents sums of relative  
 959 sterol contents in the respective groups (0 to 1), and log of  $\mu\text{g}$  per mg pollen for total  
 960 sterol content. Sterol names and groups are coloured in the same fashion as  
 961 illustrated in Fig. 1. Families: 1 - Pinaceae, 2 - Nymphaeaceae, 3 - Colchicaceae, 4 -  
 962 Cannaceae, 5 - Strelitziaceae, 6 - Iridaceae, 7 - Asphodelaceae, 8 - Asparagaceae,  
 963 9 - Amaryllidaceae, 10 - Ranunculaceae, 11 - Papaveraceae, 12 - Paeoniaceae, 13 -  
 964 Geraniaceae, 14 - Myrtaceae, 15 - Onagraceae, 16 - Cistaceae, 17 - Malvaceae, 18  
 965 - Oxalidaceae, 19 - Salicaceae, 20 - Linaceae, 21 - Euphorbiaceae, 22 - Fagaceae,  
 966 23 - Cucurbitaceae, 24 - Rosaceae, 25 - Fabaceae, 26 - Droseraceae, 27 -  
 967 Caryophyllaceae, 28 - Nyctaginaceae, 29 - Cactaceae, 30 - Hydrangeaceae, 31 -  
 968 Polemoniaceae, 32 - Theaceae, 33 - Ericaceae, 34 - Primulaceae, 35 - Araliaceae,  
 969 36 - Apiaceae, 37 - Adoxaceae, 38 - Caprifoliaceae, 39 - Campanulaceae, 40 -  
 970 Menyanthaceae, 41 - Asteraceae, 42 - Apocynaceae, 43 - Convolvulaceae, 44 -  
 971 Solanaceae, 45 - Boraginaceae, 46 - Gesneriaceae, 47 - Scrophulariaceae, 48 -  
 972 Plantaginaceae, 49 - Bignoniaceae, 50 - Phrymaceae, 51 - Lamiaceae.

973 **Fig. 3.** Hypothetical biosynthetic pathways of phytosterols identified in this study  
 974 (pathways based on Benveniste, 2004).

975 **Fig. 4.** Correlation plots of phylogenetically independent contrasts (PICs) of positions  
 976 on the environmental principal component axes (PC1-PC3) and environmental niche  
 977 breadth against PICs of total pollen sterol amounts (top row) and sterol profile  
 978 Shannon diversities H (bottom row). Blue dashed lines indicate regression lines of

979 linear models (with intercept set to zero);  $r^2$  and p-values for linear models inserted in  
980 the respective plot. PC loadings from each environmental variable see Table S3.

981 **Fig. 5.** 2D-NMDS plots of sterol profiles for plants (a) with different pollinator guilds,  
982 and (b) with/without pollen as bee reward. Distances correspond to sterol profile  
983 dissimilarity (Bray-Curtis distances). Stress of NMDS solution: 0.202.

984

985 **Tables:**

986 **Table 1.** Factor analysis identifying the covariance of 25 sterols measured across all  
987 the plant species surveyed. The main contributor(s) for each component is  
988 highlighted in grey.

For Peer Review

Component	1	2	3	4	5	6	7	8	9	10	11	12
Eigenvalue	2.146	2.053	1.874	1.761	1.581	1.460	1.421	1.356	1.234	1.105	1.098	1.069
Proportion of variance explained	0.086	0.082	0.075	0.070	0.063	0.058	0.057	0.054	0.049	0.044	0.044	0.043
Cumulative proportion of variance explained	0.086	0.168	0.243	0.313	0.376	0.434	0.491	0.545	0.594	0.638	0.682	0.725
<b>Sterol</b>												
	<b>Factor loading</b>											
Cycloartenol	-0.013	0.113	0.054	-0.683	-0.061	0.042	0.073	0.163	0.338	-0.131	0.107	-0.217
31-Norcycloartenol	-0.234	0.131	-0.022	-0.677	0.028	-0.009	-0.199	-0.251	-0.168	-0.094	-0.149	0.156
24,25-Dehydropollinastanol	0.019	0.065	-0.919	0.027	-0.022	-0.028	-0.060	0.049	-0.072	0.035	0.018	0.066
Pollinastanol	0.112	0.278	0.139	0.102	-0.046	-0.309	-0.415	0.278	-0.168	-0.286	-0.143	-0.027
Lathosterol	-0.021	0.039	-0.157	0.026	-0.004	-0.018	0.114	0.032	0.086	-0.040	-0.205	-0.740
Cholesterol	0.068	0.048	0.072	0.113	-0.194	0.670	0.090	-0.150	-0.081	0.091	-0.222	-0.080
31-Norcycloartanol	0.033	0.070	-0.924	0.032	-0.021	-0.033	0.024	0.067	0.011	0.013	-0.029	-0.222
14-Methylcholest-8-enol	0.064	-0.018	-0.038	0.048	-0.036	0.014	-0.736	-0.095	0.233	0.018	-0.001	-0.009
Desmosterol	0.051	0.077	-0.082	0.125	-0.069	-0.086	0.272	0.170	0.320	-0.110	-0.418	0.571
24-Methylenecholesterol	0.208	0.531	0.256	0.266	-0.157	-0.378	-0.036	0.282	-0.350	0.215	-0.027	-0.073
24-Methylenecycloartanol	0.035	-0.036	0.035	-0.794	-0.035	-0.066	0.009	0.130	-0.054	0.128	-0.022	0.016
Cycloaucalenol	0.001	0.076	0.037	-0.001	-0.035	-0.017	-0.013	0.053	-0.057	-0.925	0.004	-0.005
Obtusifolol	-0.022	0.115	0.067	-0.003	0.003	-0.082	-0.143	0.052	0.801	0.084	-0.014	-0.005
Iso-obtusifolol	-0.741	-0.138	-0.005	-0.049	0.017	-0.026	-0.243	0.255	-0.070	0.071	-0.006	0.001
24-Methylenelophenol	-0.823	0.085	0.018	-0.079	0.056	-0.049	0.114	-0.036	-0.086	0.027	-0.062	0.053
Episterol	0.014	0.072	-0.002	0.058	-0.043	-0.098	0.088	-0.033	0.013	-0.009	0.842	0.117
Epifungisterol	-0.414	-0.005	-0.042	-0.211	0.041	0.056	-0.641	0.016	-0.091	-0.018	-0.072	0.101
Campesterol	0.052	-0.756	0.101	0.113	-0.101	-0.131	0.079	-0.024	-0.121	0.107	0.096	0.009
Campestanol	0.013	-0.042	0.019	0.028	0.823	-0.172	0.048	-0.076	0.046	0.006	-0.064	-0.056
Avenasterol	-0.752	0.078	0.044	0.045	-0.066	0.126	-0.022	0.026	0.250	-0.090	0.090	-0.128
Schottenol	-0.101	0.133	0.032	-0.031	0.250	0.778	-0.143	0.297	-0.045	-0.069	0.077	0.064
Sitosterol	0.104	-0.746	0.085	0.028	0.143	0.027	-0.005	-0.263	-0.036	-0.024	-0.152	-0.044
Sitostanol	-0.030	0.059	0.021	0.031	0.838	0.229	-0.029	0.155	-0.053	0.038	0.032	0.044
Isofucosterol	0.162	-0.091	0.133	0.121	-0.092	-0.072	-0.090	-0.771	-0.086	0.084	0.048	-0.010
Stigmasterol	-0.076	-0.653	0.018	0.031	-0.099	-0.095	-0.110	0.411	-0.059	0.085	-0.068	0.052

990 **Table 2.** Identity of phytosterols in pollen of 122 plant species showing  
 991 those with phylogenetic signal across species. All values presented are based  
 992 on the percentage values of sterols except total sterols content ( $\mu\text{g}/\text{mg}$  sampled  
 993 pollen).  $\Delta$  and C-24 value indicates structure of ring B and on the 24<sup>th</sup> carbon (see  
 994 Fig. 1 for details). Pagel's  $\lambda$  and Blomberg's  $K$  are used for testing phylogenetic  
 995 signal. P-values for each test are given accordingly ( $p\text{-}\lambda$  and  $p\text{-}K$ ). Sterols with  
 996 significant phylogenetic signals are in bold.

Trivial Name	Semi-systematic Name	$\Delta$	C-24	$\lambda$	$p\text{-}\lambda$	$K$	$p\text{-}K$
Cycloartenol	4,4,14-trimethyl 9 $\beta$ ,19-cyclo-cholest-24-en-3b-ol	0	C0	<0.001	1.000	0.103	0.407
<b>31-Norcycloartenol</b>	4,14-dimethyl 9 $\beta$ ,19-cyclo-cholest-24-en-3b-ol	0	C0	<b>1.003</b>	<b>0.001</b>	<b>0.505</b>	<b>0.021</b>
<b>24,25-Dehydropollinastanol</b>	14-methyl 9 $\beta$ ,19-cyclo-cholest-24-en-3b-ol	0	C0	<b>1.003</b>	<b>0.001</b>	<b>0.354</b>	<b>0.036</b>
<b>Pollinastanol</b>	14-methyl 9 $\beta$ ,19-cyclo-cholestan-3b-ol	0	C0	<b>0.681</b>	<b>0.001</b>	0.169	0.107
Lathosterol	cholest-7-en-3b-ol	7	C0	<0.001	1.000	0.077	0.754
Cholesterol	cholest-5-en-3b-ol	5	C0	0.076	1.000	0.039	0.938
<b>31-Norcycloartanol</b>	4,14-dimethyl 9 $\beta$ ,19-cyclo-cholestan-3b-ol	0	C0	<b>1.003</b>	<b>0.001</b>	<b>0.208</b>	<b>0.050</b>
14-Methylcholest-8-enol	14-methyl cholest-8-en-3b-ol	8	C0	<0.001	1.000	0.093	0.615
Desmosterol	cholesta-5,24-dien-3b-ol	5	C0	0.263	1.000	0.196	0.072
<b>24-methylenecholesterol</b>	24-methyl cholesta-5,24(28)-dien-3b-ol	5	C1	<b>0.747</b>	<b>0.001</b>	<b>0.183</b>	<b>0.012</b>
24-Methylenecycloartanol	4,4,14,24-tetramethyl 9 $\beta$ ,19-cyclo-cholest-24(28)-en-3b-ol	0	C1	<0.001	1.000	0.162	0.227
Cycloeucalenol	4,14,24-trimethyl 9 $\beta$ ,19-cyclo-cholest-24(28)-en-3b-ol	0	C1	<0.001	1.000	0.219	0.147
Obtusifoliol	4,14,24-trimethyl cholesta-8,24(28)-dien-3b-ol	8	C1	<0.001	1.000	0.167	0.132
<b>Iso-obtusifoliol</b>	4,14,24-trimethyl cholesta-7,24(28)-dien-3b-ol	7	C1	<b>0.973</b>	<b>0.001</b>	<b>0.250</b>	<b>0.039</b>
24-Methylenelophenol	4,24-dimethyl cholesta-7, 24(28)-dien-3b-ol	7	C1	<0.001	1.000	0.102	0.495
<b>Episterol</b>	24-methyl cholesta-7,24(28)-dien-3b-ol	7	C1	<b>0.990</b>	<b>0.001</b>	<b>0.505</b>	<b>0.021</b>
<b>Epifungisterol</b>	24 $\alpha$ -methyl cholest-7-en-3b-ol	7	C1	<b>0.694</b>	<b>0.018</b>	0.236	0.055
<b>Campesterol</b>	24 $\alpha$ -methyl cholest-5-en-3b-ol	5	C1	<b>0.585</b>	<b>0.005</b>	<b>0.240</b>	<b>0.011</b>
Campestanol	24 $\alpha$ -methyl cholestan-3b-ol	0	C1	<0.001	1.000	0.290	0.119
Avenasterol	24-ethyl cholesta-7,24(28) trans-dien-3b-ol	7	C2	<0.001	1.000	0.069	0.814
<b>Schottenol</b>	24 $\alpha$ -ethyl cholest-7-en-3b-ol	7	C2	<b>1.003</b>	<b>0.001</b>	<b>0.440</b>	<b>0.008</b>
<b>Sitosterol</b>	24 $\alpha$ -ethyl cholest-5-en-3b-ol	5	C2	0.275	1.000	<b>0.183</b>	<b>0.012</b>
Sitostanol	24 $\alpha$ -ethyl cholestan-3b-ol	0	C2	<0.001	1.000	0.093	0.606
<b>Isofucosterol</b>	24-ethyl cholesta-5,24(28) trans-dien-3b-ol	5	C2	<b>0.634</b>	<b>0.002</b>	0.141	0.053
Stigmasterol	24 $\alpha$ -ethyl cholesta-5,22 trans-dien-3b-ol	5	C2	<0.001	1.000	0.149	0.200
<b>H-diversity</b>	Shannon Index			<b>0.834</b>	<b>0.001</b>	<b>0.177</b>	<b>0.004</b>
Grouped Sterols	Grouping description						
<b>Sum_C0</b>	No substitution at C-24			<b>0.906</b>	<b>0.001</b>	<b>0.196</b>	<b>0.025</b>
<b>Sum_C1</b>	CH <sub>n</sub> substitution at C-24			<b>0.689</b>	<b>0.001</b>	<b>0.158</b>	<b>0.012</b>
<b>Sum_C2</b>	C <sub>2</sub> H <sub>n</sub> substitution at C-24			<b>0.668</b>	<b>0.001</b>	<b>0.169</b>	<b>0.002</b>
<b>Sum_D0</b>	Phytosterols (Saturated ring B)			<b>0.880</b>	<b>0.004</b>	<b>0.177</b>	<b>0.032</b>
<b>Sum_D5</b>	$\Delta^5$ Sterols			<b>0.777</b>	<b>0.001</b>	<b>0.201</b>	<b>0.007</b>
<b>Sum_D7</b>	$\Delta^7$ Sterols			<b>0.825</b>	<b>0.001</b>	<b>0.188</b>	<b>0.035</b>
Sum_D8	$\Delta^8$ Sterols			0.022	0.114	0.167	0.152
<b>Total sterol</b>				<b>1.002</b>	<b>0.001</b>	<b>0.293</b>	<b>0.026</b>

997

998 **Table 3.** MRM (multiple regression on distance matrices) analysis results showing

999 regression coefficients and p-values for the multiple regression of pairwise distances

1000 on the three first environmental principal components (PC1-PC3) and phylogenetic

1001 distances against the sterol profile Bray-Curtis distance matrix.

Variable	Regression coefficient	p-value
<b>Intercept</b>	7.89E-01	0.24
<b>Phylogenetic distance</b>	2.27E-05	0.82
<b>PC1</b>	1.87E-02	0.0587
<b>PC2</b>	-2.0 E-02	0.0585
<b>PC3</b>	-1.07E-02	0.3635

1002

1003

1004 **Supplementary materials**

1005 **Figures**

1006 **Fig. S1.** 2D-NMDS plot: Pollen sterol profile similarities between species of different  
1007 plant families.

1008 **Fig. S2.** Boxplot: 24-MC content ( $\mu\text{g}/\text{mg}$  pollen) of plants with/without pollen as bee  
1009 reward.

1010 **Fig. S3.** Boxplot: Sterol content ( $\mu\text{g}/\text{mg}$  pollen) of plants with/without pollen as bee  
1011 reward.

1012 **Fig. S4.** GC-MS spectra of the 25 phytosterols identified in our study (after Tri-sil  
1013 derivatisation, extraction details see Materials and methods section).

1014

1015

1016 **Tables**


1017 **Table S1.** Data table (plant species, scores for different environmental  
1018 variables/principal components, pollinator guilds, sterol composition (relative &  
1019 absolute amounts)).

1020 **Table S2.** Scientific name and family for all sampled species, along with suggested  
1021 OTL synonyms (which were subsequently used) and taxon IDs; species excluded  
1022 from the phylogeny are highlighted in grey; reason for exclusion due to issues in the  
1023 data and/or the OTL taxonomy are indicated.


1024 **Table S3.** Variable contributions to axes of PCA of 13 environmental variables.

1025 **Table S4.** Test results: Linear models of phylogenetic independent contrasts (PICs)  
1026 of total sterol amount/diversity against PICs of environmental variables and niche  
1027 breadth.

## a) 24-Methylenecholesterol



## b) Delta-group (double bond position):



## c) 24C-group:




Fig. 1: Chemical structure of 24-methylenecholesterol as an illustration of phytosterols showing a) carbon numbering; b) different substitutions in ring B; and c) different substitutions at C-24.




Fig. 2

Fig. 2: Pollen sterol profiles of plant species. Phylogenetic relationships are given on the left, bold numbers indicate families. Relative contributions of individual sterols to each species' total sterol content are given in the centre; commonness (proportion of plants containing an individual sterol) and relative abundance (average proportion of individual sterol in each species) are given at the bottom; deeper reds indicate values closer to 1. Shannon diversity index (H), 24th carbon groups, delta groups, and total sterol content are given on the right; circle size represents sums of relative sterol contents in the respective groups (0 to 1), and log of g per mg pollen for total sterol content. Sterol names and groups are coloured in the same fashion as illustrated in Fig. 1.

Families: 1 - Pinaceae, 2 - Nymphaeaceae, 3 - Colchicaceae, 4 - Cannaceae, 5 - Strelitziaceae, 6 - Iridaceae, 7 - Asphodelaceae, 8 - Asparagaceae, 9 - Amaryllidaceae, 10 - Ranunculaceae, 11 - Papaveraceae, 12 - Paeoniaceae, 13 - Geraniaceae, 14 - Myrtaceae, 15 - Onagraceae, 16 - Cistaceae, 17 - Malvaceae, 18 - Oxalidaceae, 19 - Salicaceae, 20 - Linaceae, 21 - Euphorbiaceae, 22 - Fagaceae, 23 - Cucurbitaceae, 24 - Rosaceae, 25 - Fabaceae, 26 - Droseraceae, 27 - Caryophyllaceae, 28 - Nyctaginaceae, 29 - Cactaceae, 30 - Hydrangeaceae, 31 - Polemoniaceae, 32 - Theaceae, 33 - Ericaceae, 34 - Primulaceae, 35 - Araliaceae, 36 - Apiaceae, 37 - Adoxaceae, 38 - Caprifoliaceae, 39 - Campanulaceae, 40 - Menyanthaceae, 41 - Asteraceae, 42 - Apocynaceae, 43 - Convolvulaceae, 44 - Solanaceae, 45 - Boraginaceae, 46 - Gesneriaceae, 47 - Scrophulariaceae, 48 - Plantaginaceae, 49 - Bignoniaceae, 50 - Phrymaceae, 51 - Lamiaceae.




Fig. 3: Hypothetical biosynthetic pathways of phytosterols identified in this study (pathways based on Benveniste, 2004).




Fig. 4: Correlation plots of phylogenetically independent contrasts (PICs) of positions on the environmental principal component axes (PC1-PC3) and environmental niche breadth against PICs of total pollen sterol amounts (top row) and sterol profile Shannon diversities H (bottom row). Blue dashed lines indicate regression lines of linear models (with intercept set to zero);  $r^2$  and  $p$ -values for linear models inserted in the respective plot. PC loadings from each environmental variable see Table S3.




Fig. 5: 2D-NMDS plots of sterol profiles for plants (a) with different pollinator guilds, and (b) with/without pollen as bee reward. Distances correspond to sterol profile dissimilarity (Bray-Curtis distances). Stress of NMDS solution: 0.202.