The impact of COVID-19 on diet quality, food security and nutrition in Low and Middle Income Countries: A systematic review of the evidence

Fiorella Picchioni, Luis F. Goulao, Dominique Roberfroid

PII: S0261-5614(21)00395-2

DOI: https://doi.org/10.1016/j.clnu.2021.08.015

Reference: YCLNU 4945

To appear in: Clinical Nutrition

Received Date: 5 May 2021

Revised Date: 22 July 2021

Accepted Date: 18 August 2021

Please cite this article as: Picchioni F, Goulao LF, Roberfroid D, The impact of COVID-19 on diet quality, food security and nutrition in Low and Middle Income Countries: A systematic review of the evidence, *Clinical Nutrition*, https://doi.org/10.1016/j.clnu.2021.08.015.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 The Author(s). Published by Elsevier Ltd.



- 1 Title Page:
- 2 The impact of COVID-19 on diet quality, food security and nutrition in Low and Middle Income
- 3 Countries: A systematic review of the evidence
- 4 Fiorella Picchioni¹, Luis F. Goulao², Dominique Roberfroid^{3, 4}
- 5 1: Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent
- 6 ME4 4TB, UK
- 7 2: LEAF (Linking Landscape Environment Agriculture and Food) Research Centre, Instituto Superior
- 8 de Agronomia, Universidade de Lisboa, Tapada da Ajuda 1349-017, Lisbon, Portugal
- 9 3: Ghent University, Faculty of Bioscience Engineering, Department of Food Technology, Safety and
- 10 Health, Belgium
- 11 4: University of Namur, Faculty of Medicine, Department of Medicine, Belgium
- 12 *Corresponding author: Fiorella Picchioni, Natural Resources Institute, University of Greenwich,
- 13 Central Avenue, Chatham Maritime, Kent ME4 4TB, UK. Email: <u>f.picchioni@gre.ac.uk</u>

14 Abstract

- 15 Background & Aims
- 16 The current global pandemic of Coronavirus (COVID-19), and measures adopted to reduce its spread,
- 17 threaten the nutritional status of populations in Low- and middle-income countries (LMICs).
- 18 Documenting how the COVID-19 affects diets, nutrition and food security can help generating
- 19 evidence-informed recommendations for mitigating interventions and policies.
- 20 Methods
- 21 We carried out a systematic literature review. A structure search strategy was applied in MEDLINE
- 22 (Pubmed[®]), EMBASE[®], Scopus[®] and Web of Science[®]. Grey literature was retrieved by screening a

pre-set list of institutions involved in monitoring the impact of COVID-19 pandemic on nutrition and
food security. The first search was done on 20th August 2020, and updated in mid-November 2020
and mid-January 2021. All research steps were described as recommended in the PRISMA
statement.

27 Results

28 Out of the 2085 references identified, thirty-five primary studies were included. In spite of their 29 heterogeneity, studies converge to demonstrate a detrimental effect of COVID-19 pandemic and 30 associated containment measures on diet quality and food insecurity. One of the major direct effects 31 of COVID-19 on food and nutrition outcomes has been through its impact on employment, income 32 generating activities and associated purchasing power. Other channels of impact, such as physical 33 access, availability and affordability of food provided a heterogeneous picture and were assessed via 34 binary and often simplistic questions. The impacts of COVID-19 manifested with various intensity degrees, duration and in different forms. Factors contributing to these variations between and 35 36 within countries were: 1) timing, duration and stringency of national COVID-19 restriction measures 37 and policies to mitigate their adverse impacts; 2) context specific food value chain responses to 38 domestic and international containment measures; 3) differentiated impacts of restriction measures 39 on different groups, along lines of gender, age, socio-economic status and employment conditions. 40 Dietary changes and food insecurity manifested various intensity degrees, duration and in different 41 forms between and within countries. Shorter value chains and traditional smallholder farms were somewhat more resilient in the face of COVID-19 pandemic. Additionally, the impact of the 42 43 pandemic has been particularly adverse on women, individuals with a low socio-economic status, 44 informal workers and young adults that relied on daily wages. Finally, there were heterogeneous government responses to curb the virus and to mitigate the damaging effects of the pandemic. It has 45 been demonstrated that existing and well-functioning social protection programmes and public 46 47 distribution of food can buffer the adverse effects on food insecurity. But social safety nets cannot 48 be effective on their own and there is a need for broader food systems interventions and

investments to support sustainable and inclusive food systems to holistic achieve food and nutritionsecurity.

51 Conclusion

52 In conclusion, the current economic and heath crisis impact diet quality and food security, and this

raises concerns about long term impacts on access to and affordability of nutrient-rich, healthy diets

and their health implications. Women and individuals with a low socio-economic are the most at risk

55 of food insecurity. Social safety nets can be effective to protect them and must be urgently

56 implemented. We advocate for improved data collection to identify vulnerable groups and measure

57 how interventions are successful in protecting them.

58 Keywords: covid-19, Low- and Middle-Income Countries, Food Security, Nutrition, Social Safety

59 Nets

60 1. Introduction:

The current global Coronavirus pandemic (COVID-19), and measures adopted to reduce its spread, threaten the nutritional status of populations in Low- and middle-income countries (LMICs) (1). Disturbance to the food environments may ensue from changes to both external and personal dimensions of food environments (2). External dimensions include food availability and quality, prices, vendors, markets and regulations, while personal dimensions include geographical access, affordability, convenience and desirability (2).

Children under-5 years and women are expected to be particularly affected by a fall in access to food – particularly for healthy items, such as fruit and vegetables – and by potential disruption of health and nutrition-related programmes and interventions, reducing their access to care (3). The World Food Programme (WFP) has estimated that the COVID-19 pandemic will see more than a quarter of a billion people suffering acute hunger by the end of 2020, which represents a doubling of current figures (4). It was also estimated that even fairly short lockdown measures, combined with severe

73 mobility disruptions and comparatively moderate food systems' disruptions, could result in a 14,3% 74 increase in the prevalence of moderate or severe wasting among children under-5 years across 118 LMICs (5). Reduced coverage of essential maternal and child health interventions could result in an 75 76 increase of 9.8–44.7% in under-5 child deaths per month, and an 8.3–38.6% increase in maternal 77 deaths per month, across 118 LMICs (3). Poorer nutritional status may in turn expose individuals to 78 more severe COVID-19 infections and increase pressure on already vulnerable health systems (3). 79 Hence, the need to prepare and/or strengthen appropriate interventions to mitigate the effect of 80 the pandemic on nutritional outcomes are urgently needed. A response to the COVID-19 pandemic 81 may likely include support to functional and resilient food systems, sustainable healthy diets and access to public health services for all, and particularly the most vulnerable (1,6–9). 82 Such interventions should be guided by evidence. Most of the predictions of the impact of the 83 COVID-19 pandemic on diet quality and the nutritional status of populations are based on macro or 84 85 micro-level simulations (3,5,10,11). The actual impact of the COVID-19 pandemic is still to be 86 quantified. The aim of this research is to provide a preliminary assessment of the multifaceted ways 87 COVID-19 has impacted livelihoods of some of the most nutritionally vulnerable groups and, in turn, 88 their food and nutrition security. Such information is crucial to identify factors that aggravate or 89 mitigate the impact of the COVID-19 pandemic (e.g. geography, characteristics of the food 90 environment, vulnerable individuals), and to target appropriately early interventions. Documenting 91 and disseminating these lessons and emerging evidence will be key to implementing the most

- 92 appropriate and effective interventions in the face of this pandemic.
- 93 2. Material & Methods

A systematic literature review was carried out to identify documented effects of COVID-19 on diet
 quality and nutritional status of children under-5 years and women of childbearing age in LMICs. A

protocol was developed and is available on demand. All research steps were described as
 recommended in the PRISMA statement¹.

98 Evidence was sought against a pre-set list of nutrition and nutrition-related indicators. Eligibility 99 criteria of studies are presented in Table 1. The main outcomes included nutrition, diet quality and 100 food security. Data on other, more distal, indicators (e.g. consumers' behaviours, food availability 101 and affordability) was also collected. Only studies with a design allowing inferences to be made (i.e. 102 including primary data collected since the outbreak of the pandemic) on the impact of COVID-19 on 103 nutrition, diet quality and food security were included. Studies that focused on obesity as a risk 104 factor for COVID-19 infection were not included, as the scope of our research was on the effects of 105 COVID-19 on the nutritional status of individuals and not the reverse. We aknowledge the 106 importance of detailed analysis of the impacts of COVID-19 on overweight and people with obesity 107 in LMICs, which could be addressed in a different study. The search for peer-reviewed studies published up to 20th August 2020 was done in four databases: 108 109 MEDLINE (Pubmed®), EMBASE®, Scopus® and Web of Science®. Search strings can be viewed in 110 Table1-8, Supplementary Materials. MeSH terms were not used as their sensitivity was deemed to 111 be low, given that the literature on COVID-19 pandemic was recent. For retrieving grey literature we 112 screened a pre-set list of institutions for efficiency purpose. The institutions and websites were 113 selected based on their activities in monitoring the evidence-based impacts of COVID-19 on nutrition 114 and food security, publishing sound evidence-based analysis or conducting web-screening and 115 gathering evidence on this subject (see Table 9, Supplementary Materials).

Because of the rapidly accumulating new evidence, we performed an update of the review in mid-November 2020 and mid-January 2021. For pragmatic reasons, the update of the peer-reviewed literature was performed on two of the databases (Scopus® and MEDLINE (Pubmed®), which cover the bulk of the natural science and social science articles published in academic outlets. Due to a

¹ <u>http://www.prisma-statement.org/PRISMAStatement/</u>

- 120 combination of pragmatic approach and time constraints, the update of grey literature was
- 121 conducted for countries and reports already in the list.
- 122 All references were imported into Mendeley (© 2020 Mendeley) where duplicates were detected
- 123 and eliminated. Title and abstract screening was carried out on Rayyan² and irrelevant material
- 124 eliminated. All remaining reports and studies identified as potentially eligible were assessed on full-
- 125 text.
- 126 The quality appraisal of included studies was based on the grids for observational studies proposed
- 127 by the Joanna Briggs Institute³. These grids serve to appraise a number of items (e.g. appropriate
- sampling) in a systematic way with no aim of yielding an overall quality score. Data extraction
- included: 1) information about study reference(s) and author(s); 2) verification of study eligibility; 3)
- 130 study characteristics; 4) study methods; 5) participants; 6) interventions; 7) outcomes measures and
- 131 results. Studies selection, quality appraisal, and data extraction were done by one researcher (FP). A
- second researcher (DR) independently checked a sub-sample of publications and any doubtful
- 133 inclusion/exclusion and the final decision was made by consensus. No meta-analysis was
- 134 undertaken because of the wide variety of study designs and heterogeneity of outcomes reported.

135

Table 1. Inclusion & exclusion criteria for evidence retrieval

Indicators	Inclusion	Exclusion
Setting	LMIC	High income countries
Population	 Under-5y children and women/girls of childbearing age Other individuals 	
Indicators		

² <u>https://rayyan.qcri.org/welcome</u>

³ <u>https://joannabriggs.org/critical-appraisal-tools</u>

Outcome 1:	Higher wasting rate in under-5y	Studies where overweight and obesity
	• Figher wasting rate in under-by	
Nutritional status	Higher Low Birth Weight rate	were considered as risk factors for
	Higher Rate of (micronutrient) deficiencies	COVID-19 infection (reverse causality)
Outcome 2:	Lower Minimum Dietary Diversity for Women	
diet quality	(MDD-W)	
	• Lower Minimum Acceptable Diet for children	
	6-12 (MAD)	
	Lower Household Dietary Diversity Score	
	(HDDS)	
	Lower Food Consumption Score (FCS)	
	Reported changes in quantity and types of	
	food consumed	
Outcome 3:	Higher Household Food Insecurity Experience	
food security	Scale (H-FIES)	
	Higher Household Food Insecurity Access	
	Scale (HFIAS)	
	Changes in Coping Strategy Index (CSI)	
Consumers'	Source of foods (self-consumption, market,	
behaviours	non-timber forest products, etc.), type of	
	markets (open market, supermarket, etc.)	
Food availability	Reduced production	
	Changes in trade flows	
	• Disruptions in transportation of food	
Food affordability	Reduced household income	
	Higher market prices	
Food accessibility	Restricted access to markets	
	Market closures	


Disruption in health	Lower vaccination coverage	
and nutrition	Lower coverage of micronutrient	
services	supplements during pregnancy	
	Decreased treatment of acute malnutrition	
Study design	Longitudinal studies	Projection/predictive/modelling
(For outcomes 1-3)	Interrupted time series/before-after	studies
	design/repeated cross-sectional surveys/	Ecological studies
	trend studies	Individual case studies/series
	Single cross-sectional survey with questions	Opinions
	relating to outcomes before and after the	• Editorial
	pandemic	No data-based analysis
Restrictions	Language: none	
	• Type of studies (quantitative/qualitative):	
	none	
<u> </u>		

136

137 3. Results

Three searches – end of August and mid-November 2020, mid-January 2021 – for both the peer-138 139 reviewed and grey literature were conducted. The first search yielded 1079 and 139 peer-reviewed 140 and grey literature citations respectively, of which 16 (2 peer-reviewed and 14 grey literature 141 studies) were included. The second search yielded 308 peer-reviewed and 48 grey literature papers 142 and reports, of which 11 (5 peer-reviewed and 6 grey literature studies) were retained. The last 143 search in January 2021, yielded 508 peer-reviewed papers and 29 grey literature citations, of which 144 10 (5 each for both types of studies) were included. Therefore, 35 primary studies were included, of which 10 were peer-reviewed and 25 were studies and report retrieved from grey literature sources. 145 146 The overall selection process is presented in the PRISMA flow chart (Figure 1). Excluded studies after 147 full text examination are presented in <u>Annex 3</u>, with reasons for exclusion.

Figure 1. PRISMA flowchart



150

151

152 Table 2 summarises the number of studies based on the selected outcome indicators.

153

Table 2. Number of studies based on selected outcomes indicators

Indicator	Number of studies	Citations
	found	
Nutrition Indicators	0	
Diet quality (DQ) indicators: Household dietary	7	(12–18)
Diversity Score (HDDS); Food Consumption		
Score (FCS) and other non-standardized		
measures		

149

Food Security: Food Insecurity Experience Scale	114	(12–14,18–25)
(FIES); Household Food Insecurity Access Scale		
(HFIAS); direct questions on food insecurity		
perceptions		
Food availability (FAV): changes in agricultural	7	(20,26–31)
operations; changes in trade flows; disruptions		
of food transportation.		<i>C</i> .
Food accessibility (FAC): restricted access to	5	(20,21,27,32,33)
markets	0	
Food affordability (FAF): income decline and	22	(12-22,24,25,27,29,32-
food price increase	.0,1	37)
Disruption in health and nutrition services	4	(20,21,30,35)
(HNS)		

154

155 Most studies were single or repeated cross-sectional studies. The vast majority of them presented 156 the common weaknesses of not describing the sampling process nor the proportion of no respondents. Therefore, included studies were evaluated low quality except the interrupted time 157 158 series study in Bangladesh (25) which was rated high quality. Table 3 summarises the main 159 information of the search results, including outcomes measured and study design. We did not find 160 any study designed to explicitly monitor the diet quality and nutrition of children under-5 years old 161 or and women/girls of childbearing age, although they were the priority groups. Few studies 162 included the gender and urban-rural breakdown of results. In terms of geographical coverage, the 163 selected papers included Bangladesh (two studies) (25,33), Ivory Coast (two studies) (34,35), 164 Ethiopia (15 papers) (12–15,17,19,20,22,26,31), India (two papers) (16,27), Kenya (one paper) (18),

⁴ The studies we report in this table include the 6 rounds of the World Bank high-frequency phone survey conducted in Nigeria and Ethiopia that are referenced once for space reasons.

165 Mexico (one paper) (23), Nepal (two papers) (28,32), Nigeria (seven papers) (21,24), Senegal (one

- 166 paper) (32), South Africa (one paper) (37), Uganda (one paper) (18), Vanuatu (one paper) (29), and
- 167 Zambia (one paper) (30). The 15 studies on Ethiopia include 6 rounds of the World Bank high-
- 168 frequency phone survey conducted between May-October 2020. The 7 studies on Nigeria include 6
- 169 rounds of the World Bank high-frequency phone survey conducted between May-November 2020.
- 170 Each of these 2 batches was referenced in the table as one unique entry for space reason.
- 171 The majority of the studies were longitudinal (7 studies) and cross-sectional phone surveys (13
- 172 studies). The remaining studies were: interrupted time series (1 study), phone exploratory
- 173 qualitative assessment (2 studies) and repeated cross-sectional (3 studies).
- 174 Data extraction tables and quality appraisal can be found in a separate document (available on
- 175 demand).

176

*DQ: Diet quality, FAV: Food Availability, FAF: Food Affordability, FAC: Food Accessibility, FS: Food Security, HNS: Disruption in health and nutrition services, NU:

Nutrition

Sample and unit of analysis	Nationally	Survey round(s) and month	Study Design	Indicators*	Peer Review (P) or
	representative				Grey Literature (G)
2424 Mothers/female	No	1 Round: May to June 2020	Interrupted time-series	FS; FAF	Р
carers ⁵		(compared to baseline 2017-			
		2019)			
1876 households	No	1 Round: 7-15 September	Cross-sectional face-to-face	DQ, FS, FAC,	Р
		2020	and online survey ⁶	FAF	
666 Households	No ⁷	1 Round – April 2020	Cross-sectional telephone	FAF; HNS	G
		22	survey		
3,249 Households ⁸	Yes	6 Rounds (May – October	Longitudinal Phone Survey	FS; FAF; FAC;	G
		2020) ⁹		HNS	
	2424 Mothers/female carers ⁵ 1876 households 666 Households	representative 2424 Mothers/female No carers ⁵	representative2424 Mothers/femaleNo1 Round: May to June 2020carers5(compared to baseline 2017- 2019)1876 householdsNo1 Round: 7-15 September20202020666 HouseholdsNo71 Round – April 20203,249 Households ⁸ Yes6 Rounds (May – October	representativeI Round: May to June 2020 (compared to baseline 2017- 2019)Interrupted time-series1876 householdsNo1 Round: 7-15 September 2020Cross-sectional face-to-face and online survey ⁶ 666 HouseholdsNo ⁷ 1 Round – April 2020Cross-sectional telephone survey3,249 Households ⁸ Yes6 Rounds (May – OctoberLongitudinal Phone Survey	representativerepresentativeI Round: May to June 2020Interrupted time-seriesFS; FAF2424 Mothers/femaleNo1 Round: May to June 2020Interrupted time-seriesFS; FAFcarers5(compared to baseline 2017- 2019)2019)DQ, FS, FAC,1876 householdsNo1 Round: 7-15 SeptemberCross-sectional face-to-face and online survey6DQ, FS, FAC,666 HouseholdsNo71 Round – April 2020Cross-sectional telephone surveyFAF; HNS3,249 Households 8Yes6 Rounds (May – OctoberLongitudinal Phone SurveyFS; FAF; FAC;

⁵ All participants were mothers or female guardians of children enrolled in the "Benefits and risks of iron interventions in children" (BRISC) trial—a randomised controlled trial of preventive iron supplementation or placebo given to infants aged 8 months (ACTRN12617000660381) with a primary outcome of child cognitive development after 3 months of intervention. The BRISC trial was set in Rupganj upazila (county) of Narayanganj district, a rural area about 35 km northeast of Dhaka, which covers about 235 km² and comprises about 82000 households.

⁶ Face-to-face survey was conducted in areas where contagion rates were low (green and yellow zones) while online surveys were conducted in areas with high contation rates (red zones). Data from 1164 (62 %) participants were collected randomly via face-to-face interviews, and data from 712 (38 %) participants were collected using online platforms.

⁷ The study was conducted in 30 districts of Abidjan.

⁸ By the time this study was finalised, the World Bank High-frequency Phone Survey in Ethiopia had conducted 6 survey rounds (early May-October 2020). Each round included a different sample sizes: Round1: 3,249; Round 2: 3,107; Round 3: 3,058; Round 4: 2,878; Round 5: 2,770; Round 6: 2,704.

⁹ Round1: 2020-04-22/2020-05-13; Round2: 2020-05-14/2020-06-03; Round3: 2020-06-04/2020-06-26; Round4: 2020-07-27/2020-08-14; Round5: 2020-08-24/2020-09-17; Round6: 2020-09-21/2020-10-14. Survey Methodology document can be found at the following link: <u>http://documents1.worldbank.org/curated/en/107141590729601148/pdf/Survey-Methodology-</u>Document.pdf.

			Journal Pre-proof			
Ethiopia (14)	600 Households	No ¹⁰	Rounds 1 May 2020	Longitudinal Phone Survey ¹¹	DQ; FS; FAV;	G
					FAF	
Ethiopia (13)	589 Households	No	Round 2 June 2020	Longitudinal Phone Survey	DQ; FS; FAV;	G
					FAF	
Ethiopia (12)	584 Households	No	Round 3 July 2020	Longitudinal Phone Survey	DQ; FS; FAV;	G
					FAF	
Ethiopia (17)	577 Households	No	Round 4 August 2020	Longitudinal Phone Survey	DQ	G
Ethiopia (19)	2,471 young people ¹²	No	1 Round: June-July 2020	Longitudinal Phone Survey	FS; FAF	G
			(compared with 2016 data)			
Ethiopia (26)	100 value chain actors ¹³	No	1 Round: April-May 2020	Cross-sectional Phone	FAV	G
				Survey on qualitative aspects		
Ethiopia (15)	1,188 Households ¹⁴	No ¹⁵	1 Round: June 2020	Cross-sectional phone survey	DQ ¹⁶ , FAF	G
Ethiopia (31)	235 value chain actors ¹⁷	No	2 Rounds: May 2020	Repeated cross-sectional	FAV; FAF	G
		2	(compared with Jan/Feb 2020	phone survey		
			data)			

¹⁰ The study was conducted in Addis Ababa.

¹¹ Longitudinal reports conducted by IFPRI in Addis Ababa (12-14, 17) are part of the same study.

¹² This includes 1,687 Younger Cohort respondents, aged 19, and 784 Older Cohort respondents, aged 25 years old.

¹³ 100 commercial and small dairy farmers dairy processors, traders, development agents, urban retailers, and consumers in rural and urban Ethiopia.

¹⁴ Respondents were all beneficiaries of the fourth phase of Ethiopia's Productive Safety Net Program (PSNP4) and who also participate in the USAID-funded Strengthening PSNP4 Institutions and Resilience (SPIR) project.

¹⁵ The study setting was rural Ethiopia: North Wollo and Wag Himra zones in Amhara, and primarily in East and West Hararghe zones in Oromia.

¹⁶ The study included one question asked about the variations of children's egg and fresh dairy product consumption.

¹⁷ Farmers included smallholders and investors (depending on the amount of land they were renting in) and they resided in the four major vegetable producing districts in East Shewa zone in the Oromia region (Adami Tulu, Bora, Dugda, and Lume). Urban wholesalers operated in Addis Ababa and urban retailers were located in five sub-cities in Addis Ababa.

			Journal Pre-proof			
Ethiopia (22)	436 Households ¹⁸	No	1 Round: July 2020	Cross-sectional phone survey	FA; FAF	G
India (16) ¹⁹	448 Adult men and women ²⁰	No	1 Round: May 2020	Cross-sectional phone survey	DQ; FAF	Ρ
India (27)	1515 farming	No	1 Round (early-April and mid-	Cross-sectional phone survey	FS; FAV; FAC	P
	households ²¹	NO	May 2020)		13,140,140	
Kenya and Uganda	Kenya: 313 & Uganda:	No ²²	1 Round: April 2020	Cross-sectional Online	DQ; FS, FAF	P
(18)	129			Survey		
Mexico (23)	833 Adult men and	Yes ²⁴	3 Rounds: April – June 2020	Cross-sectional Phone	FS	P
	women ²³		010-4	Survey ²⁵		
Nepal and Senegal	Adult male and female	No	1 Round: June to mid-July	Cross-sectional phone survey	FAF, FAC	G
(32) ²⁶	Nepal: 656		2020			
	Senegal: 503					

¹⁸ The population sample included urban poor households and "special segment" population (i.e. particularly vulnerable groups such as day labourers). The study was conducted in 10 selected cities in Ethiopia: Addis Ababa, Mekelle, Dire Dawa, Adama, Gambela, Bahir Dar, Jigjiga, Bulehora, Logia, and Semera. Participants were part of the Urban Productive Safety Net Project (UPSNP), households who own a small-scale business (SSB), and refugees/IDPs/returnees.

¹⁹ The study was conducted in Jharkhand, Assam, Andhra Pradesh, and Karnataka.

²⁰ All respondents were farmers and producing vegetables.

²¹ 1275 farmers in Haryana State and 240 farmers in Odisha State participated in the survey.

²² The questionnaire was sent to random respondents in Kenya and Uganda using social media (WhatsApp, Facebook, Telegram, and Twitter), and via email.

²³ The study was included in 3 survey rounds with different sample sizes: Round1: 833; Round2: 850; Round3: 1,674.

²⁴ The monthly surveys were collected based on a one-stage probabilistic sample of mobile telephone numbers which are randomly selected from the publicly available National Dialing Plan.

²⁵ ENCOVID-19 is a monthly telephone cross-sectional survey, representative at a national level of individuals 18 years and older who have a mobile phone.

²⁶ Farmers in rural Nepal (Dang district of Province 5) and rural Senegal (across the country)

			Journal Pre-proof			
Nepal (28)	25 key informants ²⁷	No	2 Rounds (Mid-April and Mid- October 2020)	Repeated key informant interviews and literature	FAV	Ρ
				review		
Nigeria (21)	1,950 Households ²⁸	Yes	6 Rounds: May-November	Longitudinal Phone Survey	FS; FAC; FAF;	G
			2020 ²⁹		FAV; HNS	
Nigeria (24)	Households and adult	Yes	2 rounds: May, June 2020 +	Panel Data	FS, FAF	G
	individuals Baseline		Baseline Jul/Aug 2018	Å.		
	sample: 4976					
	COVID-19 sample: 1950		Q.Q.			
South Africa (37)	30 Adult male and	No	1 round: month not	Exploratory qualitative study	FAF	Р
	female ³⁰		mentioned (likely to be March	(via phone)		
			or April)			
Vanuatu(29)	31 Adult male and	No	1 round: April 2020	Cross sectional telephone	FAV; FAF	Р
	female ³¹	2		rapid appraisal		
Zambia (30)	40 self-employed	No	1 Round: March – July 2020	Cross-sectional telephone	FAV; HNS	P
	women ³²			semi-structured interviews		

²⁷ Online panel discussion and phone surveys were conducted between mid-April 2020 and mid- October 2020 among 10 government officers and 15 civil society and NGO officials working at different administrative levels.

²⁸ By the time this study was finalised, the World Bank High-frequency Phone Survey in Nigeria had conducted 6 survey rounds (May-November 2020). Each round included a different sample sizes: Round1: 1,950; Round 2: 1,820; Round 3: 1,737; Round 4: 1,691; Round 5: 1,656; Round 6: 1,640

²⁹ Full methodological details are reported here: https://microdata.worldbank.org/index.php/catalog/3712.

³⁰ All individuals were residents of the Informal settlement in the city of Tshwane and they are beneficiaries of relief supplies from humanitarian organisations.

³¹ 22 men and 9 women (>18 years old) who have a position of leadership in the community were interviewed in 23 coastal sites.

³² All respondents were self-employed women and resided in the district of Ndola in Zambia.

177 3.1 Nutritional Indicators

178 No study reporting on nutritional status (e.g. BMI, child stunting and wasting) was found.

179 3.2 Diet Quality Indicators

Seven studies contained information on dietary quality using a variety of indicators (12-18), 180 including: Household Dietary Diversity Score (HDDS) (57), Food Consumption Score (FCS) (58), food 181 182 consumption and diets variations since the start of the pandemic. No study reported Minimum 183 Dietary Diversity for Women (MMD-W) and Minimum Acceptable Diet (MAD). The reviewed studies 184 illustrate that the pandemic had disrupted the diets of the surveyed population, especially at the 185 beginning of the outbreak and the imposition of restriction measures. For example, aggregate HDDS 186 in Addis Ababa deteriorated at early stages of the pandemic (going from 9.3 in January-February 187 2020 to 8.5 in May and June 2020), but returned to pre-pandemic levels in August 2020 as restrictions were lifted (9.4) (12–14,17). However, even when aggregate dietary diversity indicators 188 189 improved as lockdown measures were lifted, all studies suggest that there had been a shift from 190 relatively more expensive sources of calories (e.g., legumes, nuts, animal source foods) to cheaper ones (staples) raising concerns about the long-term effects of COVID-19 on dietary diversity and 191 192 healthy diets (12,13,15–18). In Addis Ababa (Ethiopia), Food Consumption Score (FCS) collected over 193 three survey rounds (May-July 2020) (12–14,17) showed that compared to pre-pandemic baseline 194 (January-February 2020) households were consuming fruit and animal source foods less frequently. 195 In a study conducted in rural Ethiopia (regions of Oromia and Amhara), 70% and 68% of interviewed 196 parents reported children's consumption of eggs and dairy had decreased, respectively, between 197 February and June 2020 (15). In India, Harris et al. (16) showed that, 62% of surveyed farm 198 households reported changes in their diets as a result of COVID-19³³; 17% of households did report a 199 fall in ability to procure staple foods; approximately 50% and 25% reported falls in consumption of

³³ The study was conducted in May 2020, six weeks into the national lockdown and in the early stages of various states' relief packages. The paper reports that the question was asked in binary terms as: "Has your household diet changed as a result of COVID-19?" Therefore, it was unclear if the effects of COVID-19 are intended from when the first cases were registered in India (Jan 2020) or since lockdown measures took place (24th March 2020).

fruit and animal source foods (other than dairy) and pulses respectively. In Kenya and Uganda, a
study conducted in April 2020 reported that 40% and 55% of respondents in the respective countries
changed their diets involuntarily (especially to the detriment of nutritious foods) since the outbreak of
the virus (18).

204 When gendered disaggregated data was available, evidence on dietary degradations, since the COVID-

205 19 outbreak and imposition of restriction measures, showed women were affected to a larger degree

than men. HDDS and FCS in Addis Ababa was consistently lower among female-headed households

between May and July 2020(12–14); in India, women farmers were significantly more likely than men

to report a stronger reduction in consumption of vegetables, fruit, and dairy products (16). Since the

209 pandemic 16% and 30% of women farmers reported that they were eating less and purchasing

210 cheaper foods, respectively, (compared to 5% and 6% of men, respectively)

211 3.3 Food Security Indicators

Eleven³⁴ studies assessed the status of food security since the outbreak of COVID-19 (12–14,18–25).

213 Studies used a variety of indicators including: Food Insecurity Experience Scale (FIES) (59), Household

Food Insecurity Access Scale (HFIAS) (60) and non-standard food insecurity questions. The reviewed

studies agree that levels of food insecurity during the pandemic were high; when pre-pandemic data

were available, food insecurity indicators worsened during the COVID-19 outbreak (18,21,23–25)

- 217 (Table 4).
- 218 219

Table 4 . Examples of household food insecurity variation pre- and during-COVID-19 pandemic in selected countries.

Country	Pre-COVID-19	Post-COVID-19	Indicator
Nigeria (21)	51% (July/Aug 2018)	77% & 68% (Jun & Aug	FIES (Moderate and
		2020)	Severe Food
			Insecurity)

³⁴ The studies we report in this table include the 6 rounds of the World Bank high-frequency phone survey conducted in Nigeria and Ethiopia that are referenced once for space reasons.

Kenya (18)	50%	88% (April 2020)	FIES (% of food
	3370	0070 (7.0111 2020)	1123 (78 01 1000
			insecure households)*
			insecure nousenoius)
	420/		
Uganda (18)	43%	87% (April 2020)	FIES (% of food
			insecure households)*
Mexico (23)	31% (2018)	42% (May 2020)	ELCSA ³⁵ (Mild Food
			insecurity
Bangladesh (25)	6% (2017-2019)	36% (May-June 2020)	HFIAS (Moderate Food
	. ,		
			Insecurity)

220 Source: reproduction using multiple sources (18,21,23,25). *Self-assessed FIES conducted in April 2020.

221 The impact of food insecurity was differentiated among different demographic groups, such as 222 female-headed households, poorer families, young adults and workers in the informal sector. In 223 Addis Ababa, where the percentage of households in moderate and severe food insecurity status in July 2020 reduced compared to May and June 2020 (by approximately 5%), food insecurity remained 224 225 higher among female-headed and poorer households (12–14). In Ethiopia, a longitudinal study 226 conducted among young individuals (June-July 2020) the likelihood of experiencing food insecurity 227 was 27% higher among those that suffered food insecurity in the baseline survey (2016) and that 228 lived in urban areas (19). Similarly, urban vulnerable households, whose survival depends on daily 229 generated income, restrictions and lockdowns led to food insecurity: the percentage of households 230 who consumed an average of three meals a day reduced from 87.6% before COVID-19 to 62.2% at the time of the interview (22)³⁶. A study conducted in Mexico (23) for which socio-economic status 231 232 data disaggregation was available showed that, even though mild food insecurity was present at 233 every SES level, moderate and severe food insecurity increased among lower socio-economic groups.

³⁵ ELCSA is an adapted version of HFIAS and has been extensively validated for Mexico to measure multidimensional poverty (23).

³⁶ The 10 selected cities in Ethiopia include: Addis Ababa, Mekelle, Dire Dawa, Adama, Gambela, Bahir Dar, Jigjiga, Bulehora, Logia, and Semera. The study was conducted among 436 households part of Urban Productive Safety Net Project (UPSNP), households who own a small-scale business (SSB), and refugees/IDPs/returnees. The study had planned to conduct monthly interviews between July-December 2020.

234 3.4 Food Access Indicators

235 Food access information were included in five studies (20,21,27,32,33). Similarly to the previous 236 indicators, food access was not measured in a uniform way among studies. Overall, the reviewed 237 studies indicated food access was affected negatively since the start of the pandemic and hit poorer 238 household to a larger degree (32). Therefore, even if at later stages of the pandemic food access 239 improved (i.e. when lockdown measures are lifted), the percentages of households having difficulties 240 to access food are higher among lower income quintiles (21). In Bangladesh, Kundu et al. (33) 241 illustrated that 45.3% and 61.0% of interviewed households in September 2020 reported that they 242 did not access the same quantity or type of food respectively as they did prior to COVID-19, 243 respectively. The studies that reported access to different food items show that the changes were 244 product specific, suggesting heterogeneous impacts across different food value chains. For example, 245 yams and teff were the commodities less accessible by households in Nigeria and Ethiopia, 246 respectively (20,21). We observed that in Ethiopia there was a recovery a few months after the 247 beginning of the pandemic(20), likely due to the easing of lockdown measures and distribution 248 agricultural inputs.

249 3.5 Food Availability Indicators

250 In total seven studies reported food availability information (20,26–31). Because standardised 251 indicators were not used, the analysis firstly summarizes the evidence on the impacts of the 252 pandemic and lockdown on food value chains and agricultural operations (4 studies (26–28,31)). It 253 then moves to observational data on consumers self-assessment of food availability (3 studies 254 (20,29,30)). The studies on the impacts of COVID-19 and the lockdown on value chains and farming 255 operations provide snapshot of the status of a few value chains: dairy products and vegetables in Ethiopia (26,31); wheat and pulses in India (27); and a more generic overview of Nepal's food system 256 257 (28). These studies suggested that shorter value chains were better placed to survive the pandemic 258 and movement restriction measures. However, poorer farmers living in areas with lack of adequate

```
3.6 Food Affordability Indicators
261
262
       Household income reductions and variations of food prices were selected to evaluate food
263
       affordability. One study, conducted in India (the states of Haryana and Odisha) directly asked
       farmers about affordability since the start of the lockdown (27)<sup>37</sup>. The study shows large differences
264
       between the richer Haryana state, where in the period after the lockdown (April 2020)
265
       approximately 5% of farmers reported difficulties to afford sufficient variety of food, and the poorer
266
267
       state of Odisha, where baseline instances of food unaffordability were already high (approximately
268
       90%), and no significant difference in affordability before and after the lockdown was found. Authors
269
       suggested more diverse cropping patterns, a higher prevalence of homestead gardens, and shorter
270
       value chains for agricultural products helped food affordability of farmers in Odisha (27).
       3.6.1 Reduced income
271
272
       There is much consensus among the studies that the major direct effect of COVID-19 and the
273
       measures put in place by local and national authorities has been through its impact on employment
274
       and, in turn, on income (22 studies (12-22,24,25,27,29,32-35,37)). Between 80% and 58% of
275
       respondents interviewed across Ethiopia, Nigeria, Kenya, Uganda, South Africa and Senegal reported
       either their incomes had decreased compared to the pre-pandemic baseline period directly affecting
276
       their food affordability due to adverse impacts on their regular source of income caused by
277
278
       reduction or closure of business activities, cessation of remittances, and government restrictions
279
       measures (12–15,18,20,21,32,37). In Bangladesh, studies reported income decreased due to COVID-
280
       19 among 71,8 % of respondents (33); 96% of surveyed women reported a reduction in paid work
281
       and median monthly family income fall of 72% (USD 212 was the baseline level) (25).
```

storage facilities and infrastructures were adversely affected by the marketing delays posed by

259

260

restriction measures.

³⁷ The survey included questions on self-assessed food affordability in the month prior to the interview (during the lockdown) and the month before the start of lockdown. Data was collected in April 2020.

Other studies in South East Asia indicated income losses in the first half of the year: farm income reportedly dropped for 90% of vegetable farmers in India (16); a loss of income was reported among 85% of rural Nepalese households interviewed in June and mid-July 2020 (32). Several of the reviewed studies highlighted that the figures are usually higher for informal workers and younger adults in urban areas (19,22,37), and in remote areas (29).

287 3.6.2 Food prices

288 The review of the impacts of the Covid-19 pandemic on food prices suggests a nuanced picture. Food

289 price increases were reported in Nigeria (24)³⁸, where, since the outbreak of COVID-19, the food

290 consumer price index (CPI) increased by 24% and between April/May 2020 90% of households

reported facing food price increase, compared to 85% in mid-March, and 19% between January 2017

and January 2018 (21). In the capital of Côte d'Ivoire (Abidjan), 61 % of respondents reported

significant increases in food prices since the outbreak of the pandemic (35). In Bangladesh (33), 94%

of the 1876 households that took part in the study reported that they faced food price increases due

295 to COVID-19.

296 Other studies suggested a more heterogeneous and variable food price situation, where prices can

297 fluctuate over the course of several weeks. For example in Ethiopia, prices was the main reason to

explain households' inability to purchase food items at the start of the pandemic (approx. 40% of

respondents) (20). However, this gradually decreased so by October 2020, high food prices were

300 mentioned by <10% of respondents (mainly in urban areas).

301 3.7 Information on health and nutrition services disruption

302 Evidence on the impacts of COVID-19 and resulting government restriction measures on the

303 provision of health and nutrition services in LMICs is scattered. Four studies (based in Nigeria, Ivory

³⁸ The Food Consumer Price Index (CPI) employed in the study was collected and constructed by the Nigeria Bureau of Statistics (NBS), which measures the average change in prices over time consumers pay for a basket of food items. Food CPI measures changes in the retail prices of food items and was the principal indicator of changes in retail food prices. It was used to measure consumer inflation in Nigeria's economy. The paper used food CPI for May 2019 and May 2020, corresponding to both survey rounds we employ in this study.

304 Coast, Zambia and Ethiopia) reported reductions of visits to health and nutrition services

305 (20,21,30,35). When rural-urban comparison was available, reductions were more prominent in rural

areas (20). The main reasons for declining recourse to these services included: lack of available

307 medical personnel, movement restrictions and poor transport network lockdown and poor transport

308 network during lockdown and fear of infection.

309 4. Discussion:

310 Impact on diet and nutrition outcomes

311 In spite of their heterogeneity, studies converge to demonstrate a detrimental effect of COVID-19 312 pandemic and associated containment measures on diet quality and food insecurity in a range of 313 LMICs countries. Studies in Nigeria, Mexico and Bangladesh showed a significant deterioration of food security pre- and post-COVID-19 pandemic periods, based on the existence of pre-pandemic 314 315 data (21,23–25). The same trend was reported for Uganda and Kenya using self-assessed food 316 insecurity surveys. The COVID-19 pandemic may have affected diets and food security through 317 various pathways (2). There is a large consensus among the literature that one of the major direct 318 effects of COVID-19 on food and nutrition outcomes has been through its impact on the 319 employment, income and associated purchasing power. This is corroborated by the studies we 320 assessed in the report (12–14,19–22,25,32,33,37) as well as by commentaries and reports produced 321 by international organisations (38,39). However, the link between a fall in income and changes in 322 consumption behaviours and diet quality, although plausible, was not studied as such. Other 323 channels of impact, such as physical access, availability and affordability of food provided a 324 heterogeneous picture and were assessed via binary (and often simplistic) questions 325 (20,21,27,32,33).

A shift from relatively more nutritious foods groups and expensive sources of calories (e.g., legumes, nuts, animal source foods) to relatively nutrient poor and cheaper ones (staples) was observed since the start of COVID-19 (12–18). The production and distribution of perishable and more nutritious

329 foods are often more prone to disruptions during a crisis (40,41). Increased consumption of cheaper 330 sources of calories and decreased levels of dietary diversity, are increasing concerns about the 331 deepening of the triple burden of malnutrition (i.e. undernutrition; overweight and obesity; and 332 micronutrient deficiencies) especially in light of rapid urbanization in LMICs (42,43). Studies have 333 described the incidence of elevated consumption of ultra-processed foods, alcohol and lack of 334 physical activity during lockdown (44,45). While these issues were not included in the primary 335 outcomes of this review, there is urgent need to systematically assess the effects of COVID-19 on 336 overweight and obesity, as a result of changes in consumers' behaviour, access to healthy diets and 337 a general degradation in healthy diets environments.

338

339 Interacting factors

340 Dietary changes and food insecurity manifested various intensity degrees, duration and in different 341 forms between and within countries. Several interacting factors can contribute to these. Firstly, the 342 few studies on food value chains assessments suggested that shorter value chains and traditional 343 smallholder farms were somewhat more resilient in the face of COVID-19. They depended on local inputs (local indigenous seeds, compost, and family and community labour exchange) as opposed to 344 commercial or semi-commercial farms, more severely hit (26,28,46). However, with the exception of 345 346 one article (27), to our knowledge there were no other studies that linked the impacts of COVID-19 347 on agricultural processes and the dynamics and implications on rural households' incomes and food 348 insecurity. Despite food systems thinking and analysis is recognised as an important and meaningful 349 framework to conduct food security analysis, studies tended to focus separately either on food 350 production or on aspects related to food consumption.

Secondly, different food systems actors and groups have experienced and suffered from the
pandemic in different ways. Studies in Ethiopia and India have illustrated that poorer and female
headed households were among those with the lowest levels of dietary diversity and food security

indicators (12–14,20). Moderate and severe food insecurity increased among lower socio-economic
groups.(23) The impact of the pandemic has been particularly adverse on informal workers and
young adults that relied on daily wages (19,22). Given the informal nature of large sections of the
food system in LMICs (where women represent large sections of food processing and sales in wet
and formal markets), assessing the impacts on informal actors and defining targeted policies is
considered a top priority to build back more resilient food systems (42).
Finally, there have been heterogeneous government responses to curb the virus and the timing and

361 stringency of containment measures were variable. Moreover, interventions to mitigate the

362 deleterious effect of the pandemic had also variable timing and intensity.

363 Effective mitigating strategies

364 It has been demonstrated that existing and well-functioning social protection programmes and 365 public distribution of food can buffer the adverse effects on food insecurity during health crises 366 (27,47,48). For example, the evaluation of the Productive Safety Net Programme (PSNP) in Ethiopia 367 during COVID-19 demonstrated that the likelihood of becoming food insecure by 9.3 percentage 368 points in participants (47). A recent simulation on government employment and income protection 369 in Ethiopia has also demonstrated to be effective measures to protect vulnerable population food 370 security during the pandemic (48). But social safety nets cannot be effective on their own and there 371 is a need for broader food systems interventions and investments to support food and nutrition 372 security (49). These include (and not limited to): i) building resilience of health and food systems to 373 withstand shocks such as the COVID-19 pandemic; ii) strengthening and ring-fencing maternal and 374 child essential health and nutrition services so that they are not sacrificed for emergency measures; 375 iii) enhancing nutrition programme coordination and implementation; iv) engaging effectively with 376 young people and women to support both the immediate COVID-19 efforts and the long-term aim of 377 building back better (50). For example, actions on external food environment domains can go from 378 monitoring food prices, diet diversity, food security and malnutrition indicators to adopting subsidies

and taxes that promote the purchase and consumption of nutritious foods based on food-based
dietary guidelines (2,51). Actions on personal food environment domains can encompass improving
accessibility and affordability of foods by social protection programmes, or the promotion of
sustainable healthy diets. Finally, this crisis can represent a window of opportunity for positive
reforms to achieve the SDGs, including: enhancing shorter, sustainable and local food systems;
investing in primary care, especially at the local level; valuing the role of informal workers in the
food system (and other sectors).

386 Challenges and limitations

The situation is still multifaceted and sometimes difficult to interpret. A limited set of studies
included baseline pre-COVID-19 data (17,21,23,25). Without comprehensive longitudinal prepandemic data, it may be difficult to disentangle the effect of the pandemic and annual and seasonal
dietary diversity fluctuations or other factors to COVID-19 (e.g. Orthodox fasting in Ethiopia or
infestations from armyworm and desert locusts (52)).

392 COVID-19 has also posed significant obstacles to collecting information on maternal and child 393 nutritional outcomes (53), or standardized indices such as the MDD-W. We retrieved no data on 394 such outcomes. Diet diversity and food security data were collected via phone and online surveys. 395 While valuable in times of social distancing and movement restrictions such methods may have led 396 to a bias toward easily or quickly 'measurable' or quantifiable data/indicators and respondents 397 accessing digital devices. Also more data is needed from other countries and specific groups, e.g. 398 under-5 children or women of child-bearing age. The nutrition status of populations also needs to be 399 monitored and remote anthropometric assessment be done (54), possibly complemented by COVID-400 19 safe in-person visits. A thorough appraisal of mitigating policies is also needed. We acknowledge 401 that such appraisal is difficult for complex interventions in time of crisis. However, we advocate for 402 improved data collection to identify vulnerable groups and measure how interventions are 403 successful in protecting them.

404

405 5. Conclusion

In conclusion, the current economic and heath crisis impact diet quality and food security, and this raises concerns about long term impacts on access to and affordability of nutrient-rich, healthy diets and their health implications (40,41,55). Women and individuals with a low socio-economic are the most at risk of food insecurity. Social safety nets can be effective to protect them and must be urgently implemented. We advocate for improved data collection to identify vulnerable groups and measure how interventions are successful in protecting them.

412 Acknowledgements

The authors would like to thank Dimitrios Petalios and Mélanie Broin for their editorial and design support; Amélie Wood and Michelle Holdsworth for editorial support and suggestions on earlier versions of the manuscript. We are grateful to the rest of the EU Nutrition Research Facility for their work on defining the methodological approach that put the foundation to the paper. Many thanks to the Teresa Fasig for comments and suggestions on the working paper and longer version of the article.

419 Funding statement

This research was carried out in the frame of the Nutrition Research Facility, which is part of the
Knowledge and Research for Nutrition project of the European Commission. We also acknowledge
the support by Fundação para a Ciência e a Tecnologia (FCT), Portugal through LEAF research unit
(project UID/AGR/04129/2020).

424 Conflict of Interest

425 The authors declare no conflict of interest.

426 Author Contribution

DR and LG: defined the methodology; FP: searched the databases; FP and DR carried out studies
selection, quality appraisal, and data extraction; FP and DR wrote the paper; DR and LG made the
necessary recommendations; and FP, DR, LG revised the manuscript. All authors have read and
approved the final version of manuscript.

431 References

- UNSCN. Food Environments in the COVID-19 Pandemic [Internet]. 2020 [cited 2021 Mar 18].
 Available from: https://www.unscn.org/19?idnews=2040
- 434 2. Turner C, Aggarwal A, Walls H, Herforth A, Drewnowski A, Coates J, et al. Concepts and

435 critical perspectives for food environment research: A global framework with implications for

436 action in low- and middle-income countries. Vol. 18, Global Food Security. 2018. p. 93–101.

- 437 3. Roberton T, Carter ED, Chou VB, Stegmuller AR, Jackson BD, Tam Y, et al. Early estimates of
- the indirect effects of the COVID-19 pandemic on maternal and child mortality in low-income
 and middle-income countries: a modelling study. Lancet Glob Heal. 2020 Jul 1;8(7):e901–8.
- 440 4. WFP. Risk of hunger pandemic as coronavirus set to almost double acute hunger by end of
- 441 2020 [Internet]. 2020 [cited 2020 Mar 18]. Available from: https://www.wfp.org/stories/risk-
- 442 hunger-pandemic-coronavirus-set-almost-double-acute-hunger-end-2020
- 443 5. Headey D, Heidkamp R, Osendarp S, Ruel M, Scott N, Black R, et al. Impacts of COVID-19 on
- 444 childhood malnutrition and nutrition-related mortality. Lancet. 2020;396(10250):519–21.
- 6. GAIN. Impact of COVID-19 on Food Systems: A Situation Report. 2020; Available from:
- 446 https://www.gainhealth.org/resources/reports-and-publications/impact-covid-19-food-

447 systems-situation-report

448 7. FAO. Coronavirus Food Supply Chain Under Strain: What to do? [Internet]. Food Systems

- 449 Transformation. 2020 [cited 2021 Mar 23]. Available from:
- 450 http://www.fao.org/3/ca8308en/ca8308en.pdf
- 451 8. IFAD. Shaping a holistic response to COVID-19: Protecting food systems and rural producers
- 452 [Internet]. 2020 [cited 2021 Mar 23]. Available from:
- 453 https://www.ifad.org/en/web/latest/blog/asset/41863743
- 454 9. The Global Nutrition Report's Independent Expert Group. Global Nutrition Report 2020
- 455 [Internet]. 2020. Available from: https://globalnutritionreport.org/reports/2020-global-
- 456 nutrition-report/2020-global-nutrition-report-context-covid-19/
- 457 10. Hirvonen K, Bai Y, Headey D, Masters WA. Affordability of the EAT–Lancet reference diet: a
- 458 global analysis. Lancet Glob Heal [Internet]. 2020 Jan 1 [cited 2021 Mar 12];8(1):e59–66.
- 459 Available from: www.thelancet.com/lancetgh
- 460 11. Osendarp S, Akuoku J, Bank W, Black R, Headey D, Ruel M, et al. The potential impacts of the
- 461 COVID-19 crisis on maternal and child undernutrition in low and middle income countries
- 462 Results 4 Development Lawrence Haddad Global Alliance for Improved Nutrition. 2020;1–15.

463 Available from: https://www.researchsquare.com/article/rs-123716/v1

- 464 12. de Brauw A, Hirvonen K, Abate GT. Food and nutrition security in Addis Ababa, Ethiopia
- 465 during COVID-19 pandemic: July 2020 report [Internet]. Washington, DC: International Food
- 466 Policy Research Institute (IFPRI); 2020 [cited 2020 Aug 21]. (ESSP WORKING PAPER). Available
- 467 from: https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133851
- 468 13. Abate GT, de Brauw A, Hirvonen K. Food and nutrition security in Addis Ababa, Ethiopia
- 469 during COVID-19 pandemic: June 2020 report [Internet]. Washington, DC: International Food
- 470 Policy Research Institute (IFPRI); 2020 [cited 2020 Aug 21]. (ESSP WORKING PAPER). Available
- 471 from: https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133766
- 472 14. Hirvonen K, Abate GT, de Brauw A. Food and nutrition security in Addis Ababa, Ethiopia

- 473 during COVID-19 pandemic: May 2020 report [Internet]. 2020 [cited 2020 Aug 21]. (ESSP
- 474 WORKING PAPER). Available from:
- 475 https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133731
- 476 15. Alderman H, Gilligan D, Hidrobo M, Leight J, Tafesse AS, Tambet H. Short-term evidence on
- 477 wellbeing of rural Ethiopian households during the COVID-19 pandemic [Internet]. Spir
- 478 Learning Brief. 2020 [cited 2020 Aug 21]. Available from:
- 479 https://www.ifpri.org/publication/short-term-evidence-wellbeing-rural-ethiopian-
- 480 households-during-covid-19-pandemic
- 481 16. Harris J, Depenbusch L, Pal AA, Nair RM, Ramasamy S. Food system disruption: initial
- 482 livelihood and dietary effects of COVID-19 on vegetable producers in India. Food Secur.

483 2020;12(4):841–51.

- 484 17. Hirvonen K, De Brauw A, Abate GT. Food Consumption and Food Security during the COVID-
- 485 19 Pandemic in Addis Ababa [Internet]. Washington, DC: International Food Policy Research
- 486 Institute (IFPRI).; 2020. (Discussion Paper). Report No.: 1964. Available from:
- 487 https://ebrary.ifpri.org/digital/collection/p15738coll2/id/134018
- 488 18. Kansiime MK, Tambo JA, Mugambi I, Bundi M, Kara A, Owuor C. COVID-19 implications on
- 489 household income and food security in Kenya and Uganda: Findings from a rapid assessment.
 490 World Dev. 2021 Jan;137:105199.
- 491 19. Sanchez A, Cueto S, Penny M, Hidalgo A. COVID-19 Phone Survey Headlines Report Listening
 492 to Young Lives at Work in Ethiopia. 2020;(August):1–6.
- 493 20. World Bank. COVID-19 High Frequency Phone Survey of Households 2020 Ethiopia Portal
- 494 [Internet]. LSMS-Supported High-Frequency Phone Surveys on COVID-19. 2020 [cited 2020
- 495 Feb 6]. Available from: https://www.worldbank.org/en/programs/lsms/brief/lsms-launches-
- 496 high-frequency-phone-surveys-on-covid-19#2

497	21.	World Bank. COVID-19 High Frequency Phone Survey of Households 2020 - Nigeria Portal
498		[Internet]. LSMS-Supported High-Frequency Phone Surveys on COVID-19. 2020. Available
499		from: https://www.worldbank.org/en/programs/lsms/brief/lsms-launches-high-frequency-
500		phone-surveys-on-covid-19#5
501	22.	Harris D, Teklu A, Medhin G, Mitiku I, Tadesse B, Demissie M. The effect of COVID-19 and
502		government response measures on poor and vulnerable groups in urban areas in Ethiopia.
503		London, UK: Oxford Policy Management; 2020.
504	23.	Gaitán-Rossi P, Vilar-Compte M, Teruel G, Pérez-Escamilla R. Food insecurity measurement
505		and prevalence estimates during the COVID-19 pandemic in a repeated cross-sectional
506		survey in Mexico. Public Health Nutr. 2020 Oct;1–25.
507	24.	Amare M, Abay KA, Tiberti L, Chamberlin J. Impacts of COVID-19 on food security panel data
508		evidence from Nigeria [Internet]. Washington, DC: International Food Policy Research
509		Institute (IFPRI); 2020 [cited 2020 Aug 21]. (IFPRI DISCUSSION PAPER). Available from:
510		https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133866
511	25.	Hamadani JD, Hasan MI, Baldi AJ, Hossain SJ, Shiraji S, Bhuiyan MSA, et al. Immediate impact
512		of stay-at-home orders to control COVID-19 transmission on socioeconomic conditions, food
513		insecurity, mental health, and intimate partner violence in Bangladeshi women and their
514		families: an interrupted time series. Lancet Glob Heal. 2020 Nov;8(11):e1380–9.
515	26.	Tesfaye A, Habte Y, Minten B. COVID-19 is shifting consumption and disrupting dairy value
516		chains in Ethiopia. In: Swinnen J, McDermott J, editors. COVID-19 and global food security
517		[Internet]. Washington, DC: International Food Policy Research Institue (IFPRI); 2020 [cited
518		2020 Aug 21]. p. 42–5. Available from:
519		https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133815
520	27.	Ceballos F, Kannan S, Kramer B. Impacts of a national lockdown on smallholder farmers'

521		income and food security: Empirical evidence from two states in India. World Dev. 2020;136.
522	28.	Adhikari J, Timsina J, Khadka SR, Ghale Y, Ojha H. COVID-19 impacts on agriculture and food
523		systems in Nepal: Implications for SDGs. Agric Syst. 2021;186.
524	29.	Steenbergen DJ, Neihapi PT, Koran D, Sami A, Malverus V, Ephraim R, et al. COVID-19
525		restrictions amidst cyclones and volcanoes: A rapid assessment of early impacts on
526		livelihoods and food security in coastal communities in Vanuatu. Mar policy. 2020
527		Sep;104199.
528	30.	Nyashanu M, Deborah I, Karonga T, Rumbidzai C. The impact of COVID-19 lockdown in a
529		developing country: narratives of self-employed women in Ndola, Zambia. Health Care
530		Women Int. 2020 Oct;1–14.
531	31.	Hirvonen K, Mohammed B, Minten B, Tamru S. Food marketing margins during the COVID-19
532		pandemic: Evidence from vegetables in Ethiopia [Internet]. Washington, DC: International
533		Food Policy Research Institute (IFPRI); 2020 [cited 2020 Aug 21]. (IFPRI DISCUSSION PAPER).
534		Available from: https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133931
535	32.	Prapti B, Stephanie F, Wouterse, Smriti Saini, Muzna A, Rigler C. COVID-19 challenges to
536		equity: Insights from rural Nepal and Senegal IFPRI : International Food Policy Research
537		Institute. Agrilinks [Internet]. 2020 [cited 2020 Aug 21]; Available from:
538		https://www.ifpri.org/publication/covid-19-challenges-equity-insights-rural-nepal-and-
539		senegal
540	33.	Kundu S, Banna MH Al, Sayeed A, Sultana MS, Brazendale K, Harris J, et al. Determinants of
541		household food security and dietary diversity during the COVID-19 pandemic in Bangladesh.
542		Public Health Nutr. 2020 Dec;1–9.
543	34.	Ministere du Plan et du Developpement and UNDP. Mesure de l'impact socio-economique du

544 COVID-19 sur les conditions des menages en Cote d'Ivoire. 2020.

- 545 35. WFP. Enquête de référence de l'impact du covid-19 sur la situation de la sécurité alimentaire
- 546 des ménages dans le district d'Abidjan. 2020.
- 547 36. Sukhwani V, Deshkar S, Shaw R. COVID-19 Lockdown, Food Systems and Urban-Rural
- 548 Partnership: Case of Nagpur, India. Int J Environ Res Public Health [Internet]. 2020 Aug 7
- 549 [cited 2020 Aug 20];17(16):5710. Available from:
- 550 http://www.ncbi.nlm.nih.gov/pubmed/32784648
- 551 37. Nyashanu M, Simbanegavi P, Gibson L. Exploring the impact of COVID-19 pandemic lockdown
- 552 on informal settlements in Tshwane Gauteng Province, South Africa. Glob Public Health
- 553 [Internet]. 2020; Available from:
- http://www.embase.com/search/results?subaction=viewrecord&id=L2005834431&from=exp
 ort
- 556 38. FSIN Food Security Information Network. 2020 Global Report on Food Crises September
 557 update: in times of COVID-19. Rome; 2020.
- 558 39. FAO. Impact of COVID-19 on agriculture, food systems and rural livelihoods in Eastern Africa.
 559 2020.
- 40. Block SA, Kiess L, Webb P, Kosen S, Moench-Pfanner R, Bloem MW, et al. Macro shocks and
- 561 micro outcomes: Child nutrition during Indonesia's crisis. Econ Hum Biol [Internet]. 2004
- 562 [cited 2021 Mar 12];2(1):21–44. Available from: https://pubmed.ncbi.nlm.nih.gov/15463991/
- 563 41. Huber C, Finelli L, Stevens W. The Economic and Social Burden of the 2014 Ebola Outbreak in
- 564 West Africa. In: Journal of Infectious Diseases [Internet]. Oxford University Press; 2018 [cited
- 565 2021 Mar 12]. p. S698–704. Available from: https://pubmed.ncbi.nlm.nih.gov/30321368/
- 566 42. Béné C, Bakker D, Chavarro MJ, Even B, Melo J, Sonneveld A. Impacts of COVID-19 on People '
 567 s Food Security : Foundations for a more Resilient Food System. 2021;(February).
- 43. Oni T, Micklesfield LK, Wadende P, Obonyo CO, Woodcock J, Mogo ERI, et al. Implications of

	urnal	D		nı	of
U	umar			p.	U

569		COVID-19 control measures for diet and physical activity, and lessons for addressing other
570		pandemics facing rapidly urbanising countries. Glob Health Action. 2020 Dec;13(1):1810415.
571	44.	Ammar A, Brach M, Trabelsi K, Chtourou H, Boukhris O, Masmoudi L, et al. Effects of COVID-
572		19 home confinement on eating behaviour and physical activity: Results of the ECLB-COVID19
573		international online survey. Nutrients. 2020;12(6).
574	45.	Werneck AO, Silva DR, Malta DC, Gomes CS, Souza-Júnior PRB, Azevedo LO, et al. Associations
575		of sedentary behaviours and incidence of unhealthy diet during the COVID-19 quarantine in
576		Brazil. Public Health Nutr [Internet]. 2021 Feb 1 [cited 2021 Mar 16];24(3):422–6. Available
577		from: https://www.cambridge.org/core.
578	46.	Bene C. Resilience of local food systems and links to food security - A review of some
579		important concepts in the context of COVID-19 and other shocks. FOOD Secur.
580	47.	Abay K, Berhane G, Hoddinott J, Tafere K. COVID-19 and Food Security in Ethiopia: Do Social
581		Protection Programs Protect? SSRN Electron J. 2020; (November).
582	48.	Nechifor V, Boysen O, Ferrari E, Hailu K, Beshir M. COVID-19 : socioeconomic impacts and
583		recovery in Ethiopia. EUR 30484 EN, Publications Office of the European Union, Luxembourg;
584		2020.
585	49.	Ayele S, Zegeye EA, Nisbett N. Multi-Sectoral Nutrition Policy and Programme Design ,
586		Coordination and Implementation in Ethiopia. 2020.
587	50.	Save the Children. Nutrition Critical; Why we must all act now to tackle child malnutrition.
588		2020.
589	51.	FAO. Food-based dietary guidelines [Internet]. 2020. Available from:
590		http://www.fao.org/nutrition/education/food-dietary-guidelines/background/en/
591	52.	Integrated Food Security Phase Classification. IPC acute food insecurity analysis in Ethiopia:

592 Belg pastoral and agropastoral areas analysis - September 2020. 2020.

- 593 53. Sam-Agudu NA, Rabie H, Pipo MT, Byamungu LN, Masekela R, van der Zalm MM, et al. The
- 594 Critical Need for Pooled Data on COVID-19 in African Children: An AFREhealth Call for Action
- 595 through Multi-Country Research Collaboration. Clin Infect Dis [Internet]. 2021 Feb 13 [cited
- 596 2021 Feb 22]; Available from: http://www.ncbi.nlm.nih.gov/pubmed/33580256
- 597 54. Viana Bagni U, da Silva Ribeiro KD, Soares Bezerra D, Cavalcante de Barros D, de Magalhães
- 598 Fittipaldi AL, Pimenta da Silva Araújo RG, et al. Anthropometric assessment in ambulatory
- 599 nutrition amid the COVID-19 pandemic: Possibilities for the remote and in-person care. Clin
- 600 Nutr ESPEN [Internet]. 2021 Feb 1 [cited 2021 Feb 21];41:186–92. Available from:
- 601 http://clinicalnutritionespen.com/article/S2405457720311001/fulltext
- 55. Galiano A, Vera-Hernández M. Health Shocks, Household Consumption, and Child Nutrition
- 603 [Internet]. 2008 [cited 2021 Mar 12]. Available from:
- 604 http://www.dnp.gov.co/paginas_detalle.aspx?idp=760
- 605