



# Food systems thinking unpacked: a scoping review on industrial diets among adolescents in Ghana

Winnie Chepng'etich Sambu<sup>1</sup> · Fiorella Picchioni<sup>2</sup> · Sara Stevano<sup>3</sup> · Emmanuel A. Codjoe<sup>4</sup> · Paul Kwame Nkegbe<sup>5</sup> · Christopher Turner<sup>6</sup>

Received: 22 September 2022 / Accepted: 12 October 2023  
© The Author(s) 2023

## Abstract

Unhealthy diets are among the main risk factors associated with non-communicable diseases (NCDs). In Sub Saharan Africa, NCDs were responsible for 37% of deaths in 2019, rising from 24% in 2000. There is an increasing emphasis on health-harming industrial foods, such as ultra-processed foods (UPFs), in driving the incidence of diet-related NCDs. However, there is a methodological gap in food systems research to adequately account for the processes and actors that shape UPFs consumption across the different domains of the food systems framework and macro-meso-micro levels of analysis. This paper interrogates how the Food Systems Framework for Improved Nutrition (HLPE in *Nutrition and food systems. A report by the high level panel of experts on food security and nutrition of the committee on world food security, 2017*), considered the dominant framework to analyse nutrition, and language of interdisciplinarity are practised in research with regards to consumption of UPFs among adolescents in Ghana, a population group that is often at the forefront of dramatic shifts in diets and lifestyles. We conducted a scoping review of studies published between 2010 and February 2022, retrieved 25 studies, and mapped the findings against the domains and analysis levels of the Food Systems Framework for Improved Nutrition (HLPE in *Nutrition and food systems. A report by the high level panel of experts on food security and nutrition of the committee on world food security, 2017*). Our study illustrates that there is a tendency to address unhealthy diets among adolescents in a siloed manner, and as a behavioural and nutritional issue. In most cases, the analyses fail to show how domains of the food systems framework are connected and do not account for linkages across different levels of analysis. Methodologically, there is a quantitative bias. From the policy point of view, there is a disconnect between national food policies and food governance (i.e., trade and regulations) and initiatives and measures specifically targeted at adolescent's food environments and the drivers of UPFs consumption.

**Keywords** Non-communicable diseases · Ultra-processed foods (UPFs) · Adolescents · Ghana · Food systems · System-level analysis

---

✉ Fiorella Picchioni  
F.Picchioni@greenwich.ac.uk

Winnie Chepng'etich Sambu  
winniesambu@gmail.com

Sara Stevano  
ss129@soas.ac.uk

Emmanuel A. Codjoe  
ecodjoe@staff.ug.edu.gh

Paul Kwame Nkegbe  
pnkegbe@uds.edu.gh

Christopher Turner  
C.J.Turner@greenwich.ac.uk

<sup>1</sup> School of Economics, University of Cape Town, Cape Town, South Africa

<sup>2</sup> Natural Resources Institute, University of Greenwich, Chatham Maritime, Kent, UK

<sup>3</sup> SOAS University of London, Department of Economics, London, UK

<sup>4</sup> Department of Economics, University of Ghana, Accra, Ghana

<sup>5</sup> University for Development Studies, Department of Applied Economics, Tamale, Ghana

<sup>6</sup> Natural Resources Institute, University of Greenwich, Food and Markets Department, Chatham Maritime, Kent, UK

## 1 Background

In recent years, high profile reports such as the Sustainable Food Systems Framework for Improved Nutrition (HLPE, 2017) have advocated for interdisciplinary approaches and systems thinking to address the complexities of food systems and the pressing malnutrition challenges at hand. The 2017 HLPE Framework builds on earlier frameworks (Global Panel on Agriculture and Food Systems for Nutrition, 2014; Lawrence et al., 2015), and has been instrumental in enhancing the conceptual understanding of the linkages between food systems, diets, and nutrition. More distinctively, it elaborates on the role of food environments in facilitating sustainable consumer food choices (Herforth & Ahmed, 2015; Turner et al., 2018) making it a key reference point shaping national and international research and policy (see for example Brouwer et al., 2020; Clapp et al., 2022; Fanzo et al., 2020; Nordhagen et al., 2022; Raza et al., 2020; Research Institute (IFPRI), 2019, 2020). Food systems thinking recognises that actions to curb malnutrition need to engage with the complex interlinkages between processes of food production, distribution, retail, consumption, and the natural, technological, political, social, and economic environments within which they are embedded. These interlinkages are manifested across local, domestic and international levels, driven in part by the processes of globalisation, financialization of food and trade liberalisation, shaping the quality of food available to consumers (Baker et al., 2021; Clapp, 2014; Hawkes, 2006).

By mobilising the 2017 HLPE Sustainable Food Systems Framework for Improved Nutrition, this paper investigates how such framework and language of interdisciplinarity are practised in research on the consumption of “industrial diets” among adolescents in Ghana. In addition, we develop a conceptual framework to illustrate horizontal and vertical linkages across and within food systems *domains* and *levels* of analysis emerging out of studies on consumption of industrial diets among adolescents in Ghana. “Domains” are drawn from the 2017 Sustainable Food Systems Framework and refer to the different components and processes through which food is produced, distributed, consumed and disposed of. With the term “levels” we refer to the macro, meso, micro scale of analysis in which the domains are positioned and examined (Tankwanchi, 2018). The section ‘*Building an Evidence and Gap Map*’ (page 5) provides further explanation and expands the definitions of these terms. Finally, we define “industrial diets” as those characterised by high consumption of purchased ultra-processed foods and beverages (hereafter termed ‘ultra-processed foods’ or ‘UPFs’), that through the addition of sugars, salt, fats, and other edible ingredients are made “more durable, palatable, and easier

to handle and therefore help boost sales and reduce costs to processors and retailers” (Winson, 2013, pp. 173–174).

Ghana was classified a lower-middle income country (LMIC) in 2011, however, like many other countries of similar status it is characterised by high levels of poverty and growing inequality. The country is also at the tipping point of the nutrition transition. To be specific, food consumption patterns have transformed dramatically through local production and import of processed and UPFs (Andam et al., 2018). Poor nutritional outcomes affect a large proportion of children and adults. Ghana is experiencing a double burden of malnutrition with the simultaneous high prevalence of overweight and anaemia (Development Initiatives, 2021) as well as significant geographical and socio-economic nutrition inequalities within the country. Undernutrition continues to be common among young children (Jonah et al., 2018), but rising levels of overweight and obesity are becoming a key public health issue (Ghana Statistical Service et al., 2015; Ofori-Asenso et al., 2016; Ziraba et al., 2009). Further, diet-related diseases (e.g., diabetes and hypertension) historically common among adults are increasingly being observed in young children and adolescents (Crouch et al., 2022). A recent meta-analysis of over-nutrition estimates found that approximately 19% of children in Ghana are either overweight or obese (Akowuah & Kobia-Acquah, 2020). Among both children and adults, the prevalence of overweight and obesity is reportedly higher among females (Akowuah & Kobia-Acquah, 2020; Ganle et al., 2019; Manyanga et al., 2014), compared to males. Recent studies suggest high prevalence of overweight and obesity is not only a prerogative of urban areas and rates are increasing also in rural areas (Akowuah & Kobia-Acquah, 2020; Asosega et al., 2021; Mbogori et al., 2020; Muthuri et al., 2014; Ofori-Asenso et al., 2016).

Nevertheless, studies point to increased prevalence of both childhood and adulthood overweight and obesity. The heterogeneity of nutritional indicators and experiences with regards to industrial diets’ consumption, make Ghana a relevant example with regards to the transferability of findings and recommendations to other sub-Saharan Africa (SSA) contexts where NCDs were responsible for 37% of deaths in 2019, rising from 24% in 2000 (World Health Organization, 2022). With the exception of South Africa, research on health-harming industries, such as those that produce UPFs, in driving the incidence of diet-related diseases in SSA is fragmented (Reardon et al., 2021). We argue that such gap has policy repercussions. Food policies in many SSA countries tend to focus on food production, leaving a policy gap in relation to global trade, regulations and the role of the food industry in shaping processing, distribution and consumption (Masters et al., 2018a, b).

Adolescents in the LMICs are often at the forefront of dramatic shifts in diets and lifestyles (Aurino et al., 2017). Evidence suggests that adolescent intake of healthy and nutrient-rich foods such as fruit and vegetables is inadequate (Amoateng et al., 2017), and there is high consumption of high-fat, energy-dense, calorie-rich foods (Keats et al., 2018). Crucially, adolescence is an important phase in human development as it represents a transition from childhood to physical, psychological, and social maturity (Hargreaves et al., 2022; Neufeld et al., 2022; Norris et al., 2022). It includes ages 10 to 19 years and is considered a critical window of opportunity to enhance the health of the next generation (Blakemore, 2019; Prentice et al., 2013).

The effects of poor nutritional outcomes have been widely documented. Malnutrition (undernutrition, micronutrient deficiencies, and overnutrition) has been shown to have adverse negative effects on health and education outcomes in childhood and productivity in adulthood. It also contributes to the vicious cycle of poverty since malnourished children, majority of whom are poor, are more likely to remain malnourished or face other health challenges in adolescence and adulthood, affecting their productivity and participation in the labour market, and contributing to the vicious cycle of poverty (Alderman et al., 2006; Caulfield et al., 2006). Overweight and obesity in both childhood and adulthood increase the risk of non-communicable diseases like diabetes, hypertension, and some types of cancers (Koplan et al., 2005; Park et al., 2012; Simmonds et al., 2016).

Given these adverse effects, evidence on the drivers of malnutrition is necessary to determine suitable interventions. However, existing evidence from Ghana shows that while there are many studies looking at poor nutritional outcomes and links with diets have focused more on the adult population (see for example Amoateng et al., 2017; Benkeser et al., 2012; Bliznashka et al., 2021; Gyasi et al., 2020; Nelson et al., 2015; Obirikorang et al., 2017; Owiredo et al., 2019; Sarfo et al., 2018), and information on the adolescent age group may be limited.

Based on a thorough review of literature from 2010–2022, this paper aims to unpack the extent to which the analysis of recent dietary shifts among adolescents (10–19-year-olds) in Ghana are framed and adequately linked across the various domains and levels of the 2017 HLPE Food Systems Framework. The review maps the evidence and highlights the gaps within studies that have looked at adolescents' consumption of industrial diets in Ghana, with the aim of addressing the following research questions:

- What kind of evidence on UPF consumption among adolescents, quantitative and qualitative, exists from community-level or nationally representative household surveys?
- Which food systems domains and levels of analysis (micro, meso, macro) are included in the reported analysis of adolescents' diets, and more specifically, on UPFs consumption?
- Which disciplines and methodologies are mobilised to improve the understanding of the complex interactions between the different domains and levels of the food systems framework that shape UPFs consumption among adolescents?
- To what extent is the role of the food industry integrated in research on the dynamics around adolescents' food consumption and nutritional challenges?

## 2 Methodology

### 2.1 Literature review methods

This study followed the guidelines for scoping reviews outlined in the PRISMA-ScR checklist (Tricco et al., 2018) – Fig. 7 in the Appendix.<sup>1</sup> We searched three main databases between January 2021 and March 2022: Scopus<sup>®</sup>, Web of Science<sup>®</sup> and PubMed<sup>®</sup>, applying the search terms contained in Table 1. The inclusion criteria used to identify studies in the three databases are outlined here:

- Studies about consumption of UPFs in Ghana. In identifying these foods, we followed the NOVA food classification system that categorises food items based on the processing levels they have undergone (Monteiro et al., 2016).
- Studies about dietary intake among adolescents. Adolescents were defined as children aged 10–19 years, in line with the classification of the World Health Organization (WHO).<sup>2</sup>
- Articles published between 2010 and February 2022.
- Articles published in social sciences (economics, sociology, development studies), public health, nutrition, and agriculture.

We excluded studies from physical and life sciences: engineering, biological sciences, immunology, and biological sciences (including molecular biology), chemistry, mathematics, and machine sciences. In some cases, such as PubMed, we used the following words to exclude articles in the life and physical sciences: biology, bacteria, molecular,

<sup>1</sup> Preferred Reporting Items for Systematic Reviews and Meta-Analyses—Extension for Scoping Reviews.

<sup>2</sup> <https://apps.who.int/adolescent/second-decade/section2/page1/recognizing-adolescence.html>

**Table 1** Literature search terms

Category	Search terms
Food consumption	nutrition transition, unhealthy food, diet, food composition, nutritional intake, food intake, dietary intake
Food supply, availability, and acquisition	food import, food trade, food prices, food purchase, food acquisition, food demand, food consumption, supermarket
Policy environment	food policy, food policies, food governance
Food environment	food system, food environment, food advertising, advertisement, food marketing, food label, food labelling
Food processing	processed food, ultra-processed, packaged food, industrial food, retail food
Commonly consumed ultra-processed foods (identified through an initial review of other studies)	carbonated drinks, sugar sweetened beverage, sugary drinks, sweetened drinks, soda, energy drinks, processed beverages, soft drinks, coca cola, Coca-Cola, coke, juice, cakes, sweets, confectioneries, sweet spread, sweet food, sugar

biological, biochemical, candida, microbiology, microbial, immunology, bacillus, pollution, chemical, malaria, parasite, sexual, viral, virus, bats, antioxidant, lipid.


Our search initially identified 5615 articles. After excluding duplicates, we screened abstracts of the remaining studies (4568). From these, a further 4543 that did not meet the study's inclusion criteria were dropped. The remaining 185 articles were read in detail to assess suitability for inclusion. The in-depth assessment process resulted in the exclusion of a further 160 articles because they did not focus on the review's central theme of processed food consumption among adolescents, although they did cover other food and nutrition security themes such as food production, acquisition, and consumption. In total number, 25 studies were

deemed eligible for analysis. The lead author reviewed all 25 articles in full.

This findings from the search show that research on adolescents and industrial diets has developed over recent years with majority of studies (84%) published after 2014 (Fig. 1). The highest number of publications were recorded in 2019, but this was followed by a decrease in 2020 and 2021, potentially due to the impact of COVID-19 pandemic on research activities.

### 2.1.1 Building an evidence and gap map

To identify themes from the 25 included articles, we applied an Evidence and Gap Map (EAG) lens. More specifically, we



A table showing the number of articles published by year: 2020-2021. The findings presented in Fig. 1 relate to the year of publication rather than year when the data collection was carried out. Analysis in respect to the latter is not possible as nearly half of the studies do not have information on when fieldwork took place (see Table 9 in Appendix).

**Source:** Authors' analyses of reviewed articles.

**Fig. 1** Number of articles, by year of publication

**Table 2** Food systems domains and levels identified in studies on UPF consumption among adolescents in Ghana

Domain	Sub-Domain	Level of Analysis	Sub-categories
Policy environment	Food policy & governance	Macro	Food security & nutrition policies and regulations, including private sector regulations, trade policies, food governance, including public sector structures.
Food availability	Global food supply	Macro	Food imports and exports, concentration and internationalisation of agri-food businesses, financial entities.
	Local food supply chains	Meso	Local production systems, processing, distribution, food safety, retailers, and food sources <sup>a</sup> .
Affordability	Socio-economic status and food cost	Meso/Micro	Socio-economic status, including income and wealth, poverty status, access to pocket money, & food prices.
Desirability	Advertising and Marketing	Meso/Micro	Food advertising (billboards, TVs), food labelling, product placement, sensory norms and food quality issues, cultural norms and habits, cultural and peer influences, consumer knowledge and awareness.
Convenience	Physical access & mobility	Micro	Time or distance travelled to food sources, and modes of transport.
Food consumption	Power	Meso/Micro	Industry influence, e.g., through use of formal/informal channels and school programmes, decision-making around food consumption.
	Point of consumption	Micro	At-home consumption, outside-home consumption, consumption at school, including packed lunch, school feeding programmes, meals purchased inside or vicinity of schools.
	Dietary behaviours & dietary intake		Dietary behaviours (e.g., meal skipping, mealtimes, and snacking), dietary intake (food items consumed), micronutrient intake.
Nutritional and health Outcomes	Nutritional and other Health status	Micro	Nutritional status (stunting, underweight, overweight, obesity), diabetes, hypertension, Other Non-Communicable Diseases.


<sup>a</sup>Retailers and food sources include: i) local retailers such as traditional markets (including fruits and vegetable stalls), fast foods outlets and restaurants, street vendors, convenience stores, supermarkets and; ii) school food environments such as school cafeteria, school-run canteens and shops, school vendors inside and outside schools, school vendors inside and outside schools. Source: Derived from High Level Panel of Experts on Food Security and Nutrition (2017), Herforth and Ahmed (2015), and authors' analysis

drew from the EAG analysis developed by IMMANA (Innovation Methods and Metrics for Agriculture and Nutrition Actions) which summarises the extent and type of research evidence available on food systems and agriculture-nutrition linkages, focusing on tools, metrics, and methods (Sparling et al., 2021). Our analysis focuses on two vectors: i) draws on the sustainable food systems framework domains; and ii) levels of analysis, as outlined in Table 2. The domains draw from the Sustainable Food Systems Framework (Fig. 2) developed by the High-Level Panel of Experts (HLPE) on Food Security and Nutrition (HLPE, 2017). The framework outlines: i) the key drivers that affect food availability, access, utilisation, stability, and sustainability including food supply chains, food environments, and consumer behaviours; and ii) the links with diets and effects on health and nutritional outcomes as well as the broader impacts at the social, economic, and environmental level. In addition, our analysis is broken down by the level of analysis into macro, meso, micro, which refer to the scale in which a domain (or multiple domains) is positioned and examined (Tankwanchi, 2018). For example, the macro level domains include

agriculture and trade policies and institutions that influence, among others, food and agricultural markets, the set-up of food environments, and food availability. The meso level of analysis pertains to the intermediate level between macro and micro level of analysis, or the interface between them, including specific food sectors, local markets, or agricultural practices policies. Finally, micro level domains represent individual and personal factors such as food acquisition and consumption behaviours and preferences, dietary and nutritional outcomes. We recognise there can be interlinkages between micro-meso-macro scales and that domains are not isolated to a single level of analysis.

In Table 2, the first food system domain relates to the policy environment. This domain is about national and international political and institutional actors, that shape ultra-processed food availability, accessibility, and utilisation. The next set of domains are about the food environment. Our definition of food environment draws from the food systems framework (HLPE, 2017) and Herforth and Ahmed's definition and measurement (Herforth & Ahmed, 2015). Food environment domains include food availability, food acquisition patterns,





A conceptual framework showing key elements of food systems: biophysical and environmental drivers; innovation, technology, and infrastructure drivers; political and economic drivers; socio-cultural drivers; and demographic drivers. **Source:** High Level Panel of Experts on Food Security and Nutrition (2017)

**Fig. 2** Food systems framework for improved nutrition


physical access to food (convenience), affordability (income and food prices), and desirability which includes promotion, advertising, cultural norms, dietary habits, and consumer knowledge and awareness. We also include a domain to capture consumer behaviour and food consumption, for example decision making on food consumption, where food consumption occurs, and dietary behaviour. The last domain is the nutritional and health outcomes domain. These are outcomes that are directly linked to dietary intake among other factors, which are the culmination of the interaction between the consumer(s) and the food system.

In addition to these dimensions, the review also examined the research methods and metrics applied in each study, including data sources, research location (geographical setting) and target age groups, among others.

## 3 Findings

### 3.1 Findings at macro-level

Macro-level themes and analysis was only included in two studies, signalling methodological and conceptual gaps in linking consumption of UPFs among adolescents to macro processes, such as food policies, governance, and food supply. It also signals there is a gap of further systematic research on how policies regulate harmful food items, which measures address unclear labelling and harmful advertising of ultra-processed foods directed at adolescents in Ghana. No studies covered themes relating to food production, processing and distribution, trade policies, and food imports. Only two studies (Adom, De Villiers, et al., 2019; Adom,



The number of studies that include meso-level themes: food supply, local retailers, school food environment, affordability, desirability, and power.

**Source:** Authors’ analyses of reviewed articles

**Fig. 3** Research themes within meso-level studies

Kengne, et al., 2019; Pradeilles et al., 2019) covered food policy and governance themes. Adom, De Villiers et al. (2019) and Adom, Kengne et al. (2019) assessed school policies and practises in a study conducted in an urban area of Greater Accra and found that while nearly all schools (six public and six private) had policies and practises on nutrition and healthy eating, these were not associated with decreased likelihood of overweight and abdominal obesity or higher Body Mass Index (BMI) scores. Pradeilles et al. (2019) used key informant interviews with stakeholders in leadership roles in urban areas of Accra and Ho to assess community readiness to implement actions to improve diets of adolescent girls and women of reproductive ages. A community readiness model tool used to assess community awareness of interventions and motivations to implement the programme, showed that both communities were at a “vague awareness stage”, where only a few community members had some information (albeit limited) about local efforts to improve nutrition, community members perceived consumption of unhealthy foods and beverages as a concern but had little motivation to act on this, and availability of resources to mobilise community members towards tackling this issue was limited.

One of the key findings from the macro-level themes is the focus on individuals or community/school. None of the studies examine these themes at the administrative levels

of governance (either national or sub-national level – e.g., regions or municipalities).

### 3.2 Findings at meso-level

Overall, there is more focus on meso-level research themes compared to macro-level themes (see Fig. 3). Meso-level themes were included in 10 studies and covered food availability and points of sale, desirability (advertising and product placement), and power and influences over decision-making on food consumption. The meso-level themes relate mainly to the food environment: where adolescents acquire ultra-processed foods; the marketing and advertising of these foods; types of foods acquired and food safety issues; and modalities through which the food industry influences acquisition and consumption of UPFs. Surprisingly, we found no studies that examined the cost of UPFs consumed by adolescents.

Our review found that purchased foods constitute a significant fraction of adolescents’ food consumption: between 40 and 86% of those surveyed purchased food (Buxton, 2014; Fernandes et al., 2017). While these purchases constitute both UPFs and non-UPF foods, we found that a substantial share of those surveyed by the studies purchased UPFs. For example, Fernandes et al. (2017) reported that 26% of vendors selling foods to children within the school environments sold confectionary. Buxton (2014) reported that 16.5% of

adolescents purchased pastries (e.g., doughnuts, cakes and meat pies), 13.7% purchased ice cream, another 13% acquired soft drinks while 11.7% canned/package fruit juice. Nearly three quarters of studies reviewed focused on where adolescents consumed ultra-processed foods (Abiba et al., 2012; Abizari & Ali, 2019; Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019; Buxton, 2014; Darling et al., 2020; Fernandes et al., 2017; Ganle et al., 2019; Ogum-Alangea et al., 2020; Stevano et al., 2020a, b). These studies show that the main acquisition points include school vendors operating within and outside schools, canteens, fast food outlets, and supermarkets (Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019; Fernandes et al., 2017; Ogum-Alangea et al., 2020; Pradeilles et al., 2021; Stevano et al., 2020a, b). These findings indicate the extent of the proliferation of ultra-processed food within school environments, including through informal channels (such as vendors operating within school proximity).

There are variations in acquisition points and types of foods available in public and private schools, with some of the latter having improved enforcement of who and what could populate the school food environment. For example, Ogum-Alangea et al. (2020) found that, whereas there were 77 food vendors within 100-m radius of five schools (both public and private) included in their study, the vendors were disproportionately located around public schools. Additionally, while in public schools there were hawkers who sold UPFs such as confectioneries, private schools did not have any hawkers within the school environment but had canteens and snack bars instead. Similarly, Stevano et al. (2020b) found that food vendors were not allowed to operate inside private schools, while public schools did not have canteens and food vendors were free to operate within and outside the schools. They also show that in private schools, canteens sold monthly menus selected in advance by children (8–11 years) or their families, something that was not observed in public schools. Adom, De Villiers et al. (2019) and Adom, Kengne et al. (2019) found that private schools were more likely to have school cafeterias and school shops compared to public schools; 74% of private schools had a cafeteria, compared to 26% of public schools. These studies show that differentials in socio-economic status of adolescents (proxied by the types of school they attend) are likely contributors to the differences in dietary behaviours and access to UPFs.

Supermarkets did not emerge as a key UPF acquisition point, despite being considered a key distributor of these items. Only one study (Stevano et al., 2020a) examined the role of supermarkets in adolescent food consumption. They found that together with school canteens, supermarkets were the main sources of UPFs for adolescents from high socio-economic status, while those from low-income households mainly sourced these foods from street vendors.

While several studies focused on where adolescents acquired food, much fewer focused on the types of industrial foods available at purchase points. However, those that studied this found that foods sold to adolescents include ultra-processed food items such as confectioneries, soft and carbonated drinks, packaged fruit juices, biscuits, fried pastries, packaged snacks, chocolates, sweets/toffees, chips & crackers, popcorn, and ice creams (Fernandes et al., 2017; Ogum-Alangea et al., 2020). Healthier options such as eggs, cooked meals, fruits, and vegetables were also available within school environments (Fernandes et al., 2017; Ogum-Alangea et al., 2020). Some studies suggest that there are some differences in types of food items available across private and public schools. Adom, De Villiers et al. (2019) and Adom, Kengne et al. (2019) examined types of foods at cafeterias, shops, and restaurants within one kilometre radius of schools and found that compared to those in public schools, children (8–11 years) in private schools had significantly more access to both healthful (e.g., raw fruits and vegetable salads, cooked meals and 100% fresh fruit juice) and unhealthy foods (e.g., chocolates; sweets/toffees; sodas/soft drinks; packaged fruit juices; cakes, cookies). This finding potentially explains why they also found that those attending private schools had higher BMIs and were at increased likelihood of abdominal obesity and overweight, because they had more access to ultra-processed foods.

Food safety, a key concern for Ghana, emerged in three studies (Pradeilles et al., 2019, 2021; Stevano et al., 2020a), all of which showed that hygiene and food safety affected consumer choices on food acquisition points.

In this review, we found a few studies that focused on food advertising and product placement among adolescents. These studies show that ultra-processed food items are advertised through billboards (Abiba et al., 2012; Bragg et al., 2017; Holdsworth et al., 2020; Stevano et al., 2020a), small signs or displays at front of stores or at storage units (Bragg et al., 2017). Ultra-processed food product constitutes most of the food adverts. In Ghana, the most advertised food products are sugar-sweetened beverages, followed by milk-based sweet drinks targeted at adolescents and other child age groups (such as Nestle's Milo) and Indomie noodles (Bragg et al., 2017; Stevano et al., 2020a). An analysis of 77 photos of outdoor non-alcoholic beverage adverts taken in the Accra area found that over 70% of the adverts were on sugar-sweetened beverages, mainly Coca Cola products (Bragg et al., 2017).

UPFs advertisements and marketing are increasingly targeted at children (including adolescents), partly because they are attentive to food adverts and can recall which food items are commonly advertised and explain why specific adverts attract their attention (Stevano et al., 2020a). To effectively target adolescents and other child age groups, these advertisements are placed within the vicinity of school



environments (Bragg et al., 2017; Stevano et al., 2020a), feature children (including adolescents) (Bragg et al., 2017), and use miniature packaging that is attractive and cost-friendly (Stevano et al., 2020a).

A key issue that relates to food advertising is the influence of food industry in adolescents' decision-making process about food acquisition and consumption. This is a sparsely researched area, as we found only one study that examined this (Stevano et al., 2020a). The findings show that, contrary to existing literature showing supermarkets as a key contributor to the nutrition transition in many LMICs, this is not necessarily the case in Ghana. Instead, in Ghana, the food industry's influence on adolescent food consumption (especially those from poorer socio-economic backgrounds) is occurring through the informal food sector, where street vendors and other traders positioned within school environment sell packaged foods to children (including adolescents). This finding shows proliferation of the ultra-processed food industry through a complex interaction between the formal and informal ultra-processed foods sector.

### 3.3 Findings at micro-level

Micro-level themes were the most popular/dominant across studies. We found that only one study did not contain any of the micro-level research themes such as food acquisition, physical and economic access to food, and dietary habits and food intake.

Within the micro-level themes analysed, food affordability, which is related to the personal domain of food environments, was covered by most studies (Fig. 4). Several studies examined socio-economic status of households as measured by income or wealth indices and explored links with food consumption and nutritional outcomes (Abizari & Ali, 2019; Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019; Aryeetey et al., 2017; Fernandes et al., 2017; Holdsworth et al., 2020; Ogum Alangea et al., 2018; Stevano et al., 2020a, b). In most cases, however, wealth indices were the main measure of socio-economic status possibly due to difficulties in collecting data on household income from adolescents.

Some studies looked at access to pocket money and potential links with dietary behaviours and the types of food items purchased, including UPFs like soft drinks (Abizari & Ali, 2019; Buxton, 2014; Fernandes et al., 2017; Stevano et al., 2020a, b). Fernandes et al. (2017) found that adolescents (10–17-year-olds) were more likely than children aged 5–9 years to bring money to school for food purchases and less likely to consume breakfast or lunch at home. Ultra-processed foods are popular purchase options for adolescents. Abizari and Ali (2019) found that compared to those without, adolescents with pocket money were 4.7 times more likely to have a *sweet tooth pattern*, a dietary pattern

derived through principal component analysis, and which was characterised by intake of: tea and coffee, milk and milk products sugar sweetened snacks, sweets (chewing gums and toffees), energy and soft drinks, and fats and high fat-based foods. Buxton (2014) found that 41.7% of adolescents who purchased food at school bought cooked meals such as fried rice and fried chicken, 16.5% purchased pastries (cake, meat pie, sausage roll, doughnut), 13.7% purchased ice cream (FanYogo or Fan chocolate),<sup>3</sup> 13.0% purchased soft drinks (Fanta, coke, sprite), 11.7% canned/packaged fruit juice while only 3.4% purchased fruit.

Another key aspect of the food environment's personal domain is physical accessibility (convenience), examined in only four studies (Fernandes et al., 2017; Pradeilles et al., 2019; Stevano et al., 2020a). The focus in these studies is on the distance that adolescents and other children travel to and from school and how this impacts the foods they have access to and what they consume. The studies indicate distance from school is a negative factor in healthy consumption behaviour, and those who travel far to school are less likely to consume breakfast. They tend to purchase their foods from street hawkers or vendors within the vicinity of their schools. For example, Stevano et al. (2020a) found that adolescents from lower socio-economic status were more likely not to consume breakfast at home, compared to those from the high-income households, partly due to time and distance they have travel to and from school, as well as the mode of transportation (walking or use of public transport). In comparison, they found that a large percentage of adolescents who consumed breakfast were from higher socio-economic backgrounds and were driven to schools. Similarly, another study (Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019) that compared breakfast consumption among 8–11-year-olds (n = 543) in private and public schools found out that the proportion of those in private schools reporting breakfast consumption was significantly higher than those in public schools (78.3% vs 65.7%). Again, this is likely linked to distance travelled and transportation mode.

Nearly all studies (23) addressed the food consumption domain of dietary patterns and behaviours (Abiba et al., 2012; Abizari & Ali, 2019; Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019; Appiah et al., 2021; Aryeetey et al., 2017; Buxton, 2014; Darling et al., 2020; Fernandes et al., 2017; Ganle et al., 2019; Holdsworth et al., 2020; Intifal & Lartey, 2014; Ogum Alangea et al., 2018; Ogum-Alangea et al., 2020; Oyekale, 2019; Pradeilles et al., 2021; Ross et al., 2017; Sadik, 2010; Siriky et al., 2021; Smith et al., 2021; Stevano et al., 2020a, b; Yang et al., 2017). A breakdown of the food consumption domain shows that the most popular category was dietary intake which was included in

<sup>3</sup> Flavoured ice cream tubs.

all the studies except Fernandes et al. (2017) where focus was on dietary habits (e.g., breakfast consumption, packed meals vs spending at or near school).

Under dietary patterns, breakfast consumption/skipping emerged as an area of interest in some studies, potentially due to the reported positive effects of breakfast consumption on cognitive performance, moods, and nutritional outcomes. A key part of the analysis is the relationship between consuming breakfast at home and consuming purchased food before/on the way to school and whether there are age and socio-economic factors shaping these patterns. A study set in the rural areas of the country found that breakfast consumption at home was reported by 87.6% of children (aged 5–16 years) and mainly consisted of traditional cereal-based meals (Ganle et al., 2019). However, breakfast consumption at home appears to be less common in urban areas of the country. One such study of 820 adolescents in the more urbanised Cape Coast metropolis (Buxton, 2014), found that nearly two thirds (62.8%) of the adolescents did not consume breakfast at home but purchased food during commute to school using pocket money provided by their parents. Adolescents also reported their preference for bought food over home-prepared meals. Stevano et al. (2020a) found that adolescents from lower socio-economic status were more likely not to consume breakfast at home, compared to those from the high-income households, because of the long distances they travelled to school which required them to leave home early to make it to school on time. Another interesting finding that compared younger children (5–10 years) to adolescents found they are less likely to have breakfast or lunch at home and are more likely to bring money to school to buy food (Fernandes et al., 2017).

Our review found that decision-making around adolescents' food consumption is made by a wide range of actors, depending on where and when food is acquired and/or consumed. Buxton (2014) found that over half (54.6%) of adolescents surveyed reportedly made independent decisions about their food purchases within the school environment, 26.3% reported their parents made the decision, 10% were influenced by classmates, while 10% were influenced by siblings or friends. Similarly, through focus group discussions, Stevano et al. (2020a) found adolescents made decisions about food acquisition and consumption mainly in relation to their snacks and foods bought and consumed at school. They found that decisions about meals prepared and consumed at home, e.g., supper were made by parents, often the mother. While adolescents often helped with food acquisition and preparation, they were guided by their parents. Sometimes decisions about foods consumed at home were made by adolescents when their parents or caregivers were away; in these cases, some of the adolescents mentioned consuming packaged foods, such as Nestle's Milo, instead of home cooked full meals.

Studies that focused on dietary intake mainly applied recall methods to inquire about the types and frequencies of foods consumed. In the appendix, we have provided tables showing frequencies of occurrence of foods (see Table 6) and the type of food(s) discovered in each study (see Table 8). Some studies focused on just one or two food items, while others included a range of ultra-processed food items. The most popular ultra-processed food items included in nearly all studies were sugar-sweetened beverages such as carbonated/soft drinks and fruits juices. Studies found a high level of consumption of carbonated soft drinks among 12–15 adolescents, approximately between 54 and 72% (Smith et al., 2021; Yang et al., 2017). However, data on urban-rural breakdown is not available. Consumption of energy and carbonated drinks (23% and 30% respectively) was also high among pregnant adolescent girls (12–19) and reached 71% among the older age groups (16–19) (Appiah et al., 2021).<sup>4</sup> Studies that focused on the links between consumption of soft drinks and gender did not find a significant association (Darling et al., 2020).

Nearly half of studies reviewed collected data on nutritional and health outcomes. No studies covered diabetes incidence or prevalence in adolescents, presumably because the condition is more prevalent among older demographic groups. Table 3 shows that the studies that looked at nutritional outcomes find high and varying levels of overweight and obesity, partly due to the differences in characteristics of geographical settings, sample sizes, age groups included in the studies, and standards used to derive anthropometric indices. Overall, the studies show that prevalence is higher among females than males, and among those attending private schools compared to those in public schools.

Some studies examined the links between nutritional outcomes and consumption of ultra-processed food items. For example, one study (Ogum Alangea et al., 2018) found that an energy dense dietary pattern, which consisted of a typically “westernised diet”,<sup>5</sup> was positively and significantly associated with child overweight status. Another (Ganle et al., 2019) found that children (5–16 year-olds) who consumed fizzy drinks on most days had statistically significant increased odds of being obese compared to children who hardly or did not consume any fizzy drinks at all. Another study (Sirikyi et al., 2021) found that ice cream intake was predictive of overweight/obesity but did not find significant

<sup>4</sup> Both Yang et al. (2017) and Smith et al. (2021) used data from the Global School-Based Health Survey which was collected from adolescents in both urban and rural areas. However, the data is not disaggregated by geo type and so we are unable to compare estimates from urban and rural settings.

<sup>5</sup> The westernized diet included: sugar-sweetened beverages, fried foods, processed meats, spreads and toppings, fruits and fruits juices, cocoa beverages and dairy products, and high calorie snacks.

**Table 3** Overweight and obesity prevalence estimates

<b>Geography</b>	<b>Age group and sample size</b>	<b>Prevalence of overweight and obesity</b>	<b>Growth standards</b>	<b>Source</b>
Accra and Kumasi	9–15 y/o – 3089 from 121 public and private schools	Overweight: 12.4%, obese: 4.6% NB: overweight and obesity rates were higher in private schools (14.2% and 5.8% respectively) compared to public schools (8.3% and 2.5% respectively).	WHO Child Growth Standards, 2007	Aryeetey et al. (2017)
Ga-East Municipality -the Greater Accra	9–15 y/o – 487 from 12 public and 12 private schools	Overweight or obese: 17.7% Separate estimates for private/public schools unavailable.	WHO Child Growth Standards, 2007	Ogum Alangea et al. (2018)
Adentan Municipality - Greater Accra	543 children aged 8–11 years sampled from 14 schools (six public and eight private)	Overweight or obese: 16.4% Abdominal obesity: 18.8% Overweight and obesity prevalence was higher in private schools (12.9% and 1.0% respectively), compared to public schools (5.5% and 3.3% respectively).	WHO Child Growth Standards, 2007	Adom, De Villiers et al. (2019) and Adom, Kengne et al. (2019)
Tema Metropolitan District - Greater Accra	5–16 years – 285 from one public school and one private school	Overweight or obese: 46.9% Prevalence across gender: 42.2% among males and 51.7% among females. 26.8% of children from private school were obese, while 21.4% from the public school were obese.	WHO Child Growth Standards, 2007	Ganle et al. (2019)
Tamale	9–12 years - 100	Overweight: 4%	Centers for Disease Control and Prevention (CDC) growth reference charts, 2000	Abiba et al. (2012)
Tamale metropolis	366 adolescents aged 10–19 years from 10 schools in Tamale metropolis	Overweight or obese: 6%	WHO Child Growth Standards, 2007	Abizari and Ali (2019)
Cape Coast	16–20 years university students	Overweight or obese: The prevalence was higher in females where 27.7% were overweight or obese, compared to 7.08% in males	CDC growth reference charts, 2000	Sirikyi et al. (2021)
All regions – national survey (Ghana Demographic and Health Survey)	Females aged 15–49-year - 9396	Overweight or obese: 40.5% Estimates for adolescent age group unavailable.	BMI calculated by dividing weight (kilograms) by height (metres) squared (kg/m <sup>2</sup> )	Oyekale (2019)
All regions – national survey (WHO Global School Health Survey)	5–19 years - 625	overweight: 11.2%	WHO Child Growth Standards, 2007	Darling et al. (2020)

association between fast food intake and sweetened beverages intake and BMI categories.

### 3.4 Vertical and horizontal food systems linkages

We now examine the extent to which the various domains and levels of analysis classified in this study on ultra-processed food consumption among adolescents in Ghana are connected. We assess both the horizontal linkages (i.e., connections made within the same level of analysis) and vertical linkages (i.e., connections across different levels of analysis). Firstly, we describe how the papers we have reviewed conceptualise the food systems domains and depict (Fig. 5) the linkage between the macro-meso-micro levels of analysis that shape consumption of unhealthy foods among adolescents in Ghana. Detailed information on how many studies connect different domains at different level of analysis can be found in the appendix (see Table 5). Secondly, we conduct a correlation analysis of the food systems domains represented in the papers (Fig. 6).

At the macro level, only the domain of food and nutrition policy in relation to adolescents' UPF consumption was being considered. Other domains such as trade policies, food governance and food supply were not included in the analyses retrieved in this review. With regards to macro-meso vertical linkages, food and nutrition policies were connected to the domain of UPFs availability and food safety. Surprisingly, the macro domain of food and nutrition policies were not linked with the meso domains of food industry influences nor with food advertising and marketing practices. In terms of horizontal linkages, within the meso scale of analysis, food advertising and marketing were considered within the sphere of influence of the food industry. In turn, together with UPF availability, these domains shape what products are available and how they are placed in school food environments and among local retailers, where, especially among poorer socio-economic contexts, operate via informal channels. The vertical linkages that were most common in the scoping review were the meso-micro ones, with virtually all domains being connected in the analyses. Within the micro scale of analysis, horizontal linkages were made between adolescents' socio-economic status and the physical accessibility (especially in terms of convenience and time) to points of consumption or healthy or unhealthy dietary practices (i.e., skipping breakfast or not having breakfast at home). Together with the domains of personal decision-making power and food knowledge and awareness, they these micro-level domains shape food consumption. Finally, nutritional and health outcomes among the most studies micro-domains, which are assessed in relation to food consumption.

Figure 6 depicts a heatmap illustrating the correlation between different food systems domains and analysis levels. Additional information showing the values of the correlation coefficients is available in the appendix (see Table 7). Correlation heat maps are a two-dimensional plot depicting the degree of correlation between variables represented by colours. Correlation ranges from -1 to +1. The diagonals are all orange because each variable is correlated to itself (correlation coefficient = 1). The areas in dark blue on the other hand denote 0 to -1 correlation (meaning, that the two food systems domains are not positively associated to each other). The other colours in between, denote different degrees of correlation. Domains are in turn classified as covering aspects linked to the food systems at the macro, meso and micro levels.

The figure shows an overall low correlation across the different levels of analysis (macro, meso and micro). This means that there is a tendency for research on ultra-processed food consumption among adolescents in Ghana to adopt a clustered analysis approach, where different food systems domains within the same level of analysis are jointly studied. For example, convenience, a meso domain which includes time or distance travelled to food sources, and modes of transport, tends to be associated with the analysis of food supply. Correlation between 0.6 and 0.4 is found between convenience and food source, and between food consumption and place of consumption. However, there are some exceptions. Some domains, although they belong to different levels of analysis, present correlation coefficients between 0.8 and 0.6, meaning that there is a high tendency to observe the analysis of a given domain in combination with another one. An example is food policies and food supply which belong to the macro and meso levels respectively, and which have a correlation of 0.7. Very low correlation levels were found between food policy and governance and economic access (related to socio-economic aspects that shape access to food), and health and nutritional outcomes. This last domain tends to be analysed either individually or in conjunction with the food consumption domain only. Interestingly, we found that the analysis of power (that in this paper includes food industry influence, e.g., through use of formal/informal channels and school programmes as well as food choices autonomy) was not related to the food policy and governance domain nor health and nutritional outcomes.

### 3.5 Tools, methods, metrics and geographies commonly adopted in analysing adolescents' diets

Majority of studies (84%,  $n=21$ ) used quantitative methods, while only seven used qualitative methods (Table 4). Majority of studies (57%) were cross-sectional studies. We did not identify any panel (longitudinal) studies on adolescents' dietary patterns and outcomes and their associated factors

**Table 4** Number of publications, by metrics

Domains	Categories	Number of studies	% (out of 25)
<b>Methods</b>	quantitative	17	68.0%
	qualitative	4	16.0%
	mixed methods research	4	16.0%
<b>Data sources</b>	primary research (fieldwork)	22	88.0%
	secondary data sources	3	12.0%
<b>Measurement unit</b>	individual	25	100.0%
	household	0	0.0%
	community (including schools)	4	16.0%
	points of sales (e.g., shops)	0	0.0%
	other	0	0.0%
<b>Data collection point</b>	household/residence (home)	8	32.0%
	schools	16	64.0%
	community	2	8.0%
	not stated	1	4.0%
<b>Geographical area of focus</b>	urban	17	68.0%
	peri-urban	2	8.0%
	rural	2	8.0%
	all	3	12.0%
	not specified	5	20.0%
<b>Adolescent age group</b>	10 -14 years	20	80.0%
	15–19 years	18	72.0%
<b>Other focus age groups (in addition to adolescents)</b>	under 5-year-olds	2	8.0%
	5–9-year-olds	9	36.0%
	college students	1	4.0%
	adults	9	36.0%
	age not specified	1	4.0%
<b>Gender</b>	male	0	0.0%
	female	5	20.0%
	all	19	76.0%
	not specified	1	4.0%


Source: Authors’ analyses of reviewed articles

over time. Overall, primary research was the main source of data, used by 22 out of 25 of research articles. All studies that employed qualitative and mixed methods were based on primary research. Similarly, most of the quantitative methods research (82%) used primary sources of data, although a few articles (n = 3) used secondary data. Overall, only three studies made use of secondary data (Oyekale, 2019; Smith et al., 2021; Yang et al., 2017). Two of the studies (Smith et al., 2021; Yang et al., 2017) both used data from the Global School-based Student Health Survey (GSHS), a school-based surveillance survey led by the World Health Organisation (WHO) and United States CDC. The other, (Oyekale, 2019), used data from the 2014 Ghana Demographic and Health Survey.

Studies that applied quantitative methods mainly used structured questionnaires to collect data on socio-demographic characteristics (see Appendix, Table 8). For studies

that collected dietary intake data, the main data collection instruments used were food frequency questionnaires, with either 24-h or 7-day recall periods. Some studies used longer recall periods. For example, Abiba et al. (2012) applied a two-week recall period, while Darling et al. (2020) used a 30-day recall and Ross et al. (2017) used a 6-month recall period. These recall periods are particularly lengthy for adolescents, especially in cases where they are asked about quantities consumed and frequency of consumption within and outside the home. Additionally, while most studies on dietary intake collected data directly from adolescents or other child age groups where applicable, one study (Fernandes et al., 2017) collected the data from caregivers. Studies should include robust, piloted and age adequate protocols and methods and training to collect food consumption data from adolescents to reduce the risk of recall bias and ensure accurateness of the data. The number of food items included in food consumption





The number of studies that have micro-level themes such as economic and physical access, advertising and marketing, power, point of consumption, dietary behaviour and dietary intake, and nutritional and other health status/outcomes.

**Source:** Authors' analyses of reviewed articles


**Fig. 4** Research themes within micro-level studies

studies varied significantly. One study used a checklist of 229 food items (Holdsworth et al., 2020), another 60 food items (Abizari & Ali, 2019), another 100 food items (Ogum Alangea et al., 2018) while those based on GSHS data focused only on carbonated soft drinks (Smith et al., 2021; Yang et al., 2017).

The sample sizes used in the studies vary. In one case, a study that looked at food consumption in orphanages included only 40 participants purposively sampled (Sadik, 2010), while another study included all first-year students (196) who reported for medical test at a university in the Cape Coast region (Siriky et al., 2021). Other studies had much larger sample sizes. An example is a mixed-methods study (Fernandes et al., 2017) which examined children's food purchases. This study surveyed 4285 children (aged 5–17 years) in 111 schools sampled from 60 districts from the 10 regions of Ghana. However, the study covered only urban and peri-urban areas of the country. Majority of the studies do not have nationally representative samples, and those that do use secondary sources of data such as the Ghana Demographic and Health surveys (Oyekale, 2019).

In terms of geographical focus, most studies were carried out in urban areas, with some popular locations including Accra, Cape Coast, and Tamale metropolitan areas. More precisely, seven studies were conducted in Accra, five in the greater Accra regions (Ga East and Adentan municipalities) while four studies were conducted in the Cape Cost and three in Tamale. Only two studies (Darling et al., 2020; Intiful & Lartey, 2014) were conducted in rural areas alone (Manya Krobo in Eastern region and Ningo Pram-pram in Greater Accra respectively). Three studies (Bragg et al., 2017; Oyekale, 2019; Yang et al., 2017) covered both urban and rural locations (Accra & Cape Coast in Bragg et al. (Bragg et al., 2017) and national samples for both Yang & Oyekale (Oyekale, 2019; Yang et al., 2017). The large number of studies focusing on urban areas is not surprising given the higher (reported) consumption of ultra-processed foods in urban locations, compared to rural settings, in many countries in the Global South.

Nearly all studies (24) included the individual (adolescent) as the unit of measurement. Four studies used a school environment (Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019; Fernandes et al., 2017; Ogum-Alangea



A conceptual framework showing how food systems domains are conceptualised and depicted in the studies reviewed, and the linkages that emerge across level of analysis (macro-meso-micro).

Source: Authors’ analyses of reviewed articles

Fig. 5 Two-vector conceptual framework of the analysis of UPF consumption among adolescents in Ghana


et al., 2020) as the unit of measurement, while one study used outdoor spaces, including a national highway, as the unit of analysis (Bragg et al., 2017). We found that only one study conducted a mapping exercise on obesogenic environments in residential areas (Holdsworth et al., 2020). We found that in most of the studies (64%), data collection was conducted from within school environments, while in eight it was done at household residence. Majority of the studies do not exclusively cover adolescents but include them as part of a broader age group. There were 17 studies that included at least one other age group in addition to adolescents. The most common additional age group was 5–9-year-olds (Abiba et al., 2012; Adom, De Villiers, et al., 2019; Adom, Kengne, et al., 2019; Fernandes et al., 2017; Ganle et al., 2019; Intiful & Lartey, 2014; Ogum-Alangea et al., 2020). Some studies focused exclusively on the adolescent age group and did not include any other demographic age group (Abizari & Ali, 2019; Aryeetey et al., 2017; Buxton, 2014; Darling et al., 2020; Smith et al., 2021; Yang et al., 2017). Nearly all studies included both male and

female adolescents, and only three focused on female adolescents only (Oyekale, 2019; Pradeilles et al., 2019; Ross et al., 2017). Only one study (Bragg et al., 2017) did not have information on gender as the community was the measurement unit.

More than half of these studies (56%) compared public and private schools in respect to food environments, dietary patterns, and adolescents’ nutritional outcomes. However, a few studies that focused on schools did not specify whether the schools were public or private (Abiba et al., 2012; Abizari & Ali, 2019) while two studies focused only public schools (Fernandes et al., 2017; Intiful & Lartey, 2014).

#### 4 Discussion

The findings show that there are key gaps in food systems research in Ghana. Firstly, we find that, methodologically, there is a quantitative bias and absence of qualitative research that provide deeper understanding of the factors



A heatmap illustrating the correlation between different food systems domains and analysis levels.

**Source:** Authors' analyses of reviewed articles

**Fig. 6** Correlation heat matrix showing overlap across themes

underlying consumption of industrial diets among adolescents. Second, studies have mainly focused on urban areas. Third, not all aspects of food systems domains are covered equally. The macro-level themes, such as food policies and food supply, are least studied. Importantly, when it comes to adolescents' dietary patterns and nutritional outcomes, in the context of changing food environments research and practice, the literature tends to focus on the individual domain (i.e., links between food consumption and nutritional outcomes), and remains disconnected with the whole complexity of the food systems framework. In particular, the activities, and repercussions of the food industry on adolescents' food choices and practices are neglected. Future studies should focus on these under-studied domains of food systems and examine linkages between macro, meso, and micro levels of analysis. More focus should also be paid to qualitative and/or mixed methods approaches to enhance interdisciplinarity in this area of research. Finally, more attention should be given to neglected geographical settings, i.e., such as peri-urban and rural areas and the contextual marketing processes through which ultra-processed foods penetrate even the most remote food environments (i.e., via informal channels).

In recent years research and practice on nutrition has adopted the language and practice of interdisciplinarity and systems thinking. Distinctively, the HLPE report on Sustainable Food Systems Framework for Improved Nutrition (HLPE, 2017) identifies the functions of and interactions between food supply chains, food environments, and consumer behaviour as entry and exit points for nutrition and health outcomes. It recognises that the challenge to ending malnutrition is a complex and multi-layered endeavour that requires strategies to reverse the current paradigms of food production and distribution that rely on supplying overabundant energy-dense and nutrient-poor foods and damaging the environment. To meet these challenges research and practice should therefore adopt: 1) a multi-scalar approach, i.e., interlocking and relating different aggregation levels, from the individual to the global, passing through the household, community, national and global level; 2) practice interdisciplinarity that improves the understanding of complex problems, instead of studying them from a single discipline perspective. Whilst these connections are theoretically outlined, in practice, there is an operational and empirical gap in articulating the response mechanisms between the micro level factors at the "inner core" of these frameworks (e.g., food consumption, nutritional and health outcomes) and

meso (e.g., local food markets) and macro level factors (e.g., food policy and governance, and regional and global trade).

For example, in this scoping review no study covered themes relating to trade policies, which is a key research gap considering the role that trade liberalisation has played in globalisation of diets and increased availability of ultra-processed and unhealthy foods available to adolescents in the Ghanaian markets (Hawkes, 2006; Hawkes et al., 2012). We also find that where studies have examined food policies, they were few and only covered individuals or school environments. We found no studies that delved into how food policies at national or sub-national levels, where they exist, apply to the adolescent age group. For example, a key gap is the absence of research linking adolescent's unhealthy-food consumption and nutrition with food regulation, necessary to ensure domestic and imported food products are safe, sanitary, nutritious, wholesome, and properly labelled.

When it comes to adolescents' exposure to unhealthy diets, the policy environment is one of the most under-researched domains. There are existing policies in Ghana that speak to various aspects of food security; a recent policy review found 23 policy documents that either mention or are centred on food and nutrition security, from poverty reduction and economic development, to agriculture, nutrition and health, spatial demographic development, climate change, and trade (Linderhof et al., 2019). Examples include Food and Agriculture Sector Development Policy (Ministry of Food & Agriculture, 2007), and Support to the Planting for Food and Jobs Campaign (FAO, 2020). As is evident, many of these policies are about food production. The National Policy for the Prevention and Control of NCDs in Ghana, published in 2012, focuses on non-communicable diseases and lays out measures to reduce the incidence of NCDs and exposure to risk factors (Ministry of Health, 2012). The policy speaks to the role of unhealthy diets in driving the prevalence of NCDs and suggest control measures like regulating advertising and introducing price controls to discourage consumption of unhealthy foods. The policy mentions intent to discourage sale of unhealthy foods, such as fizzy drinks, in school canteens and compounds and replace them with healthier alternatives such as fruit. More recently, the government published an updated policy on NCDs (Ministry of Health, 2022). The policy focuses on the need for prevention and management of NCDs at primary, secondary and tertiary levels through nutrition interventions, among others. The policy emphasises the need to strengthen the legal regime to prevent NCDs including through regulation of labelling, advertising, and sale of UPFs such as sugar sweetened beverages. Laar et al. (2020) argue that the existence of such policies demonstrates recognition by the government and indicates some level of political will to improve food environments. However, the key challenge remains lack of implementation which hinders progress towards realising

local and global targets (Laar et al., 2020). Another key gap is the unavailability of data on government fiscal allocations towards nutrition and health interventions to support access and utilisation of healthier foods by adolescents.

At the meso-level, we find that advertising through television or print and digital media did not emerge as a key research theme in any of the studies. Some studies examined the extent of television viewing and explored links with sedentary lifestyles, but we did not find any studies that linked food advertising on television to adolescents' dietary intake.

Power features at two levels of analysis: 1) at the meso-level, where power is examined in the context of practices of the ultra-processed food industry to shape consumption of less healthy foods; and 2) at the micro-level, in terms of decision-making power on foods that are consumed by adolescents. These two conceptualizations of power have overlapping features and are interdependent, and research in this area is still sparse. We identify two other missing pieces in the analysis of power in this context: a) how national food regulation (and its absence) shapes the operation of unhealthy food industries and their impact on adolescents' nutrition; and b) how neoliberal policies account in part for some recent trends in the diets of adolescent population (i.e., foreign direct investment and global food marketing strategies). Specific dietary outcomes reflect context-specific socioeconomic and cultural features in which these policies are operating (Hawkes, 2006).

Unsurprisingly, a major gap identified by this review is the disregard of the food industry's operations in the research on adolescents' nutrition in Ghana. Despite their growing influence, food industry practices are discussed only rarely and research on the impacts of industrial food production on nutrition in the SSA is nascent (Nestle, 2013; Popkin, 2014). Food industries remain a neglected and disconnected sector in the understanding of food systems and nutrition interaction in SSA (Reardon et al., 2021), where interventions to lower consumption of UPFs are typically limited to education and information campaigns (Monteiro et al., 2013) and agricultural policies, especially in the African context, targets production (Masters, Bai, et al., 2018; Masters, Rosenblum, et al., 2018).

Our study demonstrates research in this area among Ghanaian adolescents prioritises analysis that covers individual aspects related to food consumption and dietary patterns and behaviours, and health and nutritional outcomes. In doing so, the transformations of food systems and their impacts are predominantly studied at the micro level and focus on the lower right areas of the food systems framework. This is in line with viewing food and diets as medicalised and individualised issues, that require behavioural change, personalised solutions and individual economic incentives (Scrinis, 2016). The joint expansion of medical science and the food industry over the past century have

produced the conditions in which eating and feeding are transformed from practices embedded in social, political and economic relations into explicit medical practices (Mayes, 2014). In doing so, "solutions" and interventions put forward, especially in the SSA context, are disconnected with the political economy and contextual specificities of malnutrition (Sathyamala, 2016).

One of the main observations that emerged during our review is that existing research on food consumption focuses on a small number of processed food items. The most popular items being fizzy and sweetened drinks, and fried pastries, perhaps due to their popularity among adolescents. We did encounter many studies that discussed food consumption, but they mainly used broad groups to examine dietary diversity and micronutrient composition, and mainly focused on preschool children and/or pregnant women (Ayensu et al., 2020; Bimpong et al., 2020). We suspect that this is partly due to the challenges of collection of dietary intake data: 1) the high cost of administering food consumption surveys and 2) the impact of long recall periods on accuracy of the data. Lack of detailed data on adolescent's dietary intake inhibits a comprehensive understanding of the state of ultra-processed food consumption among adolescents and the underlying drivers including across geographical locations (urban/rural) and socio-economic levels.

An additional finding that emerged from our analysis is the limited focus on ultra-processed food consumption among adolescents living in peri-urban and rural areas of Ghana. This is a key research question that needs further exploration given the strong linkages between urban and rural areas and as rural locations become increasingly urbanised. Increased advertising and growing availability in rural contexts have also likely contributed to increased demand for these foods.

Some studies show that increased demand for processed foods may be due to lower cost, compared to healthier alternatives, in addition to other factors like taste and convenience. We were unable to establish if this is the case in adolescents' dietary intake in Ghana as we found no study that examined food prices of ultra-processed foods consumed by them. There are some studies that have collected general food price data in Ghana (Masters, Bai, et al., 2018; Masters, Rosenblum, et al., 2018; Minot & Dewina, 2015), but they do not necessarily focus on UPFs, particularly those consumed by adolescents within their environments (home or school). It is possible that the administrative costs of collecting food price data may be contributing to this. This is a key research gap that needs further exploration. A related issue is that we did not find any studies that linked supermarket purchases to adolescent processed food consumption despite interest in the supermarket revolution in the Global South. In Ghana, research indicates that there has been a gradual increase in household food purchases from supermarkets (Meng et al., 2014). The studies we encountered on

food purchases in supermarkets focus on purchases by other demographic groups or the understanding of food labels and composition and not necessarily in respect to processed foods or adolescents (Aryeetey & Tay, 2015; Darkwa, 2014; Owureku-Asare et al., 2017).

## 5 Conclusion

This scoping review unpacks the disconnects present in food systems research by mapping the evidence and identifying the gaps within research undertaken on consumption of ultra-processed foods among adolescents in Ghana. More specifically, we focused on which food systems domains are more dominant, paying attention to which levels of food systems analysis (micro, meso, or macro) is more prevalent.

Our study is inclined to suggest that there is still the tendency to address consumption of unhealthy diets among adolescents in a siloed manner and that the analysis tends to fail to join the dots across the food systems framework. We also found that methodologically, there is a quantitative bias in this area of research. Despite the increasingly recognised need to systemically address harmful food consumption across the different food systems domains, research tends to remain focused on the micro actors and factors shaping food production, distribution, consumption, and waste. While the food systems approach has helped drive research and practice towards a more interdisciplinary and multi-scalar space, there are no clear guidelines to operationalise these very needed approaches. Putting in practice food systems thinking requires a deeper engagement with disciplines and methods that tend to be neglected in this area. They include mixed methods and participatory practices coupled with political economy analysis that places the politics of food at the heart of nutritional outcomes (Béné, 2022; Walls et al., 2021).

Our research showed blind spots in food systems research in Ghana and that not all food systems/environment domains are covered equally. This is particularly stark when it comes to the proliferation of unhealthy diets among adolescents, which highlights the depth of the micro-macro gap in this area of study and absence of the role of the food industry in shaping food acquisition. The disconnects within food systems research are manifested in policy. In the Ghanaian context food policy to tackle malnutrition tends to operate from a lens of food security and food production. Only recently has there been a growing recognition of the need to promote policies that take an integrated food environments approach, where food industries are playing a central role. But financial resources and political will are lagging. This is particularly problematic when it comes to adolescents' dietary patterns. The focus of research primarily on micro and individual level nutritional and health outcomes, produces policies that treat



unhealthy food consumption patterns as an individual issue that requires interventions to discourage the individual consumption of unhealthy foods. With regards to adolescents, the cross-sectoral cooperation and coordination between agri-food sector, health and education is crucial to promote

healthy and sustainable diets. Policy design and research should recognise that national and local food systems are positioned within global political economy processes that shape food environments, food access, tastes, and aspirations in multiple ways (Fig. 7).

### Appendix




Fig. 7 PRISMA Chart - evidence on industrial diets among adolescents in Ghana

**Table 5** Number of studies that connect food systems domains at different level of analysis

		Level of analysis		
		Macro	Meso	Micro
<b>Domains</b>	<b>Food and trade policies</b>	2	1	2
	<b>Food supply</b>	2	4	4
	<b>Food sources</b>	1	6	6
	<b>Affordability</b>	1	6	11
	<b>Convenience</b>	1	4	4
	<b>Desirability</b>	1	7	8
	<b>Power</b>	0	2	3
	<b>Point of consumption</b>	1	8	18
	<b>Dietary behaviours &amp; dietary intake</b>	1	8	23
	<b>Health outcomes</b>	1	2	11

Source: Authors' analyses of reviewed articles

**Table 6** Tabulation of UPFs derived from the research papers and classified using the NOVA classification (Monteiro et al., 2016)

	Frequency	Percentage
confectionery <sup>a</sup>	37	28.7%
ready-to-heat products (savory) <sup>b</sup>	28	21.7%
sweetened beverage	17	13.2%
carbonated soft drinks	16	12.4%
margarines and other spreads	8	6.2%
sweetened milk drink	8	6.2%
ice cream	4	3.1%
sugar	3	2.3%
packaged breads and buns	3	2.3%
energy drinks	1	0.8%
cereals	1	0.8%
instant sauces	1	0.8%
other	2	1.5%
<b>Total</b>	<b>129</b>	<b>100%</b>

Source: Authors' analysis

<sup>a</sup>Confectionery: sweet or savory packaged snacks; chocolate, candies, cookies (biscuits), pastries, cakes, and cake mixes

<sup>b</sup>Ready-to-heat products (savory): pies and pasta and pizza dishes; poultry and fish 'nuggets' and 'sticks', sausages, burgers, hot dogs, and other reconstituted meat products; and powdered and packaged 'instant' soups, noodles

**Table 7** Correlation across food systems domains

	Food & trade policies	Food supply	Food sources	Affordability	Convenience	Desirability	Power	Point of consumption	Dietary behaviours & dietary intake	Health outcomes
<b>Food &amp; trade policies</b>	1.000									
<b>Food supply</b>	0.676	1.000								
<b>Food sources</b>	0.180	0.521	1.000							
<b>Affordability</b>	0.036	0.273	0.445	1.000						
<b>Convenience</b>	0.274	0.702	0.521	0.273	1.000					
<b>Desirability</b>	0.086	0.355	0.164	0.175	0.355	1.000				
<b>Power</b>	-0.109	0.175	0.081	0.169	0.175	0.492	1.000			
<b>Point of consumption</b>	-0.145	0.029	0.350	0.373	0.029	0.097	0.230	1.000		
<b>Dietary behaviours &amp; dietary intake</b>	-0.457	-0.274	0.166	0.261	-0.274	-0.393	0.109	0.473	1.000	
<b>Health outcomes</b>	0.036	-0.167	-0.309	0.026	-0.387	-0.497	-0.079	0.014	0.261	1.000

Higher values depict strong correlation. Positive correlation (greater than 0) shows that both variables move in the same direction, while a negative correlation signifies that two variables move in opposite direction. Source: Authors' analyses of reviewed articles

**Table 8** Summary information on full-text articles reviewed

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Sadik	2010	2008 & 2009	Tamale	Not specified	2–18 years	40 children & 23 orphanage workers	Primary quantitative cross-sectional study	Demographic questionnaire, quantitative food frequency questionnaire, and anthropometric measurement modules	Descriptive statistics	white sugar, custard, sugar, white bread, brown bread, creamer, cold drink fanta.
Abiba et al.	2012	Not specified	Tamale Metropolis	Urban	> = 9 years	100 pupils upper primary (class 4–6) from six basic schools	Primary mixed methods cross-sectional study	2-week recall food frequency questionnaire, and anthropometric measurements module. Informal interviews with school vendors inside & outside school, and pupils. Observation methods used to record level of sanitation and food hygiene by vendors, cooks, and pupils.	Descriptive statistics	sweetened beverages, ice cream and biscuits

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Intiful and Lartey	2014	Not specified	Upper and Lower Manya-Krobo Districts of the Eastern Region of Ghana	Rural	6–10 years	359 children from 10 public schools	Primary quantitative cross-sectional study	Structured questionnaire to collect information on dietary habits, and a 24-h recall questionnaire to collect data on types, quantities and frequency of foods consumed		cocoa, fruit juice, coke/fanta, pie/chips, biscuits, condensed milk, margarine
Buxton	2014	Not specified	Cape Coast Metropolitan, Central Region of Ghana.	Not specified	13 years and over	820 adolescents in six schools (3 private and 3 public)	Primary qualitative study	Structure questionnaire to capture demographics and household characteristics, and a module on dietary practises	Simple descriptive and bivariate analysis	pastries such as cakes, meat pie, sausage roll, doughnut; ice cream (Yoghurt/fan or icefan chocolate); soft drinks (Fanta, coke, sprite); canned/packaged fruit juice
Fernandes et al.	2017	2013 & 2014	60 districts from all 10 regions of Ghana	Urban and peri-urban	5–17 years	4,258 children in 111 schools from 1,951 households located in 111 communities	cross-sectional survey applying mixed methods approach.	Household and school questionnaire administered to head teachers/caterers and parents/caregivers. Interview guide for focus group discussions/caregivers/parents	multi-level models with random effects at the household, school, and district levels for quantitative analysis. Basic content analysis for qualitative data	Confectionery, fruit, and sugar-sweetened beverages

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Bragg et al.	2017	Not specified	Accra, and areas within the Accra to Cape Coast highway	Urban area for the 4.7 km square. Not specified in the case of the 151 km highway	Not specified	151 km of highway (Accra to Cape Coast), and 4.7 km square area in Accra	Primary descriptive qualitative study	14-item codebook to capture food marketing themes in outdoor advertising	content analysis, simple descriptive statistics	Sugar-sweetened beverages (cola, fanta), bottled or canned juice, milk-based processed drinks (milo) energy drinks, water-based beverages
Ross et al.	2017	July–August 2008	Buduburam refugee settlement and Awutu, Accra	Urban	16–19 years	480 women aged 16–48 years	Primary quantitative cross-sectional survey,	A 6-month recall food frequency questionnaire to capture food consumption	Simple descriptive analysis, propensity score matching to group foods into either healthy, fats, sweets, and others	Candy, Sweet baked goods, biscuits/cookies, doughnuts, Sweetened fruit drinks (homemade/imported), soft drinks, Ready-to-eat cereal, corn flakes, processed meat, Cheese, Fried/salty snacks, Butter/margarine, sugar, tomato/tomato-based products (tin tomato), powdered or canned/tinned milk, salad dressing (mayonnaise)



Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Yang et al.	2017	2012	National study	All areas	12–15 years	1326	Quantitative study based on secondary data from Global School-Based Health Survey (GSHS) administered by the WHO and US CDC	Structured self-administered questionnaire. 30-day recall on consumption of carbonated drinks	Simple descriptive analysis	carbonated soft drinks
Aryeetey et al.	2017	December 2009–February 2012	Accra and Kumasi	Urban	9–15 years	3089 children from 121 schools	Primary quantitative cross-sectional survey	Structure questionnaire to capture demographics and household characteristics, a one week recall food frequency questionnaire, a module on physical activity, and anthropometric module	Multiple logistic regressions were used to analyse associations. Factor analysis used to derive household socio-economic status	Soft drinks, sweetened drinks
Ogum-Alangea et al.	2018	Not specified	Ga-East Municipality of the Greater Accra Region of Ghana	Urban	9–15 years	487 children from 24 schools (12 public and 12 private)	Primary quantitative cross-sectional study	Structured questionnaire to collect demographics, and a 7-day recall food frequency questionnaire used to record consumption of 100 food items. Anthropometric module to collect weight and height measurements.	Principal component analysis & linear regressions	processed meats, spreads and toppings, cocoa beverages, sugar sweetened beverages, fruit juices

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Ganle et al.	2019	Not specified	Tema Metropolitan district, 30 km East of Accra	Urban	5–16 years	285 children from two schools (1 public and 1 private)	Primary quantitative cross-sectional study	Questionnaire plus anthropometric measurements	Descriptive, bivariate and multivariable logistic regression analysis. WHO Growth standards used to convert anthropometric measurements to Z scores	junk foods, fizzy drinks
Adom et al.	2019	Not specified	Adentan Municipality of Greater Accra Region	Urban	8–11 years	543 children, attending 14 private and public primary schools	Primary quantitative cross-sectional study	Structured questionnaire to collect data on demographics and household characteristics self-reported recall questionnaire for dietary data, International Physical Activity Questionnaire for data on physical activity, and anthropometric measurements	Multivariable logistic regressions	sweets, chocolates, sweetened beverages, and soft drinks

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Adom et al.	2019	Not specified	Adentan Municipality of Greater Accra Region	Urban	8–11 years	543 children, attending 14 primary schools (8 private and 6 public)	Primary quantitative cross-sectional study	Structured questionnaires for children to self-report demographic and household characteristics, and interview administered questionnaire for school headteachers/administrators. Anthropometric module for height and weight measurements	mixed effects univariable and multivariable logistic and linear regressions models	chocolates, biscuits, cakes, cookies, sweets/toffees, packaged fruit juices sodas/soft drinks, ice creams, sausages rolls, doughnuts, pies; regular chips & crackers, and popcorn
Abizari and Ali	2019	May 2017	Tamale metropolis	Urban	10–19 years	366 pupils from 10 junior high schools	Primary quantitative cross-sectional study	Semi-structure questionnaire for socio-demographics, a 24-h recall for dietary diversity and a 7-day recall food frequency questionnaire where type and frequency of consumption of food item was consumed. Also included was an anthropometric module to collect height and weight measurements	Bivariate and multivariate logistic regressions, and principal component analysis to generate wealth index and derive dietary patterns	local beverages with added sugar; sweets such as toffees and chewing gums, fruit juices; energy and soft drinks

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Oyekale	2019	2014	National study	All areas	15–19 years	9367 women aged 15–49 years, 1346 adolescents (15–19-year-olds)	Quantitative study based on secondary data from Demographic and Health Survey	DHS survey questionnaires	Probit regression using instrumental variables	broth cubes, processed can meats, salted dried meat
Pradeilles et al.	2019	February–May 2018	Ga Mashie in Accra and Ho Central in Ho	Urban	Adolescents, although exact age group not specified	36 key informants purposefully selected to represent a wide range of sectors (schools, commerce, health, religious institutions, development agencies, traditional authorities, and youth clubs)	Primary descriptive qualitative study	Interviews with key informants, with a Consumer Readiness Module tool	Content analysis to generate consumer readiness score	unhealthy food and beverages; processed meats; sugar and sweet spreads; cakes and sweets; sodas and sweetened drinks; oils; spreading fats; cooking fats; and condiments (such as ketchup, mayonnaise)
Ogum-Alangea et al.	2020	Not specified	Ga East Municipality of Ghana	Urban and peri-urban	9–17 years	644 children in five schools (3 public and 2 private)	Primary quantitative cross-sectional study	Structured questionnaire to record children's food purchases	Simple descriptive analysis. Ghana food composition database used to weigh and analyse nutrition content of food items purchased	pastries, fried foods, and confectioneries, fizzy or sweetened drinks, fried foods, pastries, packaged snacks, fried sausages, milk & cocoa drinks, fried sausages, soybean khebab

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Darling et al.	2020	Not specified	Ningo Pram-pram district within the greater Accra region in coastal southern Ghana	Rural	10–19-year-olds	625 adolescents	Primary quantitative cross-sectional study	Structured questionnaire to collect socio-demographic data, 24-h recall for consumption of 14 food groups, 30-day recall for frequency of consumption of carbonated drinks, 7-day recall for consumption of fast foods, 7-day call for physical activity, and anthropometric module to record height and weight measurements	Simple descriptive analyses and Log-Poisson regression models	carbonated soft drinks, sweets, fast food

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Stevano et al.	2020	2015–2017	Accra Metropolitan Area	Urban	Not specified	139 schoolchildren enrolled in four Junior High Schools (2 public and 2 private), sub-sample of 63 children across four schools took part in focus group discussions	Primary mixed cross-sectional study	Student survey on food consumption and nutrition knowledge, 24-h recall dietary diversity module, focus groups with sub-groups of children, semi-structured interviews with key stakeholders (food industry, international organisations, school and street food vendors, and ministries).	Principal component analysis to derive a child asset score as measure of socio-economic status. Linear regressions to estimate associations	Fanmilk products (yoghurt-based, chocolate-based or have a fruit flavour)
Holdsworth et al.	2020	June–December 2017	Accra and Ho	Urban	13–19 years	adolescents/adults aged ≥ 13 y (male and female) in deprived neighbourhoods in Ghana and Kenya. Sample size is n = 294 in Ghana (192 in Accra and 96 in Ho).	Primary quantitative cross-sectional survey,	24-h recall questionnaire to capture dietary intake on 229 food items in Ghana	Descriptive statistics & Negative binomial regression models	fried sausage, Sweets/toffee, chocolate, bofrot, sugar, sweet pie/fart,, sugar/sweets, Sweetened tea/coffee, burkina drink, sobolo, sodas and sweetened beverages, Bread, meat pie, fried sausage, sugar, sweet spreads, biscuits, sweets and toffee, doughnuts, tapioca, vegetable oil, margarine



Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Stevano et al.	2020	2015–2017	Accra Metropolitan Area	Urban	Not specified	139 schoolchildren enrolled in four Junior High Schools (2 public and 2 private), sub-sample of 63 children across four schools took part in focus group discussions	Primary mixed methods cross-sectional study	Student survey on food consumption and nutrition knowledge, 24-h recall dietary diversity module, focus groups with sub-groups of children, semi-structured interviews with key stakeholders (food industry, international organisations, school and street food vendors, and ministries).	Principal component analysis to derive a child asset score as measure of socio-economic status	Fanmilk products (yoghurt-based, chocolate-based or have a fruit flavour), nestle's milo, indomie noodles
Smith et al.	2021	2007 & 2012	National study	Not specified but assumption is that all areas are covered	13–15 years	1648 adolescents	Quantitative study based on secondary data from Global School-Based Health Survey (GSHS) administered by the WHO and US CDC	7-day recall and self-reported questionnaire	pooled (all countries) multivariable logistic regression analysis	fast food and carbonated soft drinks

Table 8 (continued)

Authors	Publication year	Data collection year	Location	Setting	Age group	Realised sample size	Study design	Data collection tool	Data analysis methods	Processed foods reported
Sirikyi et al.	2021	August–September 2018	Cape Coast, Central Region of Ghana	Urban	16–19 years	196 students at the University of Cape Coast, aged 20 years and below	Primary cross-sectional study design	A self-administered questionnaire to capture demographics, dietary practices, substance abuse and physical activities	Multiple logistics regression	ice cream intake, fast food intake, sweetened soft drink
Pradeilles et al.	2021	May 2017–June 2018	Accra and Ho	Urban	13–19 years	98 total. Approximately 33% aged 13–18 years	Primary qualitative, study	Photovoice methodology	Coding & thematic analysis	fast food, indomie noodles
Appiah et al.	2021	October–November 2019	Ledzokuku-Krowor, Greater Accra	Urban	10–19 years	423 pregnant adolescents	Primary cross-sectional quantitative study	Food habits checklist & 24-h recall food frequency questionnaire	Descriptive and inferential statistics (chi-square & logistic regressions)	carbonated soft drinks, energy drinks, pastries

Source: Authors' analyses of reviewed articles

**Acknowledgements** We thank Prof. Gyorgy Scrinis for comments that greatly improved the manuscript and Ms Liz Millar for administrative assistance in navigating the funding process.

**Authors' contributions** Fiorella Picchioni (FP), Sara Stevano (SS), Emmanuel A. Codjoe (EC), Paul Kwame Nkegbe (PN) and Christopher Turner (CT) developed the initial concept. FP secured funding for the research project. Winnie Chepng'etich Sambu (WS) & FP developed the research design, with inputs from SS, EC, PN, and CT. WS conducted the literature search and reviewed all research articles included in the study. WS and FP analysed the data and drafted the paper. SS, EC, PN and CT reviewed the draft paper and provided critical feedback.

**Funding** This work was supported by the University of Greenwich Innovation Fund seedling fund (round 2020/21).

**Data availability** The data in this paper derives from secondary material available to all.

## Declarations

**Ethics approval** This study has used secondary sources and data, and no research participants were included. The study is a review of existing research articles that are publicly available. For this reason, it did not require ethical approval.

**Consent to participate and/or consent to publish** This study has used secondary sources and data, and no research participants were included. The study is a review of existing research articles that are publicly available. For this reason, it did not require consent to participate.

**Competing interests** The authors have no relevant financial or non-financial interests to disclose.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Abiba, A., Grace, A., & Kubreziga, C. (2012). Dietary patterns on the nutritional status of upper primary school children in Tamale metropolis. *Pakistan Journal of Nutrition*, 11(7), 689–707. <https://doi.org/10.3923/pjn.2012.689.707>
- Abizari, A. R., & Ali, Z. (2019). Dietary patterns and associated factors of schooling Ghanaian adolescents. *Journal of Health, Population, and Nutrition*, 38(1), 5–5. <https://doi.org/10.1186/s41043-019-0162-8>
- Adom, T., De Villiers, A., Puoane, T., & Kengne, A. P. (2019). Prevalence and correlates of overweight and obesity among school children in an urban district in Ghana. *BMC Obesity*, 6(1), 14. <https://doi.org/10.1186/s40608-019-0234-8>
- Adom, T., Kengne, A. P., De Villiers, A., & Puoane, T. (2019). Association between school-level attributes and weight status of Ghanaian primary school children. *BMC Public Health*, 19(1), 1–10. <https://doi.org/10.1186/s12889-019-6937-4>
- Akwuah, P. K., & Kobia-Acquah, E. (2020). Childhood obesity and overweight in Ghana: A systematic review and meta-analysis. *Journal of Nutrition and Metabolism*, 2020, 1907416. <https://doi.org/10.1155/2020/1907416>
- Alderman, H., Hoddinott, J., & Kinsey, B. (2006). Long term consequences of early childhood malnutrition. *Oxford Economic Papers*, 58(3), 450–474. <https://doi.org/10.1093/oeq/gpl008>
- Amoateng, A. Y., Doegah, P. T., & Udomboso, C. (2017). Socio-demographic factors associated with dietary behaviour among young Ghanaians aged 15–34 years. *Journal of Biosocial Science*, 49(2), 187–205. <https://doi.org/10.1017/S0021932016000456>
- Andam, K. S., Tschirley, D., Asante, S. B., Al-Hassan, R. M., & Diao, X. (2018). The transformation of urban food systems in Ghana: Findings from inventories of processed products. *Outlook on Agriculture*. <https://doi.org/10.1177/0030727018785918>
- Appiah, P. K., Naa Korklu, A. R., Bonchel, D. A., Fenu, G. A., & Wadga-Mieza Yankey, F. (2021). Nutritional knowledge and dietary intake habits among pregnant adolescents attending antenatal care clinics in urban community in Ghana. *Journal of Nutrition and Metabolism*, 2021. <https://doi.org/10.1155/2021/8835704>
- Aryeetey, R., & Tay, M. (2015). Compliance audit of processed complementary foods in urban Ghana. *Frontiers in Public Health*, 3. <https://doi.org/10.3389/fpubh.2015.00243>
- Aryeetey, R., Lartey, A., Marquis, G. S., Nti, H., Colecraft, E., & Brown, P. (2017). Prevalence and predictors of overweight and obesity among school-aged children in urban Ghana. *BMC Obesity*, 4(1), 1–8. <https://doi.org/10.1186/s40608-017-0174-0>
- Asosega, K. A., Adebajji, A. O., & Abdul, I. W. (2021). Spatial analysis of the prevalence of obesity and overweight among women in Ghana. *BMJ Open*, 11(1). <https://doi.org/10.1136/bmjopen-2020-041659>
- Aurino, E., Fernandes, M., & Penny, M. E. (2017). The nutrition transition and adolescents' diets in low- and middle-income countries: A cross-cohort comparison. *Public Health Nutrition*, 20(1), 72–81. <https://doi.org/10.1017/S1368980016001865>
- Ayensu, J., Annan, R., Lutterodt, H., Edusei, A., & Peng, L. S. (2020). Prevalence of anaemia and low intake of dietary nutrients in pregnant women living in rural and urban areas in the Ashanti region of Ghana. *PLoS ONE*, 15(1). <https://doi.org/10.1371/journal.pone.0226026>
- Baker, P., Lacy-Nichols, J., Williams, O., & Labonté, R. (2021). The political economy of healthy and sustainable food systems: An introduction to a special issue. *International Journal of Health Policy and Management*, 10(10 Special Issue on Political Economy of Food Systems), 734–744. <https://doi.org/10.34172/IJHPM.2021.156>
- Béné, C. (2022). Why the great food transformation may not happen – A deep-dive into our food systems' political economy, controversies and politics of evidence. *World Development*, 154. <https://doi.org/10.1016/j.worlddev.2022.105881>
- Benkeser, R. M., Biritwum, R., & Hill, A. G. (2012). Prevalence of overweight and obesity and perception of healthy and desirable body size in urban, Ghanaian Women. *Ghana Medical Journal*, 46(2), 66–75.
- Bimpong, K. A., Cheyuo, E. K. E., Abdul-Mumin, A., Ayanore, M. A., Kubuga, C. K., & Mogre, V. (2020). Mothers' knowledge and attitudes regarding child feeding recommendations, complementary feeding practices and determinants of adequate diet. *BMC Nutrition*, 6(1). <https://doi.org/10.1186/s40795-020-00393-0>
- Blakemore, S.-J. (2019). Adolescence and mental health. *The Lancet*, 393(10185), 2030–2031. [https://doi.org/10.1016/S0140-6736\(19\)31013-X](https://doi.org/10.1016/S0140-6736(19)31013-X)

- Bliznashka, L., Danaei, G., Fink, G., Flax, V. L., Thakwalakwa, C., & Jaacks, L. M. (2021). Cross-country comparison of dietary patterns and overweight and obesity among adult women in urban Sub-Saharan Africa. *Public Health Nutrition*, 24(6), 1393–1403. <https://doi.org/10.1017/S1368980019005202>
- Bragg, M. A., Hardoby, T., Pandit, N. G., Raji, Y. R., & Ogedegbe, G. (2017). A content analysis of outdoor non-alcoholic beverage advertisements in Ghana. *British Medical Journal Open*, 7(5), 1–6. <https://doi.org/10.1136/bmjopen-2016-012313>
- Brouwer, I. D., McDermott, J., & Ruben, R. (2020). Food systems everywhere: Improving relevance in practice. *Global Food Security*, 26, 100398. <https://doi.org/10.1016/j.gfs.2020.100398>
- Buxton, A. C. N. (2014). Ghanaian junior high school adolescents dietary practices and food preferences: Implications for public health concern. *Journal of Nutrition & Food Sciences*, 4(5). <https://doi.org/10.4172/2155-9600.1000297>
- Caulfield, L. E., Richard, S. A., Rivera, J. A., Musgrove, P., & Black, R. E. (2006). Stunting, wasting, and micronutrient deficiency disorders. In D. T. Jamison, J. G. Breman, A. R. Measham, G. Alleyne, M. Claeson, D. B. Evans, P. Jha, A. Mills, & P. Musgrove (Eds.), *Disease control priorities in developing countries* (2nd ed.). World Bank.
- Clapp, J. (2014). Financialization, distance and global food politics. *The Journal of Peasant Studies*, 41(5), 797–814. <https://doi.org/10.1080/03066150.2013.875536>
- Clapp, J., Moseley, W. G., Burlingame, B., & Termine, P. (2022). Viewpoint: The case for a six-dimensional food security framework. *Food Policy*, 106, 102164. <https://doi.org/10.1016/j.foodpol.2021.102164>
- Crouch, S. H., Soepnel, L. M., Kolkenbeck-Ruh, A., Maposa, I., Naidoo, S., Davies, J., Norris, S. A., & Ware, L. J. (2022). Paediatric hypertension in Africa: A systematic review and meta-analysis. *EClinicalMedicine*, 43, 101229–101229. <https://doi.org/10.1016/j.eclinm.2021.101229>
- Darkwa, S. (2014). Knowledge of nutrition facts on food labels and their impact on food choices on consumers in Koforidua, Ghana: A case study. *South African Journal of Clinical Nutrition*, 27(1), 13–17. <https://doi.org/10.1080/16070658.2014.11734479>
- Darling, A. M., Sunguya, B., Ismail, A., Manu, A., Canavan, C., Assefa, N., Sie, A., Fawzi, W., Sudfeld, C., & Guwattude, D. (2020). Gender differences in nutritional status, diet and physical activity among adolescents in eight countries in sub-Saharan Africa. *Tropical Medicine and International Health*, 25(1), 33–43. <https://doi.org/10.1111/tmi.13330>
- Development Initiatives. (2021). *2021 global nutrition report*.
- Fanzo, J., Covic, N., Dobermann, A., Henson, S., Herrero, M., Pingali, P., & Staal, S. (2020). A research vision for food systems in the 2020s: Defying the status quo. *Global Food Security*, 26, 100397. <https://doi.org/10.1016/j.gfs.2020.100397>
- FAO. (2020). *Support to the planting for food and jobs campaign*.
- Fernandes, M., Folsom, G., Aurino, E., & Gelli, A. (2017). A free lunch or a walk back home? The school food environment and dietary behaviours among children and adolescents in Ghana. *Food Security*, 9(5), 1073–1090. <https://doi.org/10.1007/s12571-017-0712-0>
- Ganle, J. K., Boakye, P. P., & Baatiema, L. (2019). Childhood obesity in urban Ghana: Evidence from a cross-sectional survey of in-school children aged 5–16 years. *BMC Public Health*, 19(1), 1–12. <https://doi.org/10.1186/s12889-019-7898-3>
- Ghana Statistical Service, Ghana Health Service, & ICF International. (2015). *Ghana demographic and health survey 2014* (No. 233302682661/2). Retrieved May 5, 2023, from [www.DHSprogram.com](http://www.DHSprogram.com)
- Global Panel on Agriculture and Food Systems for Nutrition. (2014). *How can agriculture and food system policies improve nutrition*.
- Gyasi, R. M., Peprah, P., & Appiah, D. O. (2020). Association of food insecurity with psychological disorders: Results of a population-based study among older people in Ghana. *Journal of Affective Disorders*, 270, 75–82. <https://doi.org/10.1016/j.jad.2020.03.088>
- Hargreaves, D., Mates, E., Menon, P., Alderman, H., Devakumar, D., Fawzi, W., Greenfield, G., Hammoudeh, W., He, S., Lahiri, A., Liu, Z., Nguyen, P. H., Sethi, V., Wang, H., Neufeld, L. M., & Patton, G. C. (2022). Strategies and interventions for healthy adolescent growth, nutrition, and development. *The Lancet*, 399(10320), 198–210. [https://doi.org/10.1016/S0140-6736\(21\)01593-2](https://doi.org/10.1016/S0140-6736(21)01593-2)
- Hawkes, C. (2006). Uneven dietary development: Linking the policies and processes of globalization with the nutrition transition, obesity and diet-related chronic diseases. *Globalization and Health*, 2(1), 4–4. <https://doi.org/10.1186/1744-8603-2-4>
- Hawkes, C., Friel, S., Lobstein, T., & Lang, T. (2012). Linking agricultural policies with obesity and noncommunicable diseases: A new perspective for a globalising world. *Food Policy*, 37(3), 343–353. <https://doi.org/10.1016/j.foodpol.2012.02.011>
- Herforth, A., & Ahmed, S. (2015). The food environment, its effects on dietary consumption, and potential for measurement within agriculture-nutrition interventions. *Food Security*, 7(3), 505–520. <https://doi.org/10.1007/s12571-015-0455-8>
- HLPE. (2017). *Nutrition and food systems. A report by the high level panel of experts on food security and nutrition of the committee on world food security*. Retrieved August 21, 2023, from [www.fao.org/cfs/cfs-hlpe](http://www.fao.org/cfs/cfs-hlpe)
- Holdsworth, M., Pradeilles, R., Tandoh, A., Green, M., Wanjohi, M., Zotor, F., Asiki, G., Klomegah, S., Abdul-Haq, Z., Osei-Kwasi, H., Akparibo, R., Bricas, N., Auma, C., Griffiths, P., & Laar, A. (2020). Unhealthy eating practices of city-dwelling Africans in deprived neighbourhoods: Evidence for policy action from Ghana and Kenya. *Global Food Security*, 26, 100452–100452. <https://doi.org/10.1016/j.gfs.2020.100452>
- International Food Policy Research Institute (IFPRI), I. F. P. (2019). *Food system innovations for healthier diets in low and middle-income countries*. <https://doi.org/10.2499/p15738coll2.133156>
- International Food Policy Research Institute (IFPRI), I. F. P. (2020). *2020 global food policy report: building inclusive food systems*. <https://doi.org/10.2499/9780896293670>
- Intiful, F. D., & Lartey, A. (2014). Breakfast habits among school children in selected communities in the eastern region of Ghana. *Ghana Medical Journal*, 48(2), 71–77. <https://doi.org/10.4314/gmj.v48i2.3>
- Jonah, C. M. P., Sambu, W. C., & May, J. D. (2018). A comparative analysis of socioeconomic inequities in stunting: A case of three middle-income African countries. *Archives of Public Health*, 76(1). <https://doi.org/10.1186/s13690-018-0320-2>
- Keats, E. C., Rappaport, A. I., Shah, S., Oh, C., Jain, R., & Bhutta, Z. A. (2018). The dietary intake and practices of adolescent girls in low-and middle-income countries: A systematic review. *Nutrients*, 10(12). <https://doi.org/10.3390/nu10121978>
- Koplan, J. P., Liverman, C. T., & Kraak, V. (2005). *Preventing childhood obesity: Health in the balance* (p. 11015). Institute of Medicine of The National Academies. <https://doi.org/10.17226/11015>
- Laar, A., Barnes, A., Aryeetey, R., Tandoh, A., Bash, K., Mensah, K., Zotor, F., Vandevijvere, S., & Holdsworth, M. (2020). Implementation of healthy food environment policies to prevent nutrition-related non-communicable diseases in Ghana: National experts' assessment of government action. *Food Policy*, 93. <https://doi.org/10.1016/j.foodpol.2020.101907>
- Lawrence, M. A., Friel, S., Wingrove, K., James, S. W., & Candy, S. (2015). Formulating policy activities to promote healthy and sustainable diets. *Public Health Nutrition*, 18(13), 2333–2340. <https://doi.org/10.1017/S1368980015002529>
- Linderhof, V., Vlijm, R., Pinto, V., Raaijmakers, I., & Dijkshoorn-Dekker, M. (2019). *Urban food security in Ghana: A policy review*. <https://doi.org/10.18174/476849>

- Manyanga, T., El-Sayed, H., Doku, D. T., & Randall, J. R. (2014). The prevalence of underweight, overweight, obesity and associated risk factors among school-going adolescents in seven African countries. *BMC Public Health*, *14*(1). <https://doi.org/10.1186/1471-2458-14-887>
- Masters, W. A., Bai, Y., Herforth, A., Sarpong, D. B., Mishili, F., Kinabo, J., & Coates, J. C. (2018). Measuring the affordability of nutritious diets in Africa: Price indexes for diet diversity and the cost of nutrient adequacy. *American Journal of Agricultural Economics*, *100*(5), 1285–1301. <https://doi.org/10.1093/ajae/aay059>
- Masters, W. A., Rosenblum, N. Z., & Alemu, R. G. (2018). Agricultural transformation, nutrition transition and food policy in Africa: Preston curves reveal new stylised facts. *Journal of Development Studies*, *54*(5), 788–802. <https://doi.org/10.1080/00220388.2018.1430768>
- Mayes, C. (2014). Medicalization of Eating and Feeding. In *Encyclopedia of Food and Agricultural Ethics* (pp. 1–8). Springer Netherlands. [https://doi.org/10.1007/978-94-007-6167-4\\_49-2](https://doi.org/10.1007/978-94-007-6167-4_49-2)
- Mbogori, T., Kimmel, K., Zhang, M., Kandiah, J., & Wang, Y. (2020). Nutrition transition and double burden of malnutrition in Africa: A case study of four selected countries with different social economic development. *AIMS Public Health*, *7*(3), 425–439. <https://doi.org/10.3934/publichealth.2020035>
- Meng, T., Florowski, W. J., Sarpong, D. B., Chinnan, M. S., & Resurreccion, A. V. A. (2014). Consumer's food shopping choice in Ghana: Supermarket or traditional outlets? *International Food and Agribusiness Management Review*, *17*(1030–2016–82985), 107–130. <https://doi.org/10.22004/ag.econ.164600>
- Ministry of Food and Agriculture. (2007). *Food and Agriculture Sector Development Policy (FASDEP II)*.
- Ministry of Health. (2012). *National policy for the prevention and control of chronic non-communicable diseases in Ghana*.
- Ministry of Health. (2022). *National Policy Non-Communicable Diseases*.
- Minot, N., & Dewina, R. (2015). Are we overestimating the negative impact of higher food prices? Evidence from Ghana. *Agricultural Economics*, *46*(4), 579–593. <https://doi.org/10.1111/agec.12183>
- Monteiro, C. A., Cannon, G., Levy, R., Moubarac, J.-C., Jaime, P., Martins, A. P., Canella, D., Louzada, M., & Parra, D. (2016). *Food Classification. Public Health*, *7*(1), 11.
- Monteiro, C. A., Moubarac, J. C., Cannon, G., Ng, S. W., & Popkin, B. (2013). Ultra-processed products are becoming dominant in the global food system. *Obesity Reviews*, *14*(S2), 21–28. <https://doi.org/10.1111/obr.12107>
- Muthuri, S. K., Francis, C. E., Wachira, L. J. M., LeBlanc, A. G., Sampson, M., Onywera, V. O., & Tremblay, M. S. (2014). Evidence of an overweight/obesity transition among school-aged children and youth in Sub-Saharan Africa: A systematic review. *PLoS ONE*, *9*(3). <https://doi.org/10.1371/journal.pone.0092846>
- Nelson, F., Nyarko, K., & Binka, F. (2015). Prevalence of risk factors for non-communicable diseases for new patients reporting to Korle-Bu Teaching Hospital. *Ghana Medical Journal*, *49*(1), 12. <https://doi.org/10.4314/gmj.v49i1.3>
- Nestle, M. (2013). *Food politics: How the food industry influences nutrition and health* (3rd ed.). University of California Press.
- Neufeld, L. M., Andrade, E. B., Ballonoff Suleiman, A., Barker, M., Beal, T., Blum, L. S., Demmler, K. M., Dogra, S., Hardy-Johnson, P., Lahiri, A., Larson, N., Roberto, C. A., Rodríguez-Ramírez, S., Sethi, V., Shamah-Levy, T., Strömmer, S., Tumilowicz, A., Weller, S., & Zou, Z. (2022). Food choice in transition: Adolescent autonomy, agency, and the food environment. *The Lancet*, *399*(10320), 185–197. [https://doi.org/10.1016/S0140-6736\(21\)01687-1](https://doi.org/10.1016/S0140-6736(21)01687-1)
- Nordhagen, S., Lambertini, E., DeWaal, C. S., McClafferty, B., & Neufeld, L. M. (2022). Integrating nutrition and food safety in food systems policy and programming. *Global Food Security*, *32*, 100593. <https://doi.org/10.1016/j.gfs.2021.100593>
- Norris, S. A., Frongillo, E. A., Black, M. M., Dong, Y., Fall, C., Lampl, M., Liese, A. D., Naguib, M., Prentice, A., Rochat, T., Stephensen, C. B., Tinago, C. B., Ward, K. A., Wrottesley, S. V., & Patton, G. C. (2022). Nutrition in adolescent growth and development. *The Lancet*, *399*(10320), 172–184. [https://doi.org/10.1016/S0140-6736\(21\)01590-7](https://doi.org/10.1016/S0140-6736(21)01590-7)
- Obirikorang, Y., Obirikorang, C., Enoch, O. A., Acheampong, E., Tboseiyefah, P., Miriam, A. A., & Emmanuella, N. B. (2017). Prevalence and risk factors of obesity among practicing nurses at three selected hospitals in Kumasi Metropolis, Ghana. *Journal of Medical and Biomedical Sciences*, *5*(3), 45–55. <https://doi.org/10.4314/jmbs.v5i3.5>
- Ofori-Asenso, R., Agyeman, A. A., Laar, A., & Boateng, D. (2016). Overweight and obesity epidemic in Ghana—A systematic review and meta-analysis. *BMC Public Health*, *16*(1). <https://doi.org/10.1186/s12889-016-3901-4>
- Ogum Alangea, D., Aryeetey, R. N., Gray, H. L., Laar, A. K., & Adanu, R. M. K. (2018). Dietary patterns and associated risk factors among school age children in urban Ghana. *BMC Nutrition*, *4*(1), 1–9. <https://doi.org/10.1186/s40795-018-0230-2>
- Ogum-Alangea, D., Aryeetey, R. N. O., Gray, H. L., Laar, A. K., & Adanu, R. M. K. (2020). Basic school pupils' food purchases during mid-morning break in urban Ghanaian schools. *PLoS ONE*, *15*(9), 1–13. <https://doi.org/10.1371/journal.pone.0238308>
- Owiredo, E. W., Dontoh, E., Essuman, S. E. S., & Bazanfara, B. B. (2019). Demographic and lifestyle predictors of prehypertension: A cross-sectional study among apparently healthy adults in Kumasi, Ghana. *BioMed Research International*, *2019*, 1–7. <https://doi.org/10.1155/2019/1764079>
- Owureku-Asare, M., Ambrose, R. P. K., Oduro, I., Tortoe, C., & Saalia, F. K. (2017). Consumer knowledge, preference, and perceived quality of dried tomato products in Ghana. *Food Science & Nutrition*, *5*(3), 617–624. <https://doi.org/10.1002/fsn3.439>
- Oyekale, A. S. (2019). Effect of obesity and other risk factors on hypertension among women of reproductive age in Ghana: An instrumental variable probit model. *International Journal of Environmental Research and Public Health*, *16*(23). <https://doi.org/10.3390/ijerph16234699>
- Park, M. H., Falconer, C., Viner, R. M., & Kinra, S. (2012). The impact of childhood obesity on morbidity and mortality in adulthood: A systematic review. *Obesity Reviews*, *13*(11), 985–1000. <https://doi.org/10.1111/j.1467-789X.2012.01015.x>
- Popkin, B. M. (2014). Nutrition, agriculture and the global food system in low and middle income countries. *Food Policy*, *47*, 91–96. <https://doi.org/10.1016/j.foodpol.2014.05.001>
- Pradeilles, R., Irache, A., Wanjohi, M. N., Holdsworth, M., Laar, A., Zotor, F., Tandoh, A., Klomegah, S., Graham, F., Muthuri, S. K., Kimani-Murage, E. W., Coleman, N., Green, M. A., Osei-Kwasi, H. A., Bohr, M., Rousham, E. K., Asiki, G., Akparibo, R., Mensah, K., & Griffiths, P. (2021). Urban physical food environments drive dietary behaviours in Ghana and Kenya: A photovoice study. *Health and Place*, *71*. <https://doi.org/10.1016/j.healthplace.2021.102647>
- Pradeilles, R., Marr, C., Laar, A., Holdsworth, M., Zotor, F., Tandoh, A., Klomegah, S., Coleman, N., Bash, K., Green, M., & Griffiths, P. L. (2019). How ready are communities to implement actions to improve diets of adolescent girls and women in urban Ghana? *BMC Public Health*, *19*(1), 1–14. <https://doi.org/10.1186/s12889-019-6989-5>
- Prentice, A. M., Ward, K. A., Goldberg, G. R., Jarjou, L. M., Moore, S. E., Fulford, A. J., & Prentice, A. (2013). Critical windows for nutritional interventions against stunting. *The American Journal of Clinical Nutrition*, *97*(5), 911–918. <https://doi.org/10.3945/ajcn.112.052332>
- Raza, A., Fox, E. L., Morris, S. S., Kupka, R., Timmer, A., Dalmiya, N., & Fanzo, J. (2020). Conceptual framework of food systems



- for children and adolescents. *Global Food Security*, 27, 100436. <https://doi.org/10.1016/j.gfs.2020.100436>
- Reardon, T., Tschirley, D., Liverpool-Tasie, L. S. O., Awokuse, T., Fanzo, J., Minten, B., Vos, R., Dolislager, M., Sauer, C., Dhar, R., Vargas, C., Lartey, A., Raza, A., & Popkin, B. M. (2021). The processed food revolution in African food systems and the double burden of malnutrition. *Global Food Security*, 28. <https://doi.org/10.1016/j.gfs.2020.100466>
- Ross, W. L., Gallego-Pérez, D. F., Lartey, A., Sandow, A., Pérez-Escamilla, R., & Hromi-Fiedler, A. (2017). Dietary patterns in Liberian refugees in Buduburam, Ghana. *Maternal and Child Nutrition*, 13(4), 1–10. <https://doi.org/10.1111/mcn.12401>
- Sadik, A. (2010). Orphanage children in Ghana- are their dietary needs met. *Pakistan Journal of Nutrition*, 9(9), 844–852.
- Sarfo, F. S., Mobula, L. M., Burnham, G., Ansong, D., Plange-Rhule, J., Sarfo-Kantanka, O., & Ofori-Adjei, D. (2018). Factors associated with uncontrolled blood pressure among Ghanaians: Evidence from a multicenter hospital-based study. *PLoS ONE*, 13(3), e0193494. <https://doi.org/10.1371/journal.pone.0193494>
- Sathyamala, C. (2016). Nutritionalizing food: A framework for capital accumulation. *Development and Change*, 47(4), 818–839. <https://doi.org/10.1111/dech.12250>
- Scrinis, G. (2016). Reformulation, fortification and functionalization: Big Food corporations' nutritional engineering and marketing strategies. *The Journal of Peasant Studies*, 43(1), 17–37. <https://doi.org/10.1080/03066150.2015.1101455>
- Simmonds, M., Llewellyn, A., Owen, C. G., & Woolacott, N. (2016). Predicting adult obesity from childhood obesity: A systematic review and meta-analysis. *Obesity Reviews*, 17(2), 95–107. <https://doi.org/10.1111/obr.12334>
- Sirikyi, I. H., Eliason, S., Ghartey, F. N., Ekenam, E., Pereko, K. K. A., Okai, E., Yiridong, F., Hagan, O. C. K., & Nsiah, P. (2021). Anthropometric indices and cardiometabolic risk factors in a Ghanaian adolescent population. *Journal of Pediatric Endocrinology and Metabolism*, 34(1), 35–44. <https://doi.org/10.1515/jpem-2020-0273>
- Smith, L., Jacob, L., Shin, J. I., Tully, M. A., Pizzol, D., López-Sánchez, G. F., Gorely, T., Yang, L., Grabovac, I., & Koyanagi, A. (2021). Bullying victimization and obesogenic behaviour among adolescents aged 12 to 15 years from 54 low- and middle-income countries. *Pediatric Obesity*, 16(1), 1–8. <https://doi.org/10.1111/jjpo.12700>
- Sparling, T. M., White, H., Boakye, S., John, D., & Kadiyala, S. (2021). Understanding pathways between agriculture, food systems, and nutrition: An evidence and gap map of research tools, metrics, and methods in the last 10 Years. *Advances in Nutrition*, 12(4), 1122–1136. <https://doi.org/10.1093/advances/nmaa158>
- Stevano, S., Johnston, D., & Codjoe, E. (2020a). Better decisions for food security? Critical reflections on the economics of food choice and decision-making in development economics. *Cambridge Journal of Economics*, 44(4), 813–833. <https://doi.org/10.1093/cje/beaa012>
- Stevano, S., Johnston, D., & Codjoe, E. (2020b). The urban food question in the context of inequality and dietary change: A study of schoolchildren in Accra. *Journal of Development Studies*, 56(6), 1177–1189. <https://doi.org/10.1080/00220388.2019.1632434>
- Tankwanchi, A. B. S. (2018). Oppression, liberation, wellbeing, and ecology: Organizing metaphors for understanding health workforce migration and other social determinants of health. *Globalization and Health*, 14(1). <https://doi.org/10.1186/s12992-018-0397-y>
- Tricco, A. C., Lillie, E., Zarin, W., O'Brien, K. K., Colquhoun, H., Levac, D., Moher, D., Peters, M. D. J., Horsley, T., Weeks, L., Hempel, S., Akl, E. A., Chang, C., McGowan, J., Stewart, L., Hartling, L., Aldcroft, A., Wilson, M. G., Garritty, C., & Straus, S. E. (2018). PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and explanation. *Annals of Internal Medicine*, 169(7), 467–473. <https://doi.org/10.7326/M18-0850>
- Turner, C., Aggarwal, A., Walls, H., Herforth, A., Drewnowski, A., Coates, J., Kalamatianou, S., & Kadiyala, S. (2018). Concepts and critical perspectives for food environment research: A global framework with implications for action in low- and middle-income countries. *Global Food Security*, 18, 93–101. <https://doi.org/10.1016/j.gfs.2018.08.003>
- Walls, H., Nisbett, N., Laar, A., Drimie, S., Zaidi, S., & Harris, J. (2021). Addressing malnutrition: The importance of political economy analysis of power. *International Journal of Health Policy and Management*, 10(12), 809–816. <https://doi.org/10.34172/ijhpm.2020.250>
- Winson, A. (2013). *The industrial diet: The degradation of food and the struggle for healthy eating* (p. 174). UBC Press.
- World Health Organization. (2022). *Deaths from noncommunicable diseases on the rise in Africa*.
- Yang, L., Bovet, P., Liu, Y., Zhao, M., Ma, C., Liang, Y., & Xi, B. (2017). Consumption of carbonated soft drinks among young adolescents aged 12 to 15 years in 53 low- and middle-income countries. *American Journal of Public Health*, 107(7), 1095–1100. <https://doi.org/10.2105/AJPH.2017.303762>
- Ziraba, A. K., Fotso, J. C., & Ochako, R. (2009). Overweight and obesity in urban Africa: A problem of the rich or the poor? *BMC Public Health*, 9(1), 465. <https://doi.org/10.1186/1471-2458-9-465>



**Winnie Chepng'etich Sambu** is a PhD student at the University of Cape Town's School of Economics where she is studying trends and dynamics of food demand in South Africa. She has nearly 10 years' experience in applied research, advocacy, and policy engagements in the areas of food & nutrition Security, poverty & inequality, transportation and mobility, early Childhood development, youth unemployment, and living conditions.



**Fiorella Picchioni** is a Senior Research Fellow in Political Economy of Food Systems at the Natural Resources Institute, University of Greenwich. She is an applied development economist and her work is positioned at the intersection of food systems thinking, gender analysis, nutrition and agrarian transitions.

Her recent research include applying a political economy lens to the analysis of the impacts of ultra-processed food, and food fortification interventions on wellbeing in the Global South.





**Sara Stevano** is a development and feminist political economist. She is a Senior Lecturer in Economics at SOAS University of London, after holding teaching and research positions at the University of the West of England, Bristol, and King's College London. Her areas of study are the political economy of work, food and nutrition, inequalities and social reproduction. Her work focuses on Africa, with primary research experience in Mozambique and Ghana. Sara is committed to expanding the boundaries of economic research and teaching through interdisciplinary approaches, qualitative methods and micro-macro bridges.

to expanding the boundaries of economic research and teaching through interdisciplinary approaches, qualitative methods and micro-macro bridges.



**Emmanuel A. Codjoe** Emmanuel A. Codjoe received his PhD from SOAS, University of London and is currently a senior lecturer in Economics at the University of Ghana, Department of Economics. His teaching focuses on economic theory, international economics, and development economics at both the undergraduate and postgraduate level. His recent research has

focused on COVID-19 and the garment industry in Ghana, food choice and the food environment.



**Paul Kwame Nkegbe** holds a PhD in Agricultural and Food Economics from the University of Reading, UK, a degree he obtained in 2011. He is currently an Associate Professor of Agricultural and Applied Economics at the Department of Development Studies, University for Development Studies, Ghana. His research interests are in applied microeconomic analysis and his recent work has concentrated on rural infrastructure and markets, and how they impact household food security and nutrition.

and how they impact household food security and nutrition.



**Christopher Turner** is an interdisciplinary food systems researcher with a background in social science, human geography and public health nutrition. His interests include the development and application of theoretical concepts and novel methodologies for food systems and food environment

research. His mixed-methods research features participatory approaches, such as qualitative-GIS and photovoice, to investigate the lived experience of food environments and understand drivers of food acquisition practices with a view to co-designing public health policies for sustainable healthy diets – particularly in low- and middle-income country settings.