

Ecosystem services from coffee agroforestry in Central America: Estimation using the CAF2021 model

Marcel van Oijen^{1*} Jeremy Haggard² Mirna Barrios³ Lucie Büchi²
Rolando Cerda⁴ Stefania Cerretelli² Erick López⁵
Elias de Melo Virginio Filho⁴ Alejandra Ospina⁴

2022-01-17

¹ Independent researcher, Edinburgh, United Kingdom

² Natural Resources Institute, University of Greenwich, Chatham Maritime, Medway, United Kingdom

³ CATIE, Centro Agronómico Tropical de Investigación y Enseñanza, Managua, Nicaragua

⁴ CATIE, Centro Agronómico Tropical de Investigación y Enseñanza, Turrialba 30501, Costa Rica

⁵ Centro Estudios Ambientales y Biodiversidad, Universidad del Valle de Guatemala, Ciudad de Guatemala, Guatemala

Abstract: The goal of sustainable coffee production requires multiple functions from agroforestry systems. Many are difficult to quantify and data are lacking, hampering the choice of shade tree species and agronomic management. Process-based modelling may help quantify ecosystem services and disservices. We introduce and apply coffee agroforestry model CAF2021 (<https://doi.org/10.5281/zenodo.5725276>). The model allows for complex systems with up to three shade tree species. It simulates coffee yield, timber and fruit production by shade trees, soil loss in erosion, C-sequestration, N-fixation, -emission and -leaching. To calibrate the model, we used multivariate data from 32 different treatments applied in two long-term coffee agroforestry experiments in Costa Rica and Nicaragua. Without any further calibration, the model was then applied to agroforestry systems on 89 farms in Costa Rica and 79 in Guatemala where yields had been reported previously in farmer interviews. Despite wide variation in environmental and agronomic conditions, the model explained 36% of yield variation in Costa Rica but only 15% in Guatemala. Model analysis quantified trade-offs between yield and other ecosystem services as a function of fertilisation and shading.

Key words: Agroforestry systems, *Coffea arabica* L., Ecosystem services, Fertilisation, Process-based modelling, Shading

* Corresponding author: ORCID 0000-0003-4028-3626, VanOijenMarcel@gmail.com

Declarations

Acknowledgments: We acknowledge support by the Institute for Coffee in Costa Rica (ICAFFE), the National Coffee Association of Guatemala (ANACAFE), Manos Campesinas in Guatemala, and the coffee farmers in both countries who enabled this research.

Funding: This research was funded by UK Research and Innovation Biotechnology and Biological Sciences Research Council (UKRI/BBSRC), from the Global Challenges Research Fund (GCRF) under the Agri-systems Research to Enhance Rural Livelihoods in Developing Countries programme, Grant No. BB/S01490X/1.

Conflicts of interest/Competing interests: None.

Ethics approval: Not applicable.

Consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and material: Requests for data should be addressed to JH.

Code availability: The coffee agroforestry model CAF2021 can be downloaded from <https://doi.org/10.5281/zenodo.5725276>.

1 Introduction

The goal of sustainable coffee production requires multiple functions from agroforestry systems. Coffee productivity and other services should be high while disservices such as negative environmental impacts are limited. Here we use the collective term *ecosystem services* for all these processes, noting that some must be maximized and others minimized. Trade-offs between ecosystem services in agroforestry systems are complex. Shade trees compete for resources, and may thus reduce coffee yields, but they also provide income (Rice 2008) and, by improving the microclimate, stabilise yields in the face of environmental change as discussed in a recent review by van Noordwijk et al. (2021).

An important issue in the evaluation of coffee agroforestry (CAF) systems is that many ecosystem services are difficult to quantify, which hampers the choice of shade tree species and agronomic management. This is a particular problem in Central America, where mountainous topography with young volcanic soils leads to strong spatial variation in weather patterns and soil fertility. Together with a wide variation in management choices, this has led to strong variability in productivity (Hagggar et al. 2021). It is likely also to cause strong variation in environmental ecosystem services, but not many data are available to estimate these.




Figure 1: Locations of sampled coffee farms in Costa Rica and Guatemala.

Here we used a combination of data analysis and process-based modelling to quantify ecosystem services from coffee agroforestry on 89 farms in Costa Rica and 79 in Guatemala (Fig. 1). The same farms were studied previously by Hagggar et al. (2021) who found that coffee productivity has no simple relationship with shade. Low-productivity farms tended to have either low or high shade.

The measurements at these farms did not include ecosystem services related to carbon and nitrogen biogeochemistry. Here we estimated these ecosystem services using the new coffee agroforestry model CAF2021, which is freely available online (<https://doi.org/10.5281/zenodo.5725276>). The model differs from its predecessors CAF2007 (Van Oijen et al. 2010a) and CAF2014 (Rahn et al. 2018; Ovalle-Rivera et al. 2020) mainly in that it allows for more complex agroforestry systems with up to three different shade tree species. The model simulates a wide range of ecosystem services: coffee yield, timber and fruit production by shade trees, soil loss by erosion, soil carbon sequestration, nitrogen fixation, loss of nitrogen in atmospheric emissions and leaching. Other process-based models of coffee agroforestry have been developed in recent years (Charbonnier et al. 2017; Vezy et al. 2019). These models focus on plant physiology, and they simulate agronomy and biogeochemistry in less detail than CAF2021, which makes them less suitable to assess a wide range of ecosystem services.

Long-term field experiments that measure a broad suite of variables are required to parameterise and evaluate complex process-based vegetation models. To calibrate CAF2021, we used multivariate data from a total of 32 different treatments applied in two long-term coffee agroforestry experiments in Costa Rica and Nicaragua. A subset of these data was used earlier in the parameterisation of CAF2014 by Ovalle-Rivera et al. (2020). These authors only used data from 10 of the treatments because their model did not simulate agroforestry systems with more than one shade tree species. We used the same method of model parameterisation, i.e. Bayesian calibration, an increasingly common method for calibration and uncertainty quantification of process-based models (Van Oijen 2020).

The application of the model to the farms in Costa Rica and Guatemala constitutes a test of the predictive capacity of the model against independent data as farm information was not used in the calibration. The farm data were less comprehensive than the experimental data, so we also compare model output against the rich literature on the impact of shading and fertilisation on delivery of ecosystem services from coffee agroforestry systems in Central America (Soto-Pinto et al. 2000; Cannavo et al. 2011; Meylan et al. 2013, 2017; Jha et al. 2014; Goodall et al. 2015; Sauvadet et al. 2019; Hagggar et al. 2021; De Leijster et al. 2021).

The goals of this study were threefold:

- 1: Identifying the drivers of observed differences in productivity between farms and CAF types.
- 2: Assessing the implications of aiming for any particular CAF type for biogeochemistry-related ecosystem services and sustainability.
- 3: Assessing CAF2021 as a tool for studying coffee agroforestry systems.

2 Materials and methods

2.1 Data from experiments at Turrialba and Masatepe

For model calibration, we used data from two long-term coffee agroforestry experiments at Turrialba in Costa Rica and Masatepe in Nicaragua. The experiments differed in various respects: choice of coffee cultivar and shade tree species, shade management, fertiliser types and amounts, weather conditions and soil properties. In both experiments, four fertilisation regimes were combined with different choices of one or two shade tree species, selected from *Erythrina poeppigiana*, *Terminalia amazonia*, *Chloroleucon eurycyclum*, *Samanea saman*, *Tabebuia rosea*, *Simarouba glauca*, *Inga laurina*. Both experiments provided data on coffee, shade trees and soil, as reported by Hagggar et al. (2011), Noponen (2012), Noponen et al. (2013), Sepúlveda (2016) and Ovalle-Rivera et al. (2020). We refer to these publications and to the Supplementary Information (S2) for further details.

2.2 Data from farms in Costa Rica and Guatemala

We used data from 89 coffee agroforestry (CAF) farms in Costa Rica and 79 farms in Guatemala collected by the SEACAF project (Hagggar et al. 2021). In the present study, these data are used for independent testing of CAF2021 without any farm-specific model calibration. The farm locations are shown in Fig. 1.

Table 1: Average environmental conditions for farms grouped by CAF system type in Costa Rica and Guatemala. The first digit of CAF type indicates relative productivity, the second shading level, both from 1=low to 4=high. Variables: altitude (m), slope (degrees), rain (mm y⁻¹), temperature (degrees C), soil water holding capacity (-), soil carbon and nitrogen stocks (kg m⁻²). Within brackets: standard deviation across farms.

Country	CAF	n	Altitude	Slope	Rain	Temp	WaterCap	Csoil	Nsoil
Costa Rica	1.2	25	1128 (185)	13.3 (7.9)	2804 (440)	20.5 (0.9)	0.25 (0.03)	14.5 (4.9)	1.8 (0.5)
	2.4	26	1148 (257)	13.3 (5.5)	2562 (345)	20.5 (1.3)	0.24 (0.03)	13.6 (2.6)	1.8 (0.4)
	3.2	24	1354 (252)	14.8 (6.7)	2480 (372)	19.3 (1.5)	0.25 (0.03)	15.9 (4.5)	2 (0.5)
	4.3	14	1332 (217)	14.8 (7.5)	2401 (293)	19.5 (1.1)	0.24 (0.04)	14.6 (4.6)	1.8 (0.6)
Guatemala	1.3	19	1490 (344)	14.9 (6)	2934 (690)	19.7 (2.5)	0.24 (0.05)	10.4 (3.9)	1.2 (0.4)
	2.4	27	1531 (320)	15.3 (10.1)	2370 (773)	19 (2.1)	0.26 (0.04)	11.1 (3.9)	1.2 (0.5)
	3.3	25	1678 (231)	10.6 (8)	2048 (600)	18.2 (1.7)	0.28 (0.03)	13.4 (5.2)	1.4 (0.5)
	4.3	8	1576	11.8	2020	18.4	0.29	13.8	1.4

Country	CAF	n	Altitude	Slope	Rain	Temp	WaterCap	Csoil	Nsoil
			(292)	(5.3)	(821)	(2.1)	(0.03)	(6.7)	(0.7)

Table 1 summarises the environmental conditions at the on-farm CAFs grouped according to the typology of Haggard et al. (2021). The table includes average temperature and precipitation, but for running CAF2021 full time series of daily weather from 2005 to 2020 were derived from the WorldClim database (Fick and Hijmans 2017) with additional information for recent years from weather stations run by Icafe and Instituto Meteorológico Nacional in Costa Rica and by Anacafe in Guatemala. Soil water holding capacity was derived from soil texture information (bulk density, silt, clay) using a pedotransfer function tested for tropical soils (Tomasella et al. 2003; Tomasella and Hodnett 2004). Soil carbon and nitrogen stocks measured in the top 26 cm were extrapolated to 1 m depth assuming the top 26 cm contained 80% of the stocks.

Table 2: Management and coffee yield under four CAF types in Costa Rica and Guatemala. Variables: N-fertilisation rate ($\text{kg ha}^{-1} \text{y}^{-1}$), density of shade trees (ha^{-1} ; E=Erythrina, I=Inga, M=Musa, P=Persea, G=Grevillea, C=Cordia), overall shade level (-) and coffee yield 2018-2020 ($\text{kg DM ha}^{-1} \text{y}^{-1}$). Within brackets: standard deviation.

Country	CAF	N-fert	E	I	M	P	G	C	Shade	Yield
Costa Rica	1.2	146 (127)	102 (136)	70 (122)	109 (156)	17 (36)	23 (64)	29 (58)	0.4 (0.17)	1044 (498)
	2.4	169 (99)	141 (157)	112 (155)	258 (303)	16 (37)	13 (29)	83 (170)	0.73 (0.08)	1787 (994)
	3.2	283 (114)	136 (156)	28 (78)	230 (243)	23 (66)	10 (17)	3 (12)	0.39 (0.12)	3145 (493)
	4.3	323 (133)	155 (105)	17 (59)	381 (340)	0 (0)	0 (0)	8 (22)	0.56 (0.09)	4583 (567)
Guatemala	1.3	75 (112)	0 (0)	103 (126)	111 (244)	54 (81)	39 (62)	49 (74)	0.47 (0.12)	558 (437)
	2.4	73 (72)	0 (0)	57 (83)	93 (169)	51 (86)	133 (149)	35 (97)	0.77 (0.09)	1071 (660)
	3.3	168 (133)	0 (0)	50 (64)	51 (166)	79 (102)	103 (118)	48 (121)	0.49 (0.15)	2334 (775)
	4.3	174 (170)	0 (0)	21 (34)	94 (227)	40 (55)	136 (193)	16 (35)	0.42 (0.17)	5433 (947)

Table 2 with on-farm CAF system properties shows management practices for fertilisation and shading, and coffee yield. Some farms had reported implausibly high fertilisation rates which we capped at $500 \text{ kg N ha}^{-1} \text{y}^{-1}$. We allocated farm-specific tree density data to the six species-types listed in the table (see also Table 3). Tree pruning and thinning regimes were not reported but had to be specified for running CAF2021. Trees were assumed to be pruned once per year, targeting each farm's reported shade level, except for Erythrina which in Costa Rica tends to follow a prescribed twice-yearly 90% pruning regime. Tree thinning was only applied to Cordia following a prescribed

calendar. Pruned branches were removed from the field in Guatemala while remaining on the field in Costa Rica. Dry coffee fruit production was calculated as one third of reported fresh yield (Sepúlveda 2016).

2.3 CAF2021 structure

CAF2021 is a process-based model for the biogeochemistry of coffee agroforestry systems. We give a short description here, focusing on how the model differs from its predecessor models. The model and a user guide are freely available online (<https://doi.org/10.5281/zenodo.5725276>).

CAF2021 simulates a coffee agroforestry system where the coffee plants are shaded by up to 3 tree species. The model allows for two lower-layer tree species, but no more than one upper-layer tree species. In typical use of the model the upper layer would be occupied by timber trees and the lower layer by service and fruit trees. The simulated tree species can have different initial planting densities and thinning regimes. The model simulates the biogeochemistry - flows of carbon, nitrogen and water - of the soil-coffee-trees system. State variables are pools of carbon, nitrogen and water as well as coffee developmental stage. Height and crown area growth are simulated for each of the tree species. The model uses a daily time step and therefore requires daily weather data (light, temperature, rain, wind, humidity) as input. Further required inputs are atmospheric $[CO_2]$, soil properties (slope, water retention parameters, initial contents of C and N) and management calendars for fertilisation, pruning and thinning. The pruning and thinning calendars can be specified separately for each plant species.

The outputs from the model are time series with daily values of state variables and processes including ecosystem services such as productivity of coffee and trees, N-leaching, N-emission, erosion in the form of soil organic matter loss (C & N) in runoff, C-sequestration in the soil.

The model is implemented in FORTRAN but called from a user-interface written in R.

CAF2021 differs from its predecessor models in the following respects:

- There can be three tree species present rather than just one. All tree variables are implemented as dynamic arrays whose length equals the number of tree species. Users with experience in programming can therefore relatively easily extend the number of shade tree species beyond three.
- Fruit production can be simulated.
- Lower-layer trees do not overlap, so the number of ground-cover conditions (vertical vegetation profiles) in any simulated field is at most six: coffee shaded by (1) no trees, (2) service trees, (3) fruit trees, (4) timber trees, (5) service + timber trees, (6) fruit + timber trees. The model keeps track of the fractional ground area covered by each of these six categories (see Supplementary Information S1 for an example).
- The relative heights, leaf area indices (LAI) and light extinction coefficients of the two competing tree species within ground cover categories (5) and (6) determine their access to light.
- Morphology (tree height as a function of stem biomass and crown width as a function of branch biomass) is simulated using allometric equations but for each species a genetic or management-induced maximum height can be specified.
- Shade management can be fully specified using calendars for dates and intensities of pruning and thinning. Alternatively, it may be simulated as being goal-directed toward a specified shade level.

Time series for selected output variables from CAF2021 are shown in the Supplementary Information (S1 and S3).

2.4 Bayesian calibration of CAF2021

We calibrated the parameters of CAF2021 using a Bayesian approach to allow for uncertainty quantification. Data from the two long-term experiments in Turrialba and Masatepe that we described above were used for this purpose. These measurements were suitable for calibration of a complex model such as CAF2021 because they comprised a wide range of coffee, tree and soil variables. This reduced the risk of tuning the model to one variable such as coffee yield production at the cost of poor simulation of other system processes.

For the calibration of CAF2021, we used data from 18 Turrialba treatments and 14 Nicaragua treatments. Some of these data had been used for Bayesian calibration of the predecessor model CAF2014 by Ovalle-Rivera et al. (2020), but that model was limited to treatments with no more than one tree species, so only six of the treatments from Turrialba and four from Masatepe could be used by them. In other respects we followed their implementation of Bayesian calibration.

We carried out a single multi-site, multi-treatment Bayesian calibration of CAF2021, using Markov Chain Monte Carlo sampling (MCMC) with a chain length of 60,000. At each iteration of the chain the model was run for all 32 treatments, so a total of 1.92 million runs was carried out during the MCMC. The calibration provided samples from the posterior distribution for 65 of the model's parameters:

- *universal* parameters (n=45) that were not allowed to vary between the treatments,
- *site-specific* parameters (n=5) that were allowed to differ between Turrialba and Masatepe,
- *tree-species specific* parameters (n=15).

The universal parameters included all 23 calibrated coffee parameters, thus ignoring any differences between cultivars in Turrialba and Masatepe. Also treated as universal were 15 tree parameters and 7 soil parameters. The motivation for treating so many parameters as universal (which in reality may show variation) was to maintain a degree of universal applicability of the model, and to constrain the need for site- or tree-specific information in future applications of the model. The 5 site-specific parameters were soil parameters governing nitrogen leaching, organic matter composition and turnover. Making these parameters site-specific allowed simulation of the greater capacity of soils in Masatepe to stabilise organic matter and minerals (Noponen 2012). The 15 tree-species specific parameters were predominantly parameters for carbon allocation and morphology. All parameters were a priori assigned wide beta probability distributions, reflecting large prior uncertainty about plausible parameter values for the largely new model CAF2021.

Bayesian calibration generates a sample from the joint posterior probability distribution for all calibrated parameters. The MAP is the '*Maximum A Posteriori*' parameter vector, i.e. the model parameterisation that achieves the highest value for the product of prior (beta distributions for parameters) and likelihood (fit to data) (Ovalle-Rivera et al. 2020; Van Oijen 2020). Results reported in this paper are for the MAP parameter vector.

2.5 CAF2021 application to farms Costa Rica and Guatemala

We used CAF2021 to simulate measured and unmeasured ecosystem services for each of the 168 farms in Costa Rica and Guatemala, using each farm's environmental conditions and CAF

management choices as model inputs. All parameters for coffee and for Erythrina and Inga trees were left at the values determined by Bayesian calibration using data from the long-term experiments at Turrialba and Masatepe. Tree species not present in the experiments were grouped according to type and management (Table 2). Parameter values for the most abundant tree species in each group (listed as the ‘Type species’ in Table 3) were taken from the literature, specifically for banana (*Musa spp.*) (Schaffer et al. 1999; Chaves C. et al. 2009; Martinez Acosta and Cayón Salinas 2011; Mustaffa and Kumar 2012; Van den Bergh et al. 2012; Damour et al. 2012), avocado (*Persea americana*) (Elzebroek and Wind 2008; CIRAD 2019; Monzón-Martinez 2019), Grevillea (Castellanos et al. 2010) and Cordia (Castellanos et al. 2010; Van Oijen et al. 2010b).

Table 3: Shade tree species and their management under CAF2021.

Tree type	Type species	Stratum	N-fixing	Pruning/thinning regime	Fate of pruned/thinned biomass and fruits
Pollarded legume service tree	Erythrina poeppigiana	Lower	Yes	Pollarded: 90% branch removal twice per year	Remain on site
Pruned legume service tree	Inga spp.	Lower	Yes	Pruned: % dependent on target tree cover	Option to remove branches
Fruit tree	<i>Persea americana</i>	Lower	No	Not pruned: biomass time coefficient 1000 days	Fruit exported
Bananas and plantains	<i>Musa spp.</i>	Lower	No	Not pruned: biomass time coefficient 365 days	Fruit exported
Service/timber tree	<i>Grevillea robusta</i>	Upper	No	Pruned: % dependent on target tree cover	Option to remove branches
Timber tree	<i>Cordia alliodora</i>	Upper	No	Thinned: 30% 5-yearly	Thinned trunks removed

Planting dates were not recorded in the farms database but coffee stands were generally mature and we simulated the years 2005-2020 for each farm. The only differences between Costa Rica and Guatemala in our model set-up were conform local practice: (1) pruned branches were removed in Guatemala but left on the field in Costa Rica, (2) the height of some tree species was differently managed.

2.5.1 Analysis of sensitivity to management

Besides the regular simulations for each farm, which used the observed and reported local conditions, we made two additional model runs to test farm sensitivity to management choices. In the first of these, we halved each farm’s fertilisation rate. In the second, we halved tree densities and shade target.

3 Results

3.1 Bayesian calibration on data from experiments in Turrialba and Masatepe




Figure 2: Observed vs. simulated coffee dry matter yields. Left: calibration experiments in Turrialba ($n=18$) and Masatepe ($n=14$). Middle: farms in Costa Rica ($n=89$). Right: farms in Guatemala ($n=79$).

The left panel of Fig. 2 shows observed coffee productivity in the 32 treatments against the yields simulated by the calibrated model. The yield values are averages over the years 2002-2017 for Turrialba and 2004-2013 for Masatepe. The model accounts for 74% of observed variation in yield across the Turrialba treatments, and 63% of variation at Masatepe. Other measured variables were simulated with similar accuracy (see Supplementary Information).

Posterior parameter uncertainty was small, with an average coefficient of variation of 5%. However, each model output depends on multiple parameters, and output uncertainty for individual variables (coffee yield, soil carbon sequestration, nitrogen leaching) was 30-36%. These are predictive uncertainties for model application to a single set of conditions and do not apply to trends across treatments.

3.2 CAF2021 outputs for different coffee agroforestry systems in Costa Rica and Guatemala

CAF2021 accounted for 36% of observed variation in coffee yields for the on-farm CAF in Costa Rica (middle panel of Fig. 2), which is less than the variation accounted for in the Costa Rican long-term experiment at Turrialba (left panel), but compares well given the fact that the farm simulations were not calibrated. The results for the Guatemalan on-farm CAF were less good, with only 15% of yield variation accounted for (right panel).

Table 4: CAF types on farms in Costa Rica and Guatemala: Simulations of shade (-), yield of coffee, timber and fruit (kg DM ha⁻¹ y⁻¹), soil carbon change and carbon runoff (kg C ha⁻¹ y⁻¹), N-fixation, leaching and emission (kg N ha⁻¹ y⁻¹).

Country	CAF	Shade	Yield	Timber	Fruit	dC	Crun	Nfix	Nlch	Nem
Costa Rica	1.2	0.48 (0.22)	2615 (1903)	696 (1042)	472 (833)	224 (1434)	408 (289)	15 (19)	84 (44)	23 (9)
	2.4	0.72 (0.09)	2623 (2125)	1269 (1432)	512 (701)	1175 (1081)	300 (176)	17 (23)	69 (31)	22 (11)
	3.2	0.48 (0.23)	5184 (1627)	198 (424)	1040 (1110)	147 (1351)	279 (168)	9 (10)	103 (37)	35 (10)
	4.3	0.6 (0.18)	5555 (2170)	202 (685)	1255 (1018)	656 (1276)	222 (119)	13 (9)	110 (29)	38 (9)
Guatemala	1.3	0.46 (0.18)	938 (1676)	721 (633)	419 (1162)	-828 (1096)	448 (236)	9 (13)	79 (80)	14 (11)
	2.4	0.64 (0.14)	697 (930)	979 (579)	183 (286)	-832 (656)	335 (281)	5 (13)	51 (36)	15 (11)
	3.3	0.5 (0.2)	2963 (2863)	668 (466)	245 (725)	-891 (945)	192 (123)	2 (4)	69 (30)	26 (19)
	4.3	0.45 (0.27)	2785 (3304)	539 (426)	431 (1051)	-971 (1268)	207 (104)	2 (3)	67 (41)	37 (43)

Table 4 summarises the on-farm simulations for coffee yield and seven other ecosystem services grouped by the CAF system types of Hagggar et al. (2021). The model correctly identified CAF types 1.2/3 and 2.4 as much lower-yielding in both countries than CAF types 3.2/3 and 4.3 (comparing Tables 2 and 4). However, the model was unable to explain the very high yield difference that was observed between the two highest-yielding types in Guatemala, which were 2334 and 5433 kg ha⁻¹ y⁻¹ of bean dry matter for CAF type 3.3 and 4.3 respectively, whereas the simulations were 2963 and 2785 kg ha⁻¹ y⁻¹. The very similar yields in these simulations are consistent with these CAF types having similar environmental conditions (Table 1), shading and fertilisation (Table 2).

The CAF2021 simulations of Table 4 suggest that there were trade-offs between coffee yield and other ecosystem services. In Costa Rica, CAF types with high coffee yield also had high fruit yields but produced less timber and had higher levels of nitrogen leaching and emission. In Guatemala, the trade-offs were less marked apart from high N-emission rates in high-yielding CAF types.

To determine whether the trade-offs were driven by management choices we plotted the simulated ecosystem services at each farm against its level of N-fertilisation (Fig. 3) and shading (Fig. 4). The results suggest that ecosystem services other than timber production were more strongly determined by level of fertilisation than by level of shading. Fertilisation had the expected effects of increasing leaching and emission but it also led to less erosion as measured by soil carbon loss in runoff. Coffee yields were more related to fertilisation level in Costa Rica than in Guatemala. This reflects the wider range and twice as high average level of fertilisation in Costa Rica (Table 2).




Figure 3: Simulations of ecosystem services at farms in Costa Rica (black) and Guatemala (red). Units as in Table 4. All plots are against N-fertilisation level (kg N ha⁻¹ y⁻¹).




Figure 4: Simulations of ecosystem services at farms in Costa Rica (black) and Guatemala (red). Units as in Table 4. All plots are against shade level (-).

3.3 Sensitivity to management

The analyses of Figures 3 and 4 show generally fairly weak correlations of the driving variables with the ecosystem variables. This was partly because of confounding of the many environmental and management differences between the farms. To highlight the effects of fertilisation and shading, we therefore carried out a sensitivity analysis where for each farm all conditions were kept the same apart from halving either fertilisation or shading. The results of this analysis are shown in Fig. 5. This shows that halving fertilisation is expected to reduce coffee yields, carbon sequestration, N-leaching and emission, while increasing (in Costa Rica) erosion through runoff. Halving shading also increases runoff but otherwise has very different effects: it increases coffee yield and reduces ecosystem services closely associated with trees (production of timber and fruit, N-fixation).




Figure 5: Changes in ecosystem services when fertilisation or shading is halved. Units as in Table 4. All changes are normalised to the country average.

4 Discussion

4.1 Bayesian calibration on data from experiments

We carried out a single Bayesian calibration - for all treatments in Turrialba and Masatepe simultaneously - to derive parameter estimates. All coffee parameters were treated as being universally applicable, and common tree species were also assumed to have the same properties in both experiments. The only experiment- and thus country-specific parameters in the calibration were five soil properties. Variation within experiments was ignored altogether. In reality, there may have been genetic and environmental variation in and between the experiments. Disregarding such plot and block effects may have hampered model fit but ensured that the model was not overfit to noise. Moreover, the calibration was conducted simultaneously for multiple variables, so model outputs were not optimised for any single variable such as coffee yield. Despite all these restrictions, the calibrated model accounted for 74% of observed coffee yield variation between treatments at Turrialba and 63% at Masatepe. This compares favourably with linear regression of yield on fertilisation level in the experiments for which the r^2 was only 0.23. It shows that models for ecosystem service evaluation in coffee agroforestry systems need to integrate the effects of not only management but also weather and soil conditions.

The results suggest that CAF2021 can be a useful tool for analysing differences in system performance between different environmental conditions. However, this does not imply that the model is a good tool for forecasting the benefits of agroforestry at any individual site, as posterior output uncertainty (driven by posterior parameter uncertainty) was high at 30-36%.

4.2 Farm simulations

After the Bayesian calibration on the data from the two experiments, no further calibration of CAF2021 was carried out. The application to farms in Costa Rica and Guatemala thus constituted an independent test of the predictive capacity of the model. This test was restricted to coffee yield as data from other ecosystem services were not available. Coffee yields on Costa Rican farms were simulated reasonably well ($r^2 = 0.36$), but yields on Guatemalan farms were not ($r^2 = 0.15$).

One possible explanation for the mixed performance of the model is the use of parameter values calibrated on data from just two experiments. The experiments may have been more representative of plant properties and environmental conditions in Costa Rica than in Guatemala. It would have been possible to calibrate the model on a subset of reported yields but the model would then be tuned as a coffee yield prediction tool to the possible detriment of predictive capacity for other ecosystem services. It is therefore of importance to find rich local data sets, with measurements of a variety of coffee, tree and soil variables, before adding a second Bayesian calibration.

A second possible explanation for uneven model performance is the greater variation in environmental conditions in Guatemala compared to Costa Rica, as measured by the larger standard deviations for temperature, precipitation and altitude (Table 1). In the simulations, all parameters for coffee and tree morphology and physiology were kept at the same values for all 168 farms (89 in Costa Rica and 79 in Guatemala). Any genetic differences between coffee- and tree-varieties were thus ignored. The only tree-parameter differences accounted for were the farm-specific initial densities of shade trees. Disregarding inter-farm variability in this way may have affected simulations in Guatemala the most because of its greater variation in growing conditions.

A third possible explanation for poor predictive capacity of the model is errors in data. In particular surprising was the observation that coffee yields in Guatemala were on average 2.3 times

higher on farm with CAF-type 4.3 than on those with type 3.3, despite similar environmental conditions and levels of fertilisation. The farms with CAF-type 4.3 had low densities of Inga trees (Table 2), so it is unlikely that nitrogen fixation explained their yield advantage. The soils of these farms also did not have higher nitrogen contents (Table 1) and their mineralisation rates were lower at $12.2 \text{ mg N kg}^{-1} \text{ soil d}^{-1}$ compared to 17.7 in CAF type 3.3 (Büchi et al., in prep.). The observed superior yield of CAF type 4.3 could not be reproduced by the model without reparameterisation. However, it must be noted that there were only 8 farms with this CAF type who reported high yields, and they also reported high variance in N application rates from over $1000 \text{ kg N ha}^{-1} \text{ y}^{-1}$ to almost zero. As stated in Hagggar et al. (2021), there were a small but significant group of farms that report high yields but use alternative low N input management and which require further study.

4.3 Analysis of ecosystem services

Except for CAF type 4.3, the observed and simulated coffee yields were higher in Costa Rica than in Guatemala, which matches the about twice as high average fertilisation rates in Costa Rica (Tables 2 and 3). The simulations suggest that the higher fertilisation in Costa Rica further stimulated carbon sequestration, nitrogen leaching and emission (Fig. 4). A positive effect of fertilisation on carbon sequestration was also shown by Körschens (2021), in their review of long-term experiments worldwide and in several empirical studies (Virto et al. 2012; Triberti et al. 2016). Other differences between the countries, such as production of more fruit but less timber in Costa Rica, were the result of different choices of shade tree species and planting density rather than differential impacts of fertilisation on tree species (Tables 2 and 3).

The model points to inevitable trade-offs between productivity of coffee, productivity of trees and environmental ecosystem services, similar to those reported by Meylan et al. (2017) for coffee agroforestry systems in Costa Rica. Changing the level of fertilisation or shading can benefit some ecosystem services but always to the detriment of others. This was apparent from the farm simulations shown in Figures 3 and 4, but was highlighted most clearly in the sensitivity analysis presented in Fig. 5. The benefits from fertilisation for yield and carbon sequestration must be balanced against the costs associated with increased nitrogen leaching and emission. Likewise the benefits for coffee yield of removing shading (consistent with literature summarised by Meylan et al. (2017) but not confirmed by Soto-Pinto et al. (2000) for an experiment in southern Mexico) must be balanced against the associated losses in tree production (fruit, timber) and carbon sequestration as well as the reduced protection against erosion. Analysis of the actual yield from the CAF systems here modelled concluded that farms with moderate shade levels were the most productive, and there was no yield advantage to further reduction in shade, but yields were suppressed by higher shade levels over 60% cover (Hagggar et al. 2021). In the simulations, the effect of shade-reduction on coffee yield varied between farms and was on average much smaller in Costa Rica than in Guatemala. De Leijster et al. (2021) summarised the literature on this and reported similarly inconsistent relationships between coffee yields and shade cover because of differences in local conditions and management. Vaast et al. (2008) concluded that moderate shading may not hamper coffee production in hot areas with suboptimal growing conditions.

Halving shading significantly reduced carbon stocks in the simulations (Fig. 5), and this is consistent with the finding of Goodall et al. (2015) that ongoing reductions in shade tree density on farms in Nicaragua are leading to loss of soil carbon stocks. Meylan et al. (2013) and Jha et al. (2014), in their reviews of impacts of shade in coffee agroforestry, also concluded that shading increases above- and belowground organic carbon sequestration. In some contrast to this, only a

minor impact of shading on carbon sequestration was reported for the Turrialba experiment by Sauvadet et al. (2019). Protection against erosion by high canopy cover, as simulated here, has repeatedly been observed in coffee agroforestry systems (Meylan et al. 2017; De Leijster et al. 2021 and references therein). Cannavo et al. (2011) found that runoff in unshaded coffee in Costa Rica was over 50% higher than runoff in coffee shaded by *Inga densiflora*. Our sensitivity analysis found a similar effect where halving shading led to 10-15% increase in carbon lost in runoff (Fig. 5).

The focus of CAF2021 is on biogeochemistry, and the effects of fertilisation on processes in soils, coffee plants and trees are simulated in considerable detail. However, the effects of shading on the coffee agroforestry system are manifold and complex, and are not simulated fully by CAF2021. The model does account for shade trees lowering light intensity and temperature below them (impacting coffee plant physiology) as well as competing with the coffee plants for light, water, and nitrogen. But the model does not represent other organisms and thus does not simulate how shade supports biodiversity (e.g. Goodall et al. 2015) and constrains pests and diseases (e.g. Jha et al. 2014). Decision support for such non-biogeochemical ecosystem services would therefore require a different model.

5 Conclusions

Returning to the three goals of this paper, we conclude:

- Variation in fertilisation and shading drives differences in productivity between farms and between CAF types.
- The modelled ecosystem services indicate trade-offs between ecosystem services that imply no single CAF type can maximize all services. The trade-offs were not only between provisioning and regulatory services, but also between provisioning services from coffee vs. those from fruit and timber trees.
- CAF2021 can be used as a tool for studying biogeochemistry-related ecosystem services in complex coffee agroforestry systems, but the availability of multivariate calibration data from long-term field experiments remains critical.

Supplementary information

S1. Output variables simulated by CAF2021

In standard use, CAF2021 simulates 169 output variables to quantify the dynamics of coffee plants, trees and soil. Fig. 6 shows time series for a small selection.




Figure 6: Selected output variables simulated by a single run of CAF2021 for Guatemala farm 79. Top row: field average soil content of C, N, water. Rows 2-3: area of coffee shaded by (1) no trees, (2) service trees, (3) fruit trees, (4) timber trees, (5) service + timber trees, (6) fruit + timber trees. Row 4-5: crown area and height of service, fruit and timber trees. Row 6: overall shade and coffee yield.

S2. Experimental treatments at Turrialba and Masatepe

CAF2021 was calibrated on data from 32 treatments applied in long-term experiments at Turrialba (Costa Rica) and Masatepe (Nicaragua). The following table shows the combinations of shading and fertilisation that were applied. For more details on the experiments, see Hagggar et al. (2011), Nojonen (2012), Nojonen et al. (2013), Sepúlveda (2016) and Ovalle-Rivera et al. (2020).

Table 5: Long-term experiments in Turrialba (treatments 1:18) and Masatepe (treatments 19:32). Shade trees: E = *Erythrina poeppigiana*, T = *Terminalia amazonia*, C = *Chloroleucon eurycyclum*, SS = *Samanea saman*, TR = *Tabebuia rosea*, SG = *Simarouba glauca*, IL = *Inga laurina*. Fertilisation regime: C = conventional, O = organic, I = intensive, M = moderate.

TURRIALBA	Shade	Fertil.	MASATEPE	Shade	Fertil.
1	E	CI	19	-	CI
2	E	CM	20	-	CM
3	E	OI	21	SSTR	CM
4	E	OM	22	SSTR	OI
5	T	CI	23	SGTR	CI
6	T	CM	24	SGTR	CM
7	T	OI	25	SGTR	OI
8	T	OM	26	SGTR	OM
9	C	CM	27	ILSS	CI
10	C	OI	28	ILSS	CM
11	TE	CM	29	ILSS	OI
12	TE	OI	30	ILSS	OM
13	CE	CI	31	ILSG	CM
14	CE	CM	32	ILSG	OI
15	CE	OI			
16	CE	OM			
17	-	CI			
18	-	CM			

S3. Data and output variables simulated in calibration of CAF2021

CAF2021 was calibrated on data from 32 treatments applied in long-term experiments at Turrialba (Costa Rica) and Masatepe (Nicaragua). Fig. 7 shows the data from treatment 1 at Turrialba together with output from the calibrated model.




Figure 7: Calibration results for treatment 1 of the Turrialba experiment. Blue points with error bars: data. Black lines: CAF2021 simulation using the MAP parameter vector. Green lines: simulation using the maximum likelihood parameter vector. For this treatment of highly fertilised coffee shaded by *Erythrina poeppigiana* trees, seven variables were measured. From top-left and row by row: coffee yield, shade, tree crown area and height, aboveground carbon in coffee and trees, soil carbon gain.

References

- Cannavo P, Sansoulet J, Harmand J-M, et al (2011) Agroforestry associating coffee and *Inga densiflora* results in complementarity for water uptake and decreases deep drainage in Costa Rica. *Agriculture, Ecosystems & Environment* 140:1–13. <https://doi.org/10.1016/j.agee.2010.11.005>
- Castellanos E, Quilo A, Pons D (2010) Estudio de linea base de carbono en cafetales. Universidad del Valle de Guatemala
- Charbonnier F, Roupsard O, le Maire G, et al (2017) Increased light-use efficiency sustains net primary productivity of shaded coffee plants in agroforestry system. *Plant, Cell & Environment* 40:1592–1608. <https://doi.org/10.1111/pce.12964>
- Chaves C. B, Cayón S. G, Jones JW (2009) Modeling plantain (*Musa AAB Simmonds*) potential yield. *Agronomía Colombiana* 27:359–366
- CIRAD (2019) Country Profile: mexico. HAB
- Damour G, Ozier-Lafontaine H, Dorel M (2012) Simulation of the growth of banana (*Musa* spp.) Cultivated on cover-crop with simplified indicators of soil water and nitrogen availability and integrated plant traits. *Field Crops Research* 130:99–108. <https://doi.org/10.1016/j.fcr.2012.02.013>
- De Leijster V, Santos MJ, Wassen MW, et al (2021) Ecosystem services trajectories in coffee agroforestry in Colombia over 40 years. *Ecosystem Services* 48:101246. <https://doi.org/10.1016/j.ecoser.2021.101246>
- Elzebroek T, Wind K (2008) *Guide to Cultivated Plants*. CABI, Wallingford, U.K.
- Fick SE, Hijmans RJ (2017) WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* 37:4302–4315. <https://doi.org/10.1002/joc.5086>
- Goodall KE, Bacon CM, Mendez VE (2015) Shade tree diversity, carbon sequestration, and epiphyte presence in coffee agroecosystems: A decade of smallholder management in San Ramón, Nicaragua. *Agriculture, Ecosystems & Environment* 199:200–206. <https://doi.org/10.1016/j.agee.2014.09.002>
- Haggar J, Barrios M, Bolaños M, et al (2011) Coffee agroecosystem performance under full sun, shade, conventional and organic management regimes in Central America. *Agroforestry Systems* 82:285–301. <https://doi.org/10.1007/s10457-011-9392-5>
- Haggar J, Casanoves F, Cerda R, et al (2021) Shade and Agronomic Intensification in Coffee Agroforestry Systems: Trade-Off or Synergy? *Frontiers in Sustainable Food Systems* 5:131. <https://doi.org/10.3389/fsufs.2021.645958>
- Jha S, Bacon CM, Philpott SM, et al (2014) Shade Coffee: Update on a Disappearing Refuge for Biodiversity. *BioScience* 64:416–428. <https://doi.org/10.1093/biosci/biu038>
- Körschens M (2021) Long-Term Field Experiments (LTEs) Importance, Overview, Soil Organic Matter. In: Mueller L, Sychev VG, Dronin NM, Eulenstein F (eds) *Exploring and Optimizing Agricultural Landscapes*. Springer International Publishing, Cham, pp 215–231
- Martinez Acosta AM, Cayón Salinas DG (2011) Dinámica del Crecimiento y Desarrollo del Banano (*Musa AAA Simmonds* cvs. Gran Enano y Valery). *RevFacNalAgrMedellin* 64:6055–6064

- Meylan L, Gary C, Allinne C, et al (2017) Evaluating the effect of shade trees on provision of ecosystem services in intensively managed coffee plantations. *Agriculture, Ecosystems & Environment* 245:32–42. <https://doi.org/10.1016/j.agee.2017.05.005>
- Meylan L, Merot A, Gary C, Rapidel B (2013) Combining a typology and a conceptual model of cropping system to explore the diversity of relationships between ecosystem services: The case of erosion control in coffee-based agroforestry systems in Costa Rica. *Agricultural Systems* 118:52–64. <https://doi.org/10.1016/j.agsy.2013.02.002>
- Monzón-Martínez ZL (2019) Perfil económico del aguacate de Guatemala y su potencial de exportación para los mercados de El Salvador y Honduras. PhD thesis, Universidad de San Carlos de Guatemala
- Mustaffa MM, Kumar V (2012) Banana production and productivity enhancement through spatial, water and nutrient management. *J Hortl Sci* 7:1–28
- Noponen MRA (2012) Carbon and economic performance of coffee agroforestry systems in Costa Rica and Nicaragua. PhD thesis
- Noponen MRA, Healey JR, Soto G, Haggard JP (2013) Sink or source? The potential of coffee agroforestry systems to sequester atmospheric CO₂ into soil organic carbon. *Agriculture, Ecosystems & Environment* 175:60–68. <https://doi.org/10.1016/j.agee.2013.04.012>
- Ovalle-Rivera O, Van Oijen M, Läderach P, et al (2020) Assessing the accuracy and robustness of a process-based model for coffee agroforestry systems in Central America. *Agroforestry Systems* 94:2033–2051. <https://doi.org/10.1007/s10457-020-00521-6>
- Rahn E, Vaast P, Läderach P, et al (2018) Exploring adaptation strategies of coffee production to climate change using a process-based model. *Ecological Modelling* 371:76–89. <https://doi.org/10.1016/j.ecolmodel.2018.01.009>
- Rice RA (2008) Agricultural intensification within agroforestry: The case of coffee and wood products. *Agriculture, Ecosystems & Environment* 128:212–218. <https://doi.org/10.1016/j.agee.2008.06.007>
- Sauvadet M, den Meersche KV, Allinne C, et al (2019) Shade trees have higher impact on soil nutrient availability and food web in organic than conventional coffee agroforestry. *Science of The Total Environment* 649:1065–1074. <https://doi.org/10.1016/j.scitotenv.2018.08.291>
- Schaffer B, Whiley AW, Searle C (1999) Atmospheric CO₂ Enrichment, Root Restriction, Photosynthesis, and Dry-matter Partitioning in Subtropical and Tropical Fruit Crops. *HortScience* 34:1033–1037. <https://doi.org/10.21273/HORTSCI.34.6.1033>
- Sepúlveda N et al. (2016) Efecto de la sombra y niveles de insumo sobre la producción de café en la región del pacífico de Nicaragua (2002-2015)
- Soto-Pinto L, Perfecto I, Castillo-Hernández J, Caballero-Nieto J (2000) Shade effect on coffee production at the northern Tzeltal zone of the state of Chiapas, Mexico. *Agriculture, Ecosystems & Environment* 80:61–69. [https://doi.org/10.1016/S0167-8809\(00\)00134-1](https://doi.org/10.1016/S0167-8809(00)00134-1)
- Tomasella J, Hodnett M (2004) Pedotransfer functions for tropical soils. In: *Developments in Soil Science*. Elsevier, pp 415–429
- Tomasella J, Pachepsky Ya, Crestana S, Rawls WJ (2003) Comparison of Two Techniques to Develop Pedotransfer Functions for Water Retention. *Soil Science Society of America Journal* 67:1085–1092. <https://doi.org/10.2136/sssaj2003.1085>

- Triberti L, Nastri A, Baldoni G (2016) Long-term effects of crop rotation, manure and mineral fertilisation on carbon sequestration and soil fertility. *European Journal of Agronomy* 74:47–55. <https://doi.org/10.1016/j.eja.2015.11.024>
- Vaast P, van Kanten R, Siles P, et al (2008) Biophysical Interactions Between Timber Trees and Arabica Coffee in Suboptimal Conditions of Central America. In: Nair PKR, Jose S, Gordon AM (eds) *Toward Agroforestry Design*. Springer Netherlands, Dordrecht, pp 133–146
- Van den Bergh I, Ramirez J, Staver C, et al (2012) Climate change in the subtropics: The impacts of projected averages and variability on banana productivity. *Acta Horticulturae* 89–99. <https://doi.org/10.17660/ActaHortic.2012.928.9>
- van Noordwijk M, Coe R, Sinclair FL, et al (2021) Climate change adaptation in and through agroforestry: Four decades of research initiated by Peter Huxley. *Mitigation and Adaptation Strategies for Global Change* 26:18. <https://doi.org/10.1007/s11027-021-09954-5>
- Van Oijen M (2020) *Bayesian Compendium*. Springer International Publishing
- Van Oijen M, Dauzat J, Harmand J-M, et al (2010a) Coffee agroforestry systems in Central America: II. Development of a simple process-based model and preliminary results. *Agroforestry Systems* 80:361–378. <https://doi.org/10.1007/s10457-010-9291-1>
- Van Oijen M, Dauzat J, Harmand J-M, et al (2010b) Coffee agroforestry systems in Central America: I. A review of quantitative information on physiological and ecological processes. *Agroforestry Systems* 80:341–359. <https://doi.org/10.1007/s10457-010-9294-y>
- Vezy R, Guerric LM, Mathias C, et al (2019) DynACof: A process-based model to study growth, yield and ecosystem services of coffee agroforestry systems. <https://doi.org/10.5281/zenodo.3246268>
- Virto I, Barré P, Burlot A, Chenu C (2012) Carbon input differences as the main factor explaining the variability in soil organic C storage in no-tilled compared to inversion tilled agrosystems. *Biogeochemistry* 108:17–26. <https://doi.org/10.1007/s10533-011-9600-4>