

What Is There to Gain? A Systematic Review on Psychosocial Co-Benefits of Climate Policies

Ahlke Kip^{1*}, Mira Tschorn^{2,3*}, Susanne Vogel^{4,5}, Michèle D. Birtel⁶, Carla L. Hendricks⁷,
Anne Voigt⁸, Silas Rooß^{2,8}, Julia Asbrand^{8,9}

*contributed equally

¹University of Freiburg, Medical Psychology and Medical Sociology, Faculty of Medicine,
Freiburg, Germany

²Freie Universität Berlin, Division of Clinical Psychological Intervention and Division of
Health Psychology, Berlin, Germany

³German Center for Mental Health (DZPG), partner site Berlin-Potsdam, Germany

⁴MSH Medical School Hamburg, Department of Psychology, Faculty of Human Sciences,
Hamburg, Germany

⁵MSH Medical School Hamburg, ICAN Institute for Cognitive and Affective Neuroscience,
Hamburg, Germany

⁶University of Greenwich, Institute for Lifecourse Development, London, UK

⁷Social and Preventive Medicine, Department of Sports and Health Sciences, Intra Faculty
Unit “Cognitive Sciences”, Faculty of Human Science and Faculty of Health Sciences
Brandenburg, Research Area Services Research and e-Health, Potsdam University,
Brandenburg, Germany

⁸Friedrich-Schiller-University Jena, Department of Psychology, Clinical Psychology of
Childhood and Adolescence, Jena, Germany

⁹German Center for Mental Health (DZPG), partner site Halle-Jena-Magdeburg, Germany

1

2 Corresponding author:

3 Julia Asbrand, PhD

4 julia.asbrand@uni-jena.de

5

6 DOI: 10.25646/14469

7 Submitted: 08.05.2026

8 Accepted: 12.08.2026

9 Published: DD.MM.2026

10

Abstract

Background: Climate change poses major risks to psychosocial health, yet little is known about potential psychosocial co-benefits of climate policies. This systematic review examines [1] psychosocial co-benefits associated with climate policy measures and [2] moderating factors shaping these effects.

Methods: Following PRISMA guidelines, $k = 21$ studies were identified and analysed using a PICOS framework and narrative synthesis.

Results: Climate policies, particularly air pollution control, environmental taxation, and nature-based urban interventions, were associated with improvements in subjective well-being, mental health, and social cohesion. Effects were mediated by environmental improvements and physical health gains, and varied by socioeconomic status, age, and contexts such as schools or rural vs. urban locations. Access to and actual use of interventions emerged as relevant but understudied mechanisms. Some studies indicated trade-offs, including unequal distribution of benefits and unintended social effects.

Conclusions: Overall, climate policies can generate psychosocial co-benefits, but evidence remains heterogeneous and methodologically limited. Future research should strengthen causal inference, standardise outcome measures, and examine long-term and distributional effects.

Keywords: Climate change, Climate policy, Psychosocial health, Mental health, Social cohesion, Well-being

Key messages

- Climate policies can generate psychosocial co-benefits, especially improvements in subjective well-being, life satisfaction, and stress-related mental health outcomes.

- Social cohesion, trust, and community engagement may improve when policies create accessible, usable, and shared public spaces.
- Physical health gains, particularly through reduced pollution exposure, better thermal comfort, sleep, and housing conditions, appear to be key pathways to psychosocial co-benefits.
- Effects are not evenly distributed: benefits vary by socioeconomic status, age, gender, vulnerability, and urban or rural context.
- The evidence base remains heterogeneous and methodologically limited, highlighting the need for stronger designs, validated outcomes, and long-term, equity-focused analyses.

1. Introduction

Human-induced climate change is one of the greatest challenges of this century. Beyond its ecological impacts, it poses numerous risks to human physical and psychosocial health [1, 2]. Simon Stevens, the former CEO of the UK National Health Service (NHS), described it as the ‘most profound long-term threat to the health of the nation’ [3, para. 5]. Given the severity of these risks, the term *climate crisis* is adopted here.

The climate crisis contributes directly to humanitarian emergencies—including wildfires, heatwaves, floods, and tropical storms—that are increasing in frequency and intensity [4]. According to the Intergovernmental Panel on Climate Change, between 3.3 and 3.6 billion people already live in areas highly vulnerable to climate change [4]. The World Health Organization (WHO) estimates that climate change will cause an additional 250,000 deaths per year between 2030 and 2050, primarily due to malaria, diarrhoea, malnutrition, and heat stress [5]. Environmental degradation—through rising sea levels, extreme weather events, and changing temperature and precipitation patterns—undermines the social and

ecological foundations of health, exacerbates inequalities, and limits access to essential services.

1.1. Psychosocial Health and the Climate Crisis

Psychosocial health is a multidimensional construct encompassing mental, social, spiritual, and material well-being, e.g. [6–8]. Building on this literature, psychosocial health can be understood as a dynamic state, shaped by the interaction between individuals and their social environments [8, 9]. In our current research, we adopt a definition of psychosocial health, which describes it as changing over time and existing on a spectrum from healthy to ill [9]. It includes a person’s experiences, perceptions, mental processes, behaviour, and lifestyle. It is closely linked with social relationships and broader social structures. A functioning body and consciousness enable interaction with the social environment. Psychosocial health can be strengthened by resources and support (e.g. emotional support) or weakened by stressors (e.g. work-related stress), with the individual being the central locus of this health. Thus, we apply a positive definition of psychosocial health as [1] mental well-being, including functioning in everyday life (e.g. sleep); [2] the absence of mental distress; and [3] social health, i.e. an individual’s connection with others and participation in all relevant aspects of life.

The impact of the climate crisis on mental health is increasingly recognised. For example, the Intergovernmental Panel on Climate Change acknowledges several associated mental health challenges and negative impacts on well-being [4]. The climate crisis affects psychosocial health through multiple pathways [10]. First, extreme weather events have direct psychological consequences, including increased rates of depression, anxiety, and post-traumatic stress disorder [10–12]. Second, gradual environmental changes, such as rising temperatures and air pollution, are associated with increased aggression, hospitalisations for mental illness, and cognitive impairments [13–15]. Third, indirect effects, including economic

instability, food insecurity, and forced migration, contribute to stress and reduced well-being [4, 16]. Finally, climate anxiety reflects the psychological burden arising from anticipated environmental loss and uncertainty and is significantly associated with mental health outcomes such as depression [17]. Younger populations may be particularly affected by climate anxiety, e.g. [18], though studies find varying levels of related impairment [19]. Other theoretical frameworks highlight community-level pathways, including the erosion of social capital and collective structures. In their causal pathways framework, Berry et al. [20] propose that climate change affects mental health through multiple routes: directly through traumatic climate-related events and indirectly through impacts on physical health as well as disruptions to environmental, social, and community conditions. These effects are further shaped by individual and contextual vulnerability, resulting in an uneven distribution of climate-related mental health impacts across individuals and communities. These impacts are unequally distributed. Vulnerable groups—including children, older adults, women, ethnic minorities, and individuals with low socioeconomic status—are disproportionately affected [5, 10, 21, 22]. This highlights the importance of considering moderators of psychosocial outcomes in climate-related research.

While the negative impact of the climate crisis and other ecological crises relating to overstepping the planet’s boundaries have repeatedly been addressed [23, 24], less focus has been on positive ‘side-effects’ of mitigation policies targeting climate and ecological protection.

1.2. Mitigation, Adaptation, and Co-Benefits

Climate policies primarily aim to mitigate emissions and support adaptation to climate impacts [4, 25]. However, beyond these primary goals, such policies may generate psychosocial co-benefits. For example, promoting active transport (e.g. cycling or walking) can improve physical health and reduce psychological distress in addition to their effect on emissions [26, 27]. Likewise, urban green spaces can enhance social cohesion and quality of

1 life [28, 29]. Participatory adaptation strategies may also increase perceived agency and
2 reduce feelings of helplessness [30]. Overall, nature-based interventions have gained
3 increasing attention in the past decade and systematic reviews and meta-analyses show
4 beneficial effects for mental health outcomes [31, 32].

5 Given the multiple pathways through which the climate crisis undermines
6 psychosocial health, climate policies may not only reduce future harm but actively promote
7 well-being. Despite these indications, the evidence base on co-benefits of climate policies
8 remains limited and fragmented. Existing research has largely focused on physical health
9 outcomes or on the perception and use of environmental features, rather than on the
10 psychosocial effects of policy interventions themselves [33]. Recent reviews and scenario-
11 based studies further suggest that mental health is rarely included in climate-related health
12 impact assessments, e.g. [34]. Similarly, reviews of environmental interventions (e.g.
13 afforestation projects) indicate that psychosocial co-benefits are seldom systematically
14 measured, highlighting a critical gap in the literature [35].

15 **1.3. The Current Study**

16 While the negative impacts of the climate crisis on psychosocial health are well-documented,
17 e.g. [1, 13], there is limited systematic evidence on whether and how the implementation of
18 climate policies contributes to psychosocial well-being. Previous reviews were limited to co-
19 benefits of prospective scenarios [34] or climate adaptation and mitigation strategies that
20 targeted wellbeing as primary outcome [35]. Addressing this gap is particularly relevant in the
21 context of rising global mental health challenges, especially among populations at risk such as
22 children and young people [18, 21, 36]. This review provides a systematic overview of
23 existing research on psychosocial co-benefits of climate policy measures. By synthesising
24 evidence across diverse interventions and contexts, it aims to identify key outcomes,
25 underlying mechanisms, and existing research gaps. Specifically, the review addresses the
26 following research questions:

(A) What psychosocial co-benefits are associated with climate policy measures?

(B) How do moderators—such as age, gender, socioeconomic status, and vulnerability—shape these relationships?

2. Methods

2.1. Search Strategy and Selection Process

We conducted multi-field searches in the databases Web of Science, PubMed, and PsycInfo. To identify relevant studies, search terms indicative of climate protection measures (e.g. environment* AND policy* OR “green infrastructure”) were combined with search terms indicative of psychosocial outcomes (e.g. well-being OR stress), and co-benefits (e.g. gain* OR benefit*) in the keyword search (see Supplementary material Table S1). Besides, we repeated the search with the MeSH term ‘Climatic Processes’ instead of keywords for climate protection measures. The search was restricted to human participants and peer-reviewed articles in English and German given the language proficiencies in the research team. The search included all publications up until November 22, 2024. We furthermore cross-checked references of related reviews [33, 35, 37–40] and searched Google Scholar for articles on psychosocial co-benefits of established climate-protection measures, e.g. Reducing Emissions from Deforestation and Forest Degradation (REDD+), Barcelona’s Superblocks, or the Regeneración Urbana, Calidad de Vida y Salud (RUCAS) [41–43]. All 39,349 hits were uploaded into Rayyan [44] for de-duplication and title-abstract screening.

A random sample of 1,350 hits was assessed by five blinded and independent reviewers each. Because Fleiss’s kappa, a chance-corrected measure of agreement among multiple raters, is sensitive to low inclusion rates [45], agreement was also assessed using the overall agreement rate. The overall intra-class correlation coefficient was 0.42 (0.16 – 0.64 between individual raters). However, the overall agreement rate between all reviewers was high at 93.6 % (95.5 – 97.4 between individual raters), meaning that raters made the same inclusion or exclusion decision in most cases. The remaining 30,482 hits were divided

1 between reviewers with all unclear cases being discussed with at least one other reviewer.
2 Full texts were assessed by two reviewers each with unclear cases being discussed among all
3 five reviewers.

4 The inclusion criteria according to the PICOS framework (Population, Intervention,
5 Comparison, Outcome, Study design); [46] were: P) human respondents (e.g. residents or
6 stakeholders) of all ages or human-related data (e.g. psychiatry admission data); I) mitigation
7 interventions or adaptation interventions that have an additional mitigation effect with a focus
8 on climate protection (requiring a change of environment instead of maintenance of a status
9 quo); C) studies with or without a control condition; O) psychosocial outcomes assessed as
10 co-benefits with a focus on health indicators versus health determinants (e.g. feelings of
11 safety); S) pre-post, controlled studies, and retrospective studies. We excluded P) outcomes
12 related to e.g. companies or households rather than individuals; I) pure adaptation
13 interventions without concurrent mitigation (e.g. installation of air conditioning),
14 interventions focusing on individual behaviour (e.g. physical exercise, change of nutrition),
15 disaster response programs, interventions that are not related to climatic environments (e.g.
16 school environments); C) n.a.; O) human-related measures that are not directly linked to
17 psychosocial outcomes (e.g. finances, physical health, education), non-human-related
18 outcomes (e.g. air pollution scores), outcomes on use, perception, or experience of natural
19 environments; ineligible operationalisation of psychosocial outcome (e.g. not inviting others
20 home as indicator of social isolation; expert opinion without direct involvement such as
21 NGOs); S) simulation or modelling studies, correlative studies (e.g. measuring closeness to
22 green space with mental health; see Table 1).

23 -- Table 1 about here --

2.2. Data Extraction


Data extraction was completed by one reviewer per study with all data being checked by a second reviewer. Any discrepancies were resolved by consensus. Extracted data included information on the study (e.g. quantitative vs. qualitative data, study design), sample (e.g. sampling strategy, age, gender), intervention (e.g. primary aim, year of implementation), outcome (e.g. psychosocial construct, instrument), and results (e.g. type of data analysis, primary and secondary findings such as on moderators).

3. Results

3.1. Study selection

The database search yielded 31,832 records, supplemented by 13 additional studies identified through other sources (mainly reference lists of other reviews or studies). After title and abstract screening, 229 full-text articles were assessed for eligibility, resulting in the inclusion of 21 studies. The study selection process is illustrated in Figure 1.

Figure 1: Flowchart of the literature selection process based on PRISMA. Source: Own presentation based on Page et al. [47]



3 3.2. Study Characteristics

4 *Sociodemographics.* Most studies were conducted in Asia ($k = 7$), followed by North
5 America ($k = 6$) and Europe ($k = 5$), while only a small number originated from South
6 America ($k = 1$) or Oceania ($k = 2$). Study designs included quantitative ($k = 12$), qualitative
7 ($k = 3$), and mixed-method approaches ($k = 6$) and comprised both cross-sectional and
8 longitudinal assessments. Only a minority of studies reported the use of a control group ($k =$
9 4). Sample sizes ranged from $n = 17$ [48] to $n = 61,100$ [49]. Most studies did not report a
10 mean age but instead described participants broadly as adults or youths. Samples
11 predominantly consisted of convenience samples such as students and teachers in schools, e.g.
12 [50], or residents of neighbourhoods, e.g. [51]. Gender distributions ranged from balanced
13 samples to predominantly female samples [52, 53], while several studies did not report
14 participants' gender, e.g. [54].

15 *Methodology*

16 Most studies were conducted from 2010 onwards, with one earlier study published in 1995
17 [55]. Outcomes covered a range of psychosocial constructs, including subjective well-being
18 or related constructs such as social cohesion or quality of life. Most studies applied an
19 observational or quasi-experimental design and used study-specific items to assess outcomes
20 (e.g. 'Generally, do you think your life is happy?', rated on a scale from 1 = not happy at all
21 to 5 = very happy [56]). Assessment methods included self-report measures administered via
22 interviews or questionnaires, e.g. [52, 53], other-reports [57], focus group discussions [43, 55,
23 57], and behavioural observations [58]. Follow-up periods varied widely across studies, with
24 most studies applying post-assessments directly after the intervention or within a year after.

25 *Types of interventions and environmental changes*

The included studies examined a diverse set of environmental interventions targeting both the built and natural environment. Several studies focused on *housing and building retrofits*, including energy-efficiency upgrades and comprehensive green renovations of residential buildings, e.g. [52, 59]. Another group of studies investigated *urban greening interventions*, such as green schoolyard renovations, e.g. [58], the creation of community gardens, e.g. [60], street tree planting, e.g. [57], green roofs, e.g. [60], and the development of urban green infrastructure including greenways and accessible workplace green spaces, e.g. [43, 51]. A smaller number of studies examined *nature-based interventions*, including ecosystem restoration projects such as mangrove rehabilitation [61]. Finally, some studies analysed *policy and governance interventions*, including environmental protection taxes [49], air pollution control policies, e.g. [62], and renewable energy programs [63], as well as one urban mobility intervention involving the implementation of superblocks [43].

3.3. Risk of Bias and Study Quality

The methodological quality of the included studies varied considerably. High-quality evidence was provided by quasi-experimental designs (e.g. difference-in-differences, regression discontinuity), which strengthened causal inference, e.g. [62, 64]. However, many studies relied on unvalidated outcome assessments, including single-item measures of psychosocial outcomes, e.g. [53, 62], while several employed qualitative approaches, either as stand-alone designs or through focus groups, interviews, or ethnographic methods, e.g. [43, 55, 57]. A smaller number of studies used validated instruments, such as quality-of-life measures including the Short Form Health-36 (SF-36) [54] the EQ-5D-3L and EQ-5D-5L by the EuroQol Research Foundation [43, 53], and the General Health Questionnaire (GHQ-12) [43]. Some studies applied multi-method designs combining quantitative and qualitative data, e.g. [43, 57], which allowed for a more comprehensive assessment of psychosocial outcomes. No randomised controlled trials were identified.

Methodological quality was additionally assessed using the Quality Assessment with Diverse Studies (QuADS) tool, which is specifically suited to reviews including heterogeneous quantitative, qualitative, and mixed-methods evidence. The QuADS comprises 13 criteria, each rated on a scale from 0 to 3, with 0 generally indicating no mention of the criterion and 3 indicating that the criterion was explicitly described, considered, or addressed. Given the aims of the present review, ratings were applied pragmatically with particular emphasis on the quality of methods used to assess psychosocial outcomes, rather than on all aspects of the broader environmental interventions. Assessments were conducted by one reviewer, with ambiguous cases discussed within the research team, and should therefore be interpreted as a structured and supportive overview rather than a strict hierarchical judgement of study quality.

QuADS ratings indicated overall moderate methodological quality across the included studies, with substantial variation reflecting the heterogeneous evidence base (see Table 2). Higher ratings were typically observed in studies with stronger quasi-experimental or longitudinal designs, clearer reporting of procedures, and the use of validated outcome measures, e.g. [51, 54, 62, 64]. Moderate ratings were common among studies using qualitative or mixed-methods approaches that were well aligned with their aims but more limited in sampling strategies or analytic transparency, e.g. [43, 57]. Lower ratings were generally associated with insufficient reporting of recruitment, limited justification of analytic methods, or the use of less rigorous and predominantly descriptive designs. Across studies, research aims and study settings were usually well-described, whereas stakeholder involvement and detailed reporting of recruitment or analytic decision-making were less consistently addressed.

3.4. Main Findings of Co-Benefits for Psychosocial Outcomes

Well-being and Life Satisfaction

Thirteen studies examined the effects of climate-protection interventions on well-being. Well-being was operationalised using indicators such as happiness, life satisfaction, morale, perceived quality of life, positive affect, and perceived psychological functioning [43, 51, 52, 54, 56, 57, 59, 60, 62, 64–67].

Overall, most studies reported positive associations between environmental improvements and well-being-related outcomes. For instance, environmental interventions to reduce excessive temperatures were further associated with improvements in quality of life ($t = 3.179, p < .01$) and increased sleep duration ($t = 2.150, p < .05$) [52]. Similarly, several urban greening interventions reported improvements in perceived well-being and morale. Perceived well-being encompasses positive psychological functioning, including emotional stability, life satisfaction, and the ability to cope with daily demands, e.g. [51], whereas morale reflects an individual's overall psychological state, including feelings of motivation, satisfaction, and being valued within their environment [57].

In workplace settings, the introduction of visible and accessible green infrastructure was perceived as contributing positively to staff well-being, with employees reporting increased happiness and a greater sense of being valued by their employer. In one study, respondents were asked whether working in an environment with more green spaces makes them happier (1-5 scale). Both before and after the intervention, 86% of respondents selected one of the two most positive response categories. However, the proportion selecting the most positive response category increased by 13% from 54% to 67% ($p = .05$) [57]. Evidence from neighbourhood and urban-scale interventions, altering the built environment, similarly pointed towards improvements in well-being [e.g., 57]. Superblock mobility schemes that reduced traffic and increased pedestrian space, were reported to enhance residents' perceived

well-being and feelings of calmness and pleasure through improved aesthetics, reduced noise and pollution, and increased opportunities for social interaction and outdoor activities [43]. Reductions in air pollution might be a critical mechanism as evidence from policy-based environmental interventions showed that while air pollution was related to reduced happiness and life satisfaction [67], new air pollution governance led to improvements in subjective well-being and self-rated health [62, 66].

However, findings were not entirely consistent. Some studies reported no statistically significant changes in mental well-being despite improvements in self-reported quality of life which can be considered a broader, multidimensional construct not limited to mental health (e.g. pre–post difference $p = .226$ unadjusted; $p = .524$ adjusted, controlling for sociodemographic and other factors) [51]. One other study reported small declines in overall well-being scores following environmental changes, although these differences were not statistically tested [57]. Further, one study reported that certain policy measures such as pollutant discharge fees could reduce well-being due to inconvenience in daily life, despite environmental benefits [56]. Overall, evidence suggests net positive but context-dependent effects on well-being, e.g. limited to urban areas [64]. For an overview of all studies see Table 2.

-- Table 2 about here --

Psychopathology Outcomes

Nine studies examined the effects of environmental interventions on mental health outcomes, including indicators of emotional distress, perceived stress and anxiety, depressive symptoms, sleep, and cognitive functioning [48, 49, 51–54, 57, 59, 64]. Overall, findings were heterogeneous, with some studies reporting improvements, while others found no significant changes or even increases in symptoms.

Some evidence suggests that environmental improvements may contribute to reductions in stress and emotional distress. For instance, environmental interventions to reduce excessive temperatures were associated with decreased emotional distress ($t = -2.085$, $p < .05$) [52]. Similarly, improvements in housing conditions were linked to reductions in stress and anxiety related to energy insecurity and difficulties paying energy bills [59]. One study further reported improvements in sleep duration and memory following environmental interventions, with increases reaching statistical significance [49]. Qualitative evidence from urban environmental interventions further supported these findings, with participants describing a more relaxed atmosphere and reduced feelings of stress following environmental improvements [59].

However, quantitative evidence across studies was not consistent. One longitudinal, quasi-experimental study found no statistically significant pre–post changes in mental health indicators within intervention groups, including measures of psychological well-being and the number of days with poor mental health [53]. Taken together, the evidence suggests that environmental interventions may contribute to reductions in stress-related mental health outcomes, particularly through improvements in thermal comfort, housing conditions, and neighbourhood environments.

Social Health Outcomes

Eleven studies examined the effects of environmental interventions on social and community outcomes, including indicators such as social interactions, bullying or teasing, neighbourhood climate, and perceived community cohesion [43, 48, 50, 51, 55, 57–59, 61, 63, 65].

Observational studies of schoolyard and neighbourhood interventions indicated increases in positive social interactions after environmental changes. For example, the proportion of children observed interacting with others increased from 63.0 % at baseline to 73.3 % following a green schoolyard renovation, with a corresponding rise in positive interactions (27.1 % to 35.2 %) and only very low levels of negative interactions [50]. Similarly, another

1 observational study reported shifts in the quality of interactions, including a decrease in
2 negative interactions (2.4 % to 1.1 %) and an increase in positive interactions (44.3 % to 48.7
3 %) following a green schoolyard renovation [58]. Retrospective reports from caregivers and
4 school staff also suggested reductions in teasing and bullying after schoolyard renovations,
5 although these changes were not statistically tested and no further changes were observed
6 from baseline to follow-up assessments [50]. Several studies further reported improvements in
7 perceptions of social climate and community relationships. For instance, one study found a
8 significant improvement in perceptions of neighbourhood climate ($t = -2.62, p < .05$) and a
9 reduction in overall school bullying ($d = 0.405, p < .01$), alongside improvements in school–
10 community relationships ($d = 0.408, p < .01$) [58]. The effect sizes are moderate in size
11 according to Cohen’s conventions [68]. In some cases, perceptions of social benefits such as
12 community strengthening, empowerment, and social recognition were reported to outweigh
13 potential social costs of the intervention [61]. Overall, qualitative findings suggested that
14 environmental interventions may facilitate social interaction by creating shared spaces for
15 social activities and reducing barriers to informal encounters among residents [55].

16 Taken together, the available evidence suggests that environmental interventions may
17 contribute to improvements in social interactions and perceptions of community cohesion,
18 particularly where interventions create shared public spaces or opportunities for community
19 engagement, although quantitative findings remain heterogeneous across studies.

21 *Negative Side-Effects*

22 Overall, systematic evidence on adverse psychosocial effects of environmental interventions
23 remains scarce, as relatively few studies explicitly examined negative outcomes. In a small
24 number of studies, negative trends were observed; however, these were typically not
25 attributable to the intervention itself. For example, one quasi-experimental study reported a
26 small but statistically significant decline in reported social networks over time (4.0 to 3.9; $p <$

.001), although similar changes were observed in the control group and differences between groups were not significant [51]. Additionally, some findings suggest that effects may vary depending on contextual factors such as region and time-lag to implementation, with both positive and negative associations observed [56]. Qualitative evidence further indicates that environmental interventions can give rise to unintended social tensions or unmet expectations [61].

Potential Moderators

Only few studies investigated the influence of potential moderators. Age [49, 50, 66] and gender [50, 61] repeatedly appeared as sociodemographically relevant constructs. Further, individuals with higher income were more likely to derive well-being benefits from environmental improvements, whereas lower-income groups prioritised material conditions [67]. However, individual attitudes towards and perceptions of the implemented interventions were also relevant, e.g. [63]. Finally, differences between urban and rural populations suggest that different benefits may emerge depending on the context, with urban residents reporting improvements in subjective well-being, while rural populations showed more pronounced physical health gains [66].

4. Discussion

This review aimed to [1] identify psychosocial co-benefits associated with climate policy measures and [2] examine moderators shaping these relationships. Surprisingly few studies have targeted this specific subject to date. Overall, the evidence indicates that climate policies—particularly air pollution control, environmental taxation, and nature-based urban interventions—are associated with positive psychosocial outcomes, although effects are heterogeneous and context-dependent. Across the included studies, environmental interventions were most consistently associated with improvements in subjective well-being and mental health, including reductions in stress, anxiety, and sleep disturbances, as well as increases in life satisfaction and emotional stability [49, 62, 64]. These findings extend prior

1 literature linking environmental quality to well-being [69] by demonstrating that policy-
2 driven environmental improvements can actively enhance psychosocial health. Also, several
3 studies highlight social co-benefits, including increased social cohesion, trust, and community
4 engagement, particularly in the context of community-based or nature-based interventions
5 [63]. These findings suggest that climate policies can generate collective psychosocial
6 benefits, not limited to individual-level outcomes.

7 Consistent with broader health–well-being frameworks [8], physical health
8 improvements may function as a central mediating pathway. Reductions in pollution exposure
9 are associated with lower morbidity and improved self-rated health, which in turn contribute
10 to higher well-being [49, 62]. However, it has to be considered, that several studies found no
11 statistically significant changes in mental health measures, e.g. [51], and in some cases, a
12 slight decline in overall well-being [57] or a decrease in social networks, e.g. [51], was
13 reported. These findings can be understood through a resource-based framework: climate
14 policies may strengthen psychosocial health by improving individual resources such as
15 physical health, social resources such as cohesion and community engagement, and material
16 resources such as housing conditions and reduced economic burden [70]. This aligns with
17 planetary health approaches, which emphasises the interdependence of human well-being,
18 social conditions, and healthy environments e.g. [71].

19 **4.1. Potential Moderators**

20 Socioeconomic status emerged as a key moderator, although findings were mixed:
21 some studies suggest that individuals with higher socioeconomic resources derive greater
22 psychosocial benefits, potentially due to increased capacity to utilise improved environments
23 [67]. Conversely, other findings indicate stronger health gains among socioeconomically
24 disadvantaged groups due to higher baseline exposure, suggesting differential benefit
25 pathways rather than uniform effects. Older adults appear to benefit substantially from
26 environmental policies, particularly through improvements in physical and cognitive health

[49]. Evidence for children and adolescents—primarily from green infrastructure studies—indicates benefits for psychological development and well-being, although these are less frequently examined in policy contexts. Finally, the observed difference in co-benefits for urban and rural residents [49] highlight the importance of contextual and environmental baseline conditions. Several studies indicate that effects vary depending on policy design, implementation intensity, and regional conditions, including non-linear relationships between regulation and well-being [56]. This suggests that psychosocial co-benefits are not inherent to policies per se but depend on how policies are implemented and experienced.

A cross-cutting issue relevant to both research questions is the distinction between access to climate-related interventions and their actual use. Evidence from urban green space research indicates that availability alone does not guarantee psychosocial benefits, which depend on engagement, perceived safety, and usability [72, 73]. This implies that some null or weak findings in the included studies may reflect limited use rather than absence of effect, suggesting an important behavioural mechanism that is insufficiently captured in current research.

Additionally, the identified co-benefits may be incompletely captured due to the predominance of short-term study designs. Climate policies can trigger longer-term systemic changes, such as increased urban attractiveness, migration, and social development, which may further influence well-being over time. At the same time, self-selection processes [74, 75] complicate interpretation, as individuals may sort into environments that match their preferences or resources. Additionally, urban greening may contribute to gentrification and rising housing costs, potentially offsetting benefits for vulnerable populations, e.g. [76, 77]. Finally, certain interventions may impose economic or psychosocial burdens, such as increased labour demands or effects on income inequality, e.g. [78]. These findings indicate that psychosocial co-benefits should be considered alongside distributional and contextual trade-offs.

4.2. Methodological Considerations

Original studies. This review provides a first synthesis of the evidence, indicating that the field of psychosocial co-benefits of climate policies remains in its infancy. While the limited number of high-quality studies restricts the conclusions that can currently be drawn, a systematic synthesis can be particularly valuable at this early stage of the field. It allows for an identification of first indications of potential co-benefits and, importantly, a specification of methodological improvements needed to establish a more robust evidence base in the future. This interpretation is consistent with the QuADS assessment, which indicated predominantly moderate methodological quality across the included studies. Moderate ratings often reflected a reasonable alignment between study aims and methods, particularly in qualitative and mixed-methods studies, but also limitations in sampling strategies, analytic transparency, or reporting detail. Across studies, aims and settings were usually well described, whereas recruitment procedures, stakeholder involvement, and analytic decision-making were less consistently addressed. While negative effects of the climate crisis on both general symptoms of psychopathology and established mental disorders have been described quite regularly [13], more thorough attention has yet to be directed towards preventing and decreasing clinical levels of psychopathology through climate policies. This crucial interdisciplinary topic faces several key challenges that need to be addressed in future research to improve the reliability of its conclusions. First, psychosocial constructs were defined very differently between studies and not always as intended by psychological and health sciences [8]. They were further limited to narrow levels of psychosocial health, e.g. evidence for spiritual or existential dimensions (e.g. meaning, place attachment) has so far been less systematically assessed and remains an area for future research. Second, given that the main goal of the interventions were changes in the environment, assessments of co-benefits were sometimes only added post-hoc in a retrospective manner, e.g., [50], or assessed only indirectly via observation [50, 58], resulting in an enormous heterogeneity of methods.

For example, our review focused on individual-level psychosocial health, while several studies assessed this outcome at household level [79, 80]. As differences in reporters' perspectives are the norm rather than the exception [81, 82], a multi-person approach should be preferred over a single-respondent household perspective. Additionally, outcome measures were often insufficient, with most studies relying on single-item indicators rather than validated, psychometrically robust scales [53, 62]. Third, only a few studies applied mediation analyses to investigate the indirect pathways from environmental changes to psychosocial benefits. For example, climate protection measures could induce a sense of security [83–86], reinstate attention, and increase physical health [4, 30, 65, 86]. Thus, all findings should be interpreted very cautiously and rather serve to generate impulses for future research. Finally, assessments of long-term effects, including follow-up, remain limited [49, 62, 64], although such effects may be particularly important, given that climate protection measures are likely to unfold their impact over the longer term rather than the short term [1, 87, 88].

Current study. Regarding limitations of our systematic review, this review was not pre-registered, and inclusion criteria were iteratively refined due to the conceptual diversity of climate-related interventions. Further, as the field is very broad and heterogeneous, we focused on a psychosocial perspective [8], which could be broadened to societal as well as somatic levels and did not include mediators, thus providing only a first glimpse into the field and presenting potential future lines of research.

4.3. Implications for Research and Society

The findings suggest that climate policies can generate psychosocial co-benefits across multiple domains, but these are not uniform and depend on contextual and individual factors. Future research should employ longitudinal and experimental designs [89], systematically assess moderators and vulnerable groups, including a broad perspective on psychosocial health and regions which might suffer even more from negative effects of the climate crisis such as sub-Saharan Africa, e.g. [90, 91]. Of note, none of the studies included here has

1 focused on the continent of Africa. Also, next to indicators for psychosocial health, a broader
2 network of health-related determinants should be considered such as gender empowerment
3 [92], food and nutrition security [93], and days in school/at work [94]. Current evidence also
4 highlights limitations in interdisciplinary integration and measurement consistency, which
5 should be addressed to improve evidence quality. This calls for interdisciplinary research
6 combining expertise on both climate interventions and high-quality psychosocial assessments.

7 For society and policy, the findings suggest that climate policies may generate benefits
8 beyond environmental and physical health outcomes. Measures that reduce pollution, improve
9 housing or thermal conditions, or provide accessible shared spaces may support everyday
10 well-being and social interaction. While psychosocial outcomes are typically not the primary
11 objective of climate policy, some measures may explicitly pursue well-being or social
12 benefits alongside environmental goals. More generally, psychosocial effects should be
13 considered as potential co-benefits of climate action rather than assumed to occur across all
14 policies. Their assessment should consider accessibility, usability, and the distribution of
15 benefits and burdens across population groups. This highlights the need for standardised
16 frameworks to assess psychosocial impacts, similar to existing approaches in urban health
17 research [91, 95].

19 **Data protection and ethics**

20 Not applicable.

22 **Data availability**

23 Not applicable.

25 **Funding**

26 No funding.

28 **Individual author contributions**

29 Substantial contributions to the conception or design of the work: AK, MT, SV, MDB, CH,
30 AV, SR, JA; to the acquisition of the data for the work: AK, MT, SV, MDB, SR, JA; to the
31 analysis of the data for the work: AK, MT, SV, MDB, CH, AV, SR, JA; to the interpretation
32 of the data for the work: AK, MT, SV, MDB, CH, AV, SR, JA. Drafting the work: AK, MT,

1 JA; revising the work critically for important intellectual content: SV, MDB, CH, AV, SR.
2 Final version of the manuscript read and approved for publication: AK, MT, SV, MDB, CH,
3 AV, SR, JA.

4
5 **Bitte für die deutsche Version wörtlich übernehmen:**

6 Maßgebliche Beiträge zu Konzeption oder Design der Arbeit: AK, MT, SV, MDB, CH, AV,
7 SR, JA; zur Erhebung der verwendeten Daten: AK, MT, SV, MDB, SR, JA; zur Analyse der
8 verwendeten Daten: AK, MT, SV, MDB, CH, AV, SR, JA; zur Interpretation der
9 verwendeten Daten: AK, MT, SV, MDB, CH, AV, SR, JA. Ausarbeitung des Manuskripts:
10 AK, MT, JA; kritische Überarbeitung bedeutender Inhalte: SV, MDB, CH, AV, SR. Finale
11 Version des Manuskripts gelesen und der Veröffentlichung zugestimmt: AK, MT, SV, MDB,
12 CH, AV, SR, JA.

13
14 **Declaration of generative artificial intelligence (AI) and AI-assisted
15 technologies**

16 Not applicable.

17
18 **Conflicts of interest**

19 The authors declare that there is no conflict of interest.
20
21

References

1. Romanello M, Napoli CD, Green C, Kennard H, Lampard P, Scamman D, et al. The 2023 report of the Lancet Countdown on health and climate change: the imperative for a health-centred response in a world facing irreversible harms. *Lancet*. 2023;402(10419):2346–94. doi: 10.1016/S0140-6736(23)01859-7.
2. Cuijpers P, Kumar M, Karyotaki E. Climate Change and Mental Health—Time to Act Now. *JAMA Psychiatry*. 2023;80(12):1183–4. doi: 10.1001/jamapsychiatry.2023.3413.
3. National Health System (NHS). NHS England » NHS becomes the world's first national health system to commit to become 'carbon net zero', backed by clear deliverables and milestones. 2020 [cited 07.04.2026]. Available from: <https://www.england.nhs.uk/2020/10/nhs-becomes-the-worlds-national-health-system-to-commit-to-become-carbon-net-zero-backed-by-clear-deliverables-and-milestones/>
4. IPCC. Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. In: Core Writing Team, H. Lee and J. Romero, editors. Intergovernmental Panel on Climate Change (IPCC). 2023. p. 35–115. doi: 10.59327/IPCC/AR6-9789291691647.5. World Health Organization. Climate change. 2023 [cited 05.11.2024]. Available from: <https://www.who.int/news-room/fact-sheets/detail/climate-change-and-health>
5. Eiroa-Orosa FJ. Understanding Psychosocial Wellbeing in the Context of Complex and Multidimensional Problems. *Int J Environ Res Public Health*. 2020;17(16):5937. doi: 10.3390/ijerph17165937.
6. Martikainen P, Bartley M, Lahelma E. Psychosocial determinants of health in social epidemiology. *Int J Epidemiol*. 2002;31(6):1091–3. doi: 10.1093/ije/31.6.1091.
7. Peter KA, Helfer T, Golz C, Halfens RJG, Hahn S. Development of an Interrelated Definition of Psychosocial Health for the Health Sciences Using Concept Analysis. *J Psychosoc Nurs Ment Health Serv*. 2022;60(6):19–26. doi: 10.3928/02793695-20211214-02.
8. Helfer T, Faeh D, Luijckx E, Frey S, Berg A, Peter KA. The Connection between Health Promotion, Prevention, and Psychosocial Health: An Innovative Action Model. *Open Public Health J*. 2020;13(1):850–5. doi: 10.2174/1874944502013010850.
9. Clayton S. Climate change and mental health. *Curr Environ Health Rep*. 2021;8(1):1–6. doi: 10.1007/s40572-020-00303-3.
10. Beaglehole B, Mulder RT, Frampton CM, Boden JM, Newton-Howes G, Bell CJ. Psychological distress and psychiatric disorder after natural disasters: systematic review and meta-analysis. *Br J Psychiatry*. 2018;213(6):716–22. doi: 10.1192/bjp.2018.210.
11. Kip A, Hoppen TH, Tam KP, Bryant R, Lowe SR, Morina N. Prevalence of common mental disorders in survivors of natural hazards meta-analytic comparison of exposed vs. non-exposed samples. *J Environ Psychol*. 2025;106:102690. doi: 10.1016/j.jenvp.2025.102690.
12. Cuijpers P, Miguel C, Ciharova M, Kumar M, Brander L, Kumar P, et al. Impact of climate events, pollution, and green spaces on mental health: an umbrella review of meta-analyses. *Psychol Med*. 2023;53(3):638–53. doi: 10.1017/S0033291722003890.

14. Hansen AL, Bi P, Ryan P, Nitschke M, Pisaniello D, Tucker G. The effect of heat waves on hospital admissions for renal disease in a temperate city of Australia. *Int J Epidemiol*. 2008;37(6):1359–65. doi: 10.1093/ije/dyn165.
15. Miles-Novelo A, Anderson CA. Climate change and psychology: Effects of rapid global warming on violence and aggression. *Curr Clim Change Rep*. 2019;5(1):36–46. doi: 10.1007/s40641-019-00121-2.
16. Carleton TA, Hsiang SM. Social and economic impacts of climate. *Science*. 2016;353(6304):aad9837. doi: 10.1126/science.aad9837.
17. Cosh SM, Ryan R, Fallander K, Robinson K, Tognela J, Tully PJ, et al. The relationship between climate change and mental health: a systematic review of the association between eco-anxiety, psychological distress, and symptoms of major affective disorders. *BMC Psychiatry*. 2024;24(1):833. doi: 10.1186/s12888-024-06274-1.
18. Hickman C, Marks E, Pihkala P, Clayton S, Lewandowski RE, Mayall EE, et al. Climate anxiety in children and young people and their beliefs about government responses to climate change: a global survey. *Lancet Planet Health*. 2021;5(12):e863–73. doi: 10.1016/S2542-5196(21)00278-3.
19. Wullenkord MC, Johansson M, Loy LS, Menzel C, Reese G. Go out or stress out? Exploring nature connectedness and cumulative stressors as resilience and vulnerability factors in different manifestations of climate anxiety. *J Environ Psychol*. 2024;95:102278. doi: 10.1016/j.jenvp.2024.102278.
20. Berry HL, Bowen K, Kjellstrom T. Climate change and mental health: a causal pathways framework. *Int J Public Health*. 2010;55(2):123–32. doi: 10.1007/s00038-009-0112-0.
21. Anjum G, Aziz M. Climate change and gendered vulnerability: A systematic review of women’s health. *Womens Health (Lond)*. 2025;21:17455057251323645. doi: 10.1177/17455057251323645.
22. Benevolenza MA, DeRigne L. The impact of climate change and natural disasters on vulnerable populations: A systematic review of literature. *J Hum Behav Soc Environ*. 2019;29(2):266–81. doi: 10.1080/10911359.2018.1527739.
23. Richardson K, Steffen W, Lucht W, Bendtsen J, Cornell SE, Donges JF, et al. Earth beyond six of nine planetary boundaries. *Sci Adv*. 2023;9(37):eadh2458. doi: 10.1126/sciadv.adh2458.
24. Rockström J, Steffen W, Noone K, Persson Å, Chapin FS, Lambin E, et al. Planetary boundaries: Exploring the safe operating space for humanity. *Ecology and Society*. 2009 [cited 05.11.2024];14(2). Available from: <https://www.jstor.org/stable/26268316>
25. Tosun J, Peters BG. The politics of climate change: Domestic and international responses to a global challenge. *Int Political Sci Rev*. 2021;42(1):3–15. doi: 10.1177/0192512120975659.
26. Green S, Sakuls P, Levitt S. Cycling for health: Improving health and mitigating the climate crisis. *Can Fam Physician*. 2021;67(10):739–42. doi: 10.46747/cfp.6710739.
27. Smith PJ, Merwin RM. The role of exercise in management of mental health disorders: An integrative review. *Annu Rev Med*. 2021;72:45–62. doi: 10.1146/annurev-med-060619-022943.
28. Castillo MD, Anenberg SC, Chafe ZA, Huxley R, Johnson LS, Kheirbek I, et al. Quantifying the health benefits of urban climate mitigation actions: Current state of the epidemiological evidence and application in health impact assessments. *Front Sustain Cities*. 2021;3. doi: 10.3389/frsc.2021.768227.

29. Liu M, Huang Y, Hiscock R, Li Q, Bi J, Kinney PL, et al. Do climate change policies promote or conflict with subjective wellbeing: A case study of Suzhou, China. *Int J Environ Res Public Health*. 2016;13(3):3. doi: 10.3390/ijerph13030344.
30. Meenar MR. Integrating placemaking concepts into Green Stormwater Infrastructure design in the City of Philadelphia. *Environ Pract*. 2019;21(1):4–19. doi: 10.1080/14660466.2019.1568121.
31. Coventry PA, Brown JVE, Pervin J, Brabyn S, Pateman R, Breedvelt J, et al. Nature-based outdoor activities for mental and physical health: Systematic review and meta-analysis. *SSM Popul Health*. 2021;16:100934. doi: 10.1016/j.ssmph.2021.100934.
32. Heinen-Stach D, Herbold E, Hendricks CL, Zuch N, Rapp MA, Birtel MD, et al. The effectiveness of nature-based interventions on mental health: A systematic review. *Appl Psychol Health Well Being*. 2026;18(4):e70164. doi: 10.1111/aphw.70164.
33. Lamb WF, Steinberger JK. Human well-being and climate change mitigation. *Wiley Interdiscip Rev Clim Change*. 2017;8(6):e485. doi: 10.1002/wcc.485.
34. Moutet L, Bernard P, Green R, Milner J, Haines A, Slama R, et al. The public health co-benefits of strategies consistent with net-zero emissions: a systematic review. *Lancet Planet Health*. 2025;9(2):e145–56. doi: 10.1016/S2542-5196(24)00330-9.
35. Flores EC, Brown LJ, Kakuma R, Eaton J, Dangour AD. Mental health and wellbeing outcomes of climate change mitigation and adaptation strategies: a systematic review. *Environ Res Lett*. 2023;19(1):014056. doi: 10.1088/1748-9326/ad153f.
36. McGorry PD, Mei C, Dalal N, Alvarez-Jimenez M, Blakemore SJ, Browne V, et al. The Lancet Psychiatry Commission on youth mental health. *Lancet Psychiatry*. 2024;11(9):731–74. doi: 10.1016/S2215-0366(24)00163-9.
37. Gao J, Kovats S, Vardoulakis S, Wilkinson P, Woodward A, Li J, et al. Public health co-benefits of greenhouse gas emissions reduction: A systematic review. *Sci Total Environ*. 2018;627:388–402. doi: 10.1016/j.scitotenv.2018.01.193.
38. Karlsson M, Alfredsson E, Westling N. Climate policy co-benefits: a review. *Clim Policy*. 2020;20(3):292–316. doi: 10.1080/14693062.2020.1724070.
39. Lamb WF, Antal M, Bohnenberger K, Brand-Correa LI, Müller-Hansen F, Jakob M, et al. What are the social outcomes of climate policies? A systematic map and review of the ex-post literature. *Environ Res Lett*. 2020;15(11):113006. doi: 10.1088/1748-9326/abc11f.
40. Shonkoff SB, Morello-Frosch R, Pastor M, Sadd J. The climate gap: environmental health and equity implications of climate change and mitigation policies in California—a review of the literature. *Clim Change*. 2011;109(1):485–503. doi: 10.1007/s10584-011-0310-7.
41. Alexander S, Nelson CR, Aronson J, Lamb D, Cliquet A, Erwin KL, et al. Opportunities and Challenges for Ecological Restoration within REDD+. *Restor Ecol*. 2011;19(6):683–9. doi: 10.1111/j.1526-100X.2011.00822.x.
42. González F, Baeza F, Valdebenito R, Sánchez BN, Diez-Roux A, Vives A. Improvements in habitability and housing satisfaction after dwelling regeneration in social housing complexes. The RUCAS study. *Soc Sci Med*. 2024;355:117090. doi: 10.1016/j.socscimed.2024.117090.
43. Pérez K, Palència L, López MJ, León-Gómez BB, Puig-Ribera A, Gómez-Gutiérrez A, et al. Environmental and health effects of the Barcelona superblocks. *BMC Public Health*. 2025;25(1):634. doi: 10.1186/s12889-025-21835-z.

44. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016;5(1):210. doi: 10.1186/s13643-016-0384-4.
45. McKenzie KA, Mahnken JD. Simulating and estimating agreement in the presence of multiple raters and covariates. *Stat Med*. 2023;42(11):1687–98. doi: 10.1002/sim.9694.
46. Higgins JPT, Cochrane Collaboration, editors. *Cochrane handbook for systematic reviews of interventions*. Second edition. Hoboken, NJ: Wiley-Blackwell; 2019. 694 p. (Cochrane book series).
47. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi: 10.1136/bmj.n71.
48. Hoffman AJ. Going “green” from gray: Providing opportunities of community development and sustainability within a correctional facilities program. *J Prev Interv Community*. 2020;48(3):272–80. doi: 10.1080/10852352.2019.1625605.
49. Wang H, Zhou R, Luo H, Liang S, Kong Y. The impact of environmental protection tax on the health of middle-aged and older adults: evidence from CHARLS data in China. *Front Public Health*. 2024;12. doi: 10.3389/fpubh.2024.1446248.
50. Bates CR, Bohnert AM, Gerstein DE. Green Schoolyards in Low-Income Urban Neighborhoods: Natural Spaces for Positive Youth Development Outcomes. *Front Psychol*. 2018;9. doi: 10.3389/fpsyg.2018.00805.
51. Hunter RF, Adlakha D, Cardwell C, Cupples ME, Donnelly M, Ellis G, et al. Investigating the physical activity, health, wellbeing, social and environmental effects of a new urban greenway: a natural experiment (the PARC study). *Int J Behav Nutr Phys Act*. 2021;18(1):142. doi: 10.1186/s12966-021-01213-9.
52. Ahrentzen S, Erickson J, Fonseca E. Thermal and health outcomes of energy efficiency retrofits of homes of older adults. *Indoor Air*. 2016;26(4):582–93. doi: 10.1111/ina.12239.
53. Breyse J, Dixon SL, Jacobs DE, Lopez J, Weber W. Self-Reported Health Outcomes Associated With Green-Renovated Public Housing Among Primarily Elderly Residents. *J Public Health Manag Pract*. 2015;21(4):355. doi: 10.1097/PHH.0000000000000199.
54. Howden-Chapman P, Matheson A, Crane J, Viggers H, Cunningham M, Blakely T, et al. Effect of insulating existing houses on health inequality: cluster randomised study in the community. *BMJ*. 2007;334(7591):460. doi: 10.1136/bmj.39070.573032.80.
55. Gill B, Simeoni E. Residents’ perceptions of an environmental enhancement project in Australia. *Health Promot Int*. 1995;10(4):253–9. doi: 10.1093/heapro/10.4.253.
56. Song Y, Guo S, Zhang M. Will environmental regulations affect subjective well-being?—a cross-region analysis in China. *Environ Sci Pollut Res*. 2019;26(28):29191–211. doi: 10.1007/s11356-019-06147-1.
57. Cinderby S, Bagwell S. Exploring the co-benefits of urban green infrastructure improvements for businesses and workers’ wellbeing. *Area*. 2018;50(1):126–35. doi: 10.1111/area.12361.
58. Bohnert AM, Nicholson LM, Mertz L, Bates CR, Gerstein DE. Green schoolyard renovations in low-income urban neighborhoods: Benefits to students, schools, and the surrounding community. *Am J Community Psychol*. 2022;69(3–4):463–73. doi: 10.1002/ajcp.12559.

59. Hernández D, Phillips D. Benefit or burden? Perceptions of energy efficiency efforts among low-income housing residents in New York City. *Energy Res Soc Sci*. 2015;8:52–9. doi: 10.1016/j.erss.2015.04.010.
60. Kim E, Jung J, Hapsari G, Kang S, Kim K, Yoon S, et al. Economic and environmental sustainability and public perceptions of rooftop farm versus extensive garden. *Build Environ*. 2018;146:206–15. doi: 10.1016/j.buildenv.2018.09.046.
61. Richerzhagen C, Rodríguez de Francisco JC, Weinsheimer F, Döhnert A, Kleiner L, Mayer M, et al. Ecosystem-Based Adaptation Projects, More than just Adaptation: Analysis of Social Benefits and Costs in Colombia. *Int J Environ Res Public Health*. 2019;16(21):4248. doi: 10.3390/ijerph16214248.
62. Xie T, Wang Y, Yuan Y. Health Benefits from Improved Air Quality: Evidence from Pollution Regulations in China's 2+26 Cities. *Environ Resour Econ*. 2024;87(5):1175–221. doi: 10.1007/s10640-024-00860-3.
63. Tsagkari M, Roca J, Stephanides P. Sustainability of local renewable energy projects: A comprehensive framework and an empirical analysis on two islands. *Sustain Dev*. 2022;30(5):1155–68. doi: 10.1002/sd.2308.
64. Zhao C, Li X. Understanding government pollution control actions and well-being: Evidence from micro individuals. *Air Qual Atmos Health*. 2024;17(9):1917–35. doi: 10.1007/s11869-024-01551-0.
65. Marshall CAM, Wilkinson MT, Hadfield PM, Rogers SM, Shanklin JD, Eversham BC, et al. Urban wildflower meadow planting for biodiversity, climate and society: An evaluation at King's College, Cambridge. *Ecol Solut Evid*. 2023;4(2):e12243. doi: 10.1002/2688-8319.12243.
66. Song Y, Li M, Zhang M, Sun X. Study on the impact of air pollution control on urban residents' happiness from microscopic perspective. *J Clean Prod*. 2019;229:1307–18. doi: 10.1016/j.jclepro.2019.05.070.
67. Wang J, Tang D. Air Pollution, Environmental Protection Tax and Well-Being. *Int J Environ Res Public Health*. 2023;20(3):2599. doi: 10.3390/ijerph20032599.
68. Cohen J. Statistical power analysis for the behavioral sciences. Routledge; 2013 [cited 09.04.2026]. Available from: <https://api.taylorfrancis.com/content/books/mono/download?identifierName=doi&identifierValue=10.4324/9780203771587&type=googlepdf>
69. Levinson A. Valuing public goods using happiness data: The case of air quality. *J Public Econ*. 2012;96(9):869–80. doi: 10.1016/j.jpubeco.2012.06.007.
70. Hobfoll SE. Conservation of resources: A new attempt at conceptualizing stress. *Am Psychol*. 1989;44(3):513–24. doi: 10.1037/0003-066X.44.3.513.
71. MacNeill AJ, McGain F, Sherman JD. Planetary health care: a framework for sustainable health systems. *Lancet Planet Health*. 2021;5(2):e66–8. doi: 10.1016/S2542-5196(21)00005-X.
72. Bressane A, da Cunha Pinto JP, Garcia Goulart AP, César de Castro Medeiros L. Which dimensions of nature contact in Urban Green Spaces most significantly contribute to mental wellbeing? A multidimensional analysis in Brazilian metropolitan cities. *Health Place*. 2024;89:103335. doi: 10.1016/j.healthplace.2024.103335.
73. Nutsford D, Pearson AL, Kingham S. An ecological study investigating the association between access to urban green space and mental health. *Public Health*. 2013;127(11):1005–11. doi: 10.1016/j.puhe.2013.08.016.

74. Paluš H, Parobek J, Šulek R, Lichý J, Šálka J. Understanding Sustainable Forest Management Certification in Slovakia: Forest Owners' Perception of Expectations, Benefits and Problems. *Sustainability*. 2018;10(7):2470. doi: 10.3390/su10072470.
75. Pham VT, Roongtawanreongsri S, Ho TQ, Tran PHN. Can payments for forest environmental services help improve income and attitudes toward forest conservation? Household-level evaluation in the Central Highlands of Vietnam. *For Policy Econ*. 2021;132:102578. doi: 10.1016/j.forpol.2021.102578.
76. Cole HVS, Lamarca MG, Connolly JJT, Anguelovski I. Are green cities healthy and equitable? Unpacking the relationship between health, green space and gentrification. *J Epidemiol Community Health*. 2017;71(11):1118–21. doi: 10.1136/jech-2017-209201.
77. Li L, Wang H, Guo C, Ning J. Can low-carbon cities increase urban housing prices? Evidence from China's low-carbon city pilot. *Environ Sci Pollut Res*. 2024;31(34):46703–21. doi: 10.1007/s11356-023-30203-6.
78. Wei X, Zhao C. Does Chinese Environmental Policy Affect Income Inequality? Evidence from the Central Environmental Protection Inspection. *China World Econ*. 2024;32(3):130–61. doi: 10.1111/cwe.12534.
79. Duchelle AE, de Sassi C, Jagger P, Cromberg M, Larson AM, Sunderlin WD, et al. Balancing carrots and sticks in REDD+: implications for social safeguards. *Ecology and Society*. 2017 [cited 10.04.2026];22(3). Available from: <https://www.jstor.org/stable/26270145>
80. Gurney GG, Pressey RL, Cinner JE, Pollnac R, Campbell SJ. Integrated conservation and development: evaluating a community-based marine protected area project for equality of socioeconomic impacts. *Philos Trans R Soc Lond B Biol Sci*. 2015;370(1681):20140277. doi: 10.1098/rstb.2014.0277.
81. De Los Reyes A, Augenstein TM, Wang M, Thomas SA, Drabick DAG, Burgers DE, et al. The validity of the multi-informant approach to assessing child and adolescent mental health. *Psychol Bull*. 2015;141(4):858–900. doi: 10.1037/a0038498.
82. Groves RM, Fowler FJ Jr, Couper MP, Lepkowski JM, Singer E, Tourangeau R. *Survey Methodology*. John Wiley & Sons; 2011. 487 p.
83. Gros C, Pronti A, Sheikh K, Hassan A, Shahjahan M. Effects of anticipatory humanitarian cash assistance to households forecasted to experience extreme flooding: evidence from Bangladesh. *Hydrol Res*. 2023;54(11):1315–28. doi: 10.2166/nh.2023.111.
84. Lei Y, Finlayson CM, Thwaites R, Shi G, Cui L. Using Government Resettlement Projects as a Sustainable Adaptation Strategy for Climate Change. *Sustainability*. 2017;9(8):1373. doi: 10.3390/su9081373.
85. McMichael C, Powell T. Planned Relocation and Health: A Case Study from Fiji. *Int J Environ Res Public Health*. 2021;18(8):4355. doi: 10.3390/ijerph18084355.
86. Nguyen Dang HA, Legg R, Khan A, Wilkinson S, Ibbett N, Doan AT. Users' Perceptions of the Contribution of a University Green Roof to Sustain Dev. *Sustainability*. 2023;15(8):6772. doi: 10.3390/su15086772.
87. Haines A, McMichael AJ, Smith KR, Roberts I, Woodcock J, Markandya A, et al. Public health benefits of strategies to reduce greenhouse-gas emissions: overview and implications for policy makers. *Lancet*. 2009;374(9707):2104–14. doi: 10.1016/S0140-6736(09)61759-1.
88. Tebaldi C, Friedlingstein P. Delayed detection of climate mitigation benefits due to climate inertia and variability. *Proc Natl Acad Sci U S A*. 2013;110(43):17229–34. doi: 10.1073/pnas.1300005110.

89. Bax V, van de Lageweg WI, Terpstra T, Buijs JM, de Reus K, de Groot F, et al. The impact of coastal realignment on the availability of ecosystem services: gains, losses and trade-offs from a local community perspective. *J Environ Manage.* 2023;345:118675. doi: 10.1016/j.jenvman.2023.118675.
90. Baninla Y, Sharifi A, Allam Z, Tume SJP, Gangtar NN, George N. An overview of climate change adaptation and mitigation research in Africa. *Front Clim.* 2022;4. doi: 10.3389/fclim.2022.976427.
91. Saber M, Koroma A, Posite RV, LY A, Tejan Bah AM, Abdulai A, et al. A comprehensive review of climate change adaptation and disaster risk reduction in Africa. *J Water Clim Change.* 2025;16(5):1831–62. doi: 10.2166/wcc.2025.741.
92. Tesfaye A, Radeny M, Ogada MJ, Recha JW, Ambaw G, Chanana N, et al. Gender empowerment and parity in East Africa: evidence from climate-smart agriculture in Ethiopia and Kenya. *Clim Dev.* 2023;15(9):768–78. doi: 10.1080/17565529.2022.2154124.
93. Radeny M, Rao EJO, Ogada MJ, Recha JW, Solomon D. Impacts of climate-smart crop varieties and livestock breeds on the food security of smallholder farmers in Kenya. *Food Secur.* 2022;14(6):1511–35. doi: 10.1007/s12571-022-01307-7.
94. Chapman R, Howden-Chapman P, Viggers H, O’Dea D, Kennedy M. Retrofitting houses with insulation: a cost–benefit analysis of a randomised community trial. *J Epidemiol Community Health.* 2009;63(4):271–7. doi: 10.1136/jech.2007.070037.
95. Baeza F, Vives Vergara A, González F, Orlando L, Valdebenito R, Cortinez-O’Ryan A, et al. The Regeneración Urbana, Calidad de Vida y Salud - RUCAS project: a Chilean multi-methods study to evaluate the impact of urban regeneration on resident health and wellbeing. *BMC Public Health.* 2021;21(1):728. doi: 10.1186/s12889-021-10739-3.

1 Tables

2

3

1 **Table 1:** Overview of All Included Studies Based on PICOS

Study	Population (P)	Intervention / Exposure (I)	Comparator (C)	Psychosocial Outcomes (O) ¹	Study Design (S)	Moderators	Results	Major limitations
Ahrentzen et al. (2016) [52]	Older adults, lower income, convenience sample (USA), n = 57	Energy retrofit housing	Pre/post	General health/quality of life, emotional distress, sleep	Quantitative, longitudinal, self-report applied as interview	-	Reduction in excessive temperature associated with improved quality of life/health, reduced emotional distress, and increased sleep duration (all $p < .05$).	Single-case study, assessment via single-item surveys
Bates et al. (2018) [50]	Children, self-selected by schools, reports by caregivers and teachers (USA), observations T1: n = 3,345, T2: n = 3,710; participants T1: n = 64/33, T3: n = 61/40	Green schoolyard renovation	Pre/post, retrospective	Social interactions, teasing/bullying	Mixed methods, longitudinal, behavioural observation and self-report	Age, gender	Intervention associated with increased social interaction and more positive interactions; bullying perceived as reduced (not significant). Interaction patterns differed by ethnicity and age ($p < .01$), with no gender effects.	Retrospective assessments of changes in student safety etc.
Bohnert et al. (2022) [58]	Children, self-selected by schools, reports by caregivers and teachers (USA), observations T1: n = 1,713, T2: n = 1,937, participants: T1: n = 94/66/27; T2: n = 84/45/25	Green schoolyard renovation	Control schools / pre-post	Social interactions, social cohesion	Mixed methods, longitudinal, behavioural observation and self-report	School context	Intervention associated with fewer interactions overall but higher proportion of positive interactions and reduced negative/neutral interactions;	No report if T1 and T2 respondents were identical

Breyse et al. (2015) [53]	Older adults, lower income, convenience sample (USA), n = 40	Green renovation/building renovation	Pre/post	General mental health	Quantitative, longitudinal, self-report applied as interview	-	improved neighbourhood climate ($p < .05$), reduced bullying and stronger school–community ties ($p < .01$); heterogeneous effects across schools, some outcomes non-significant. Intervention showed no significant within-group changes, but significant improvements relative to control (which declined) for selected outcomes ($p < .05$); no effects for other measures.	home conditions of control group unknown, enrolling 13 new tenants at T2, assessment via single-item
Cinderby & Bagwell (2017) [57]	Business community, convenience sample (UK), T1: n = 28, T2: n = 32	Green urban infrastructure	Exposure vs. non-exposure	Enjoyment, well-being, happiness	Qualitative, others-report (manager), self-report, interview	-	Green infrastructure linked to higher reported happiness ($p = .05$), improved morale, social interaction, and environmental awareness; overall well-being ratings slightly decreased (not tested), with	Change scores calculated based on different respondent; other massive development projects conducted parallel, some assessments via single-item

Gill and Simeoni (1995) [55]	Community residents, convenience sample (Australia), n = 55	Environmental enhancement project including urban park development	Pre/post exposure	Communal congeniality	Observational, cross-sectional, interview and focus group discussions	-	stable social support. Participants reported enhanced community interaction, greater use of shared social spaces, and improved intercultural exchange following the park intervention.	Representativeness of data
Hernández & Phillips (2015) [59]	Low-income households and landlords, convenience sample (USA), n = 20	Energy efficiency improvements of low-income housing units	Pre/post intervention	Stress, quality of relationships, comfort, positive emotions	Mixed methods, cross-sectional, interview	-	Intervention reduced energy-related stress and financial strain, increased well-being, and fostered positive feelings and improved tenant–landlord relationships.	Interviews conducted with household heads instead of single members; discrepancies between perspective of building owners and tenants
Hoffman (2020) [48]	Incarcerated individuals and community members, self-selection (USA), n = 17	Green space orchard project in correctional setting	No formal control; post-project survey associations	Community connectedness	Mixed methods, quasi-experimental, cross-sectional, self-report	-	Strong correlations between perceived importance of volunteering and community connectedness, and between continued engagement and prosocial impact ($r \approx .91-.93$, $p < 0.001$); qualitative data suggests	small sample; self-selection, no distinction between groups (inmates and community members), self-developed survey

Howden-Chapman et al. (2007) [54]	Low-income households with respiratory vulnerability, convenience sample (New Zealand), experimental: n = 960, control: n = 944	Home insulation retrofit	Randomised control households	General health, social functioning, emotional role functioning happiness	Quantitative, longitudinal, cluster-randomised, self-report	Analyses controlled for sex, ethnic origin, age group	therapeutic benefits and enhanced physical, mental, and social well-being. Intervention led to significant improvements in social and emotional functioning ($p < .0001$), reduced odds of poor mental health ($OR \approx 0.56$), and lower likelihood of reporting poor/fair health ($OR \approx 0.50$).	Potential argument that household income rather than substandard housing is fundamental problem underlying health issues
Hunter et al. (2021) [51]	Urban adult residents, random selection (UK), T1: n = 1,037, T2: n = 968	Urban greenway regeneration	Control areas and distance bands	Quality of life, mental well-being, social networks	Quantitative, longitudinal, self-report as interview	Analyses controlled for age, season, education, car ownership and deprivation	Significant increase in quality of life compared to declining control group ($p < .001$), no significant change in mental well-being, and small but significant decrease in social networks (no difference compared to control).	Comparison group was originally intended but not conducted
Kim et al. (2018) [60]	Urban residents / stakeholders (South Korea), n = 218	Rooftop farm vs. roof garden on university building	Comparison across roof types	Emotions and well-being	Quantitative, cross-sectional, self-report applied as interview	Individual preferences for intervention	Majority perceived green roofs as needed (> 80 %), with a third reporting restorative use; preference for	no pre-test, no control group

Marshall et al. (2023) [65]	University/community respondents and urban biodiversity setting, random sampling (UK), n = 278	Urban wildflower meadow replacing lawn	Remaining lawn and pre-sowing baseline	Mental well-being	Mixed methods, cross-sectional, self-report applied as interview	-	green roofs associated with higher motivation and satisfaction. Significantly higher mental well-being reported for meadows vs. lawns (83.5% vs. 45%, $p < .001$), with qualitative data highlighting strong restorative and emotional benefits.	Primary focus on biology and climate aspects, low survey quality
Pérez et al. (2025) [43]	Urban residents and workers (Barcelona, Spain), focus groups: n $\approx$ 20, quantitative study: final n not reported	Superblocks urban redesign	Pre/post, affected vs. control areas, mixed methods	General health, well-being, mental health, social support	Mixed-methods natural experiment, longitudinal and cross-sectional, self-report applied as interview, observational measures, discussion	-	Participants reported improved well-being ($\sim$ 45 – 55 %), with qualitative evidence indicating enhanced aesthetics, reduced pollution and noise, increased opportunities for physical activity and outdoor use, and stronger social interaction and neighbourhood cohesion.	Assessment in parallel to COVID-19 pandemic
Richerzhagen et al. (2019) [61]	Rural and peri-urban communities, convenience sampling (Colombia), n = 71	Ecosystem-based adaptation projects	Project vs. nearby non-project experience /	Perceived social benefits	Qualitative, cross-sectional, self-report applied as individual and group interview, focus groups, observation	Gender	Social benefits outweighed costs, with reported improvements in	interviewees might be biased by institutional affiliations and social standing, no

			qualitative comparison					community cohesion, empowerment, and perceived appreciation; gender differences observed, with stronger empowerment effects among women.	confidentiality during some interviews, no pre- test, no control group
Song, Li et al. (2019) [66]	Urban residents in Chinese air-pollution transmission channel cities, random sampling (China), n = 1,016	Specific air pollution control measures	Policy exposure variation	Subjective well-being, happiness, life satisfaction	Quantitative, cross- sectional, self-report	Income, education, age		Policies targeting air pollution (e.g. heating regulation, EV subsidies) linked to increased well-being; results consistent in robustness analyses.	assessment of well- being and life- satisfaction with only 1 item each
Song, Guo et al. (2019) [56]	General population (China), sample size not reported	Environmental regulations (command-and- control, market- based, informal)	Cross-region comparison	Subjective well-being	Quantitative, cross- sectional, self-report	National region, time- lag		Command-and- control and market-based regulations showed region- specific, non- linear or null associations with well-being; effects varied depending on temporal lag to intervention.	No sample size reported, assessment of subjective wellbeing with only 1 item
Tsagkari et al. (2022) [63]	Island communities, random sampling (Greece and Spain), n = 50/145	Local renewable energy projects including wind, photovoltaic, hydro-wind power	Cross-case / post- implementation perceptions	Social cohesion	Quantitative, cross- sectional, self-report	Perception on success of the project		Participants reported overall agreement with social benefits; institutional benefits were perceived as stronger, while	No pre-test, no control group

Wang & Tang (2023) [67]	Representative data, conducted annually (China), sample size not reported	Environmental protection tax and subjective air pollution context	Cross-regional variation	Subjective well-being	Quantitative, cross-sectional, self-report	Income, region, urban-rural, gender, age	economic benefits were rated lower than social benefits. Subjective air pollution significantly decreased well-being, while environmental protection tax increased happiness; negative effects were larger for low-income individuals and present in both urban and rural populations.	Other relevant health factors such as work status or education might be responsible for effects
Wang et al. (2024) [49]	Middle-aged and older adults (China), n ≈ 61,100	Environmental protection tax policy	Treatment vs. non-treatment cities over time	Sleep quality, memory condition	Quantitative, longitudinal, self-report	Age, marital status	Significant increase in sleep duration ($p < .05$) and marginal improvement in memory function ($p < .10$) following the intervention.	Assessments using only 1 item per construct, environmental protection tax policy involves multiple factors (tax rate settings, collection methods, and the implementation strength of local governments) reliance on observational data; difficulty measuring morbidity impacts
Xie et al. (2024) [62]	General population / households, representative survey (China), n = not reported	Large-scale air pollution control policy	Policy boundary and non-policy regions	Happiness, self-rated health	Quasi-experimental, quantitative, longitudinal, observational	Regional location	Air quality improvements linked to increased happiness ($\beta \approx -0.017$, $p < .01$) and improved perceived health ($\beta \approx 0.007$, $p < .01$).	

Zhao & Li (2024) [64]	Working-age adults, random sampling (China), observations in different subsamples n = 9,332-30,918	Atmosphere Ten Articles Plan (ATAP) to target pollution	Staggered city implementation / pre-post	Subjective well-being	Quantitative, longitudinal, self-report applied as interview	Emotional stability, mental health	ATAP significantly increased happiness and likelihood of being very happy ($p < .001$), with effects mediated by improved mental health, emotional stability, and reduced insomnia; benefits were greater for low- income, urban, and middle- aged/older individuals, with no effects for rural or younger groups.	assessment of subjective well- being with only 1 item
--------------------------	--	--	--	--------------------------	--	--	--	--

-
- 1
 - 2 ¹limited to relevant psychosocial co-benefits
 - 3 PICOS = Population, Intervention/Exposure, Comparator, Outcome, Study design
 - 4
 - 5

1 **Table 2:** Quality appraisal of included studies based on QuADS criteria

2

Criterion	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.	13.
	Theoretical or conceptual underpinning to the research	Statement of research aim/s	Clear description of research setting and target population	Appropriate study design to address the stated research aim/s	Appropriate sampling to address the research aim/s	Rationale for choice of data collection tool/s	Format & content of data collection tool is appropriate to address the stated research aim/s	Description of data collection procedure	Recruitment data provided	Justification for analytic method selected	The method of analysis was appropriate to answer the research aim/s	Evidence that the research stakeholders have been considered in research design or conduct	Strengths and limitations critically discussed
Ahrentzen et al. (2016) [52]	2	3	3	2	2	2	3	3	2	2	3	1	3
Bates et al. (2018) [50]	2	3	3	2	2	2	2	2	2	2	3	1	2
Bohnert et al. (2022) [58]	2	3	3	3	2	2	3	3	2	2	3	1	2
Breyse et al. (2015) [53]	1	3	3	1	2	1	3	3	2	1	2	0	3
Cinderby & Bagwell (2017) [57]	3	3	3	3	2	2	3	3	3	2	3	2	3
Gill and Simeoni (1995) [55]	3	3	3	3	2	2	3	3	1	1	2	2	1
Hernández & Phillips (2015) [59]	1	3	3	2	2	2	2	3	2	2	3	2	3
Hoffman (2020) [48]	1	3	2	1	1	1	1	2	1	1	1	1	0
Howden-Chapman et al. (2007) [54]	2	3	3	3	3	2	3	3	3	3	3	1	3
Hunter et al. (2021) [51]	3	3	3	3	3	3	3	3	3	2	3	2	3
Kim et al. (2018) [60]	3	3	3	2	1	0	2	1	0	0	3	0	1

Marshall et al. (2023) [65]	2	3	2	3	1	1	1	1	1	0	3	0	0
Pérez et al. (2025) [43] quantitative	3	3	3	2	1	3	2	2	1	2	3	2	1
Pérez et al. (2025) [43] qualitative	3	3	3	3	2	2	3	3	2	1	3	2	2
Richerzhagen et al. (2019) [61]	3	3	3	2	1	2	1	2	1	0	1	1	1
Song, Li et al. (2019) [66]	3	3	3	3	3	2	1	1	1	3	3	0	1
Song, Guo et al. (2019) [56]	3	2	2	3	2	3	1	3	3	3	3	3	1
Tsagkari et al. (2022) [63]	3	3	3	2	3	3	2	2	1	0	2	1	2
Wang & Tang (2023) [67]	2	3	2	2	1	1	2	1	1	2	2	0	2
Wang et al. (2024) [49]	3	3	3	3	2	1	1	1	1	2	3	0	2
Xie et al. (2024) [62]	3	3	3	3	3	2	3	3	2	3	3	0	3
Zhao & Li (2024) [64]	3	3	3	3	2	3	2	2	1	3	3	1	3

1

2

3 Pérez et al., 2025: rated separately due to distinct subprojects with different methodologies: quantitative components (survey) and qualitative components (focus groups, ethnographic studies)

4 QuADS = Quality assessment with diverse studies; the QuADS includes 13 criteria, each rated from 0 to 3. A score of 0 generally indicates that the criterion was not mentioned, whereas a score of 3 indicates that the criterion was explicitly described, considered, or addressed. Higher total scores indicate more complete reporting and/or stronger methodological appropriateness. Cell shading corresponds to the respective QuADS rating and is provided to facilitate visual interpretation across studies and criteria.

8